



A global method for deterministic and stochastic homogenisation in BV

Filippo Cagnetti¹ · Gianni Dal Maso² · Lucia Scardia³ · Caterina Ida Zeppieri⁴

Received: 3 May 2021 / Accepted: 4 March 2022 / Published online: 7 April 2022
© The Author(s) 2022

Abstract

In this paper we study the deterministic and stochastic homogenisation of free-discontinuity functionals under *linear* growth and coercivity conditions. The main novelty of our deterministic result is that we work under very general assumptions on the integrands which, in particular, are not required to be periodic in the space variable. Combining this result with the pointwise Subadditive Ergodic Theorem by Akcoglu and Krengel, we prove a stochastic homogenisation result, in the case of stationary random integrands. In particular, we characterise the limit integrands in terms of asymptotic cell formulas, as in the classical case of periodic homogenisation.

Keywords Free-discontinuity problems · Γ -convergence · Stochastic homogenisation · Blow-up method

Mathematics Subject Classification 49J45 · 49Q20 · 74Q05 · 74E30 · 60K35

✉ Filippo Cagnetti
F.Cagnetti@sussex.ac.uk

Gianni Dal Maso
dalmaso@sissa.it

Lucia Scardia
L.Scardia@hw.ac.uk

Caterina Ida Zeppieri
caterina.zeppieri@uni-muenster.de

¹ Department of Mathematics, University of Sussex, Brighton, UK

² SISSA, Trieste, Italy

³ Department of Mathematics, Heriot-Watt University, Edinburgh, UK

⁴ Angewandte Mathematik, WWU Münster, Münster, Germany

1 Introduction

In this paper we derive deterministic and stochastic homogenisation results for *free-discontinuity functionals* in the space of functions of bounded variation.

We consider families of free-discontinuity functionals of the form

$$E_\varepsilon(\omega)(u, A) = \int_A f\left(\omega, \frac{x}{\varepsilon}, \nabla u\right) dx + \int_{S_u \cap A} g\left(\omega, \frac{x}{\varepsilon}, [u], \nu_u\right) d\mathcal{H}^{n-1}, \quad (1.1)$$

where ω belongs to the sample space of a given probability space (Ω, \mathcal{T}, P) and labels the realisations of the random integrands f and g , while $\varepsilon > 0$ is a parameter of either geometrical or physical nature, and sets the scale of the problem. In (1.1) the set A belongs to the class \mathcal{A} of bounded open subsets of \mathbb{R}^n and the function u belongs to the space $SBV(A, \mathbb{R}^m)$ of special \mathbb{R}^m -valued functions of bounded variation on A (see [21] and [5, Section 4.5]). Moreover, ∇u denotes the approximate gradient of u , $[u]$ stands for the difference $u^+ - u^-$ between the approximate limits of u on both sides of the discontinuity set S_u , ν_u denotes the (generalised) normal to S_u , and \mathcal{H}^{n-1} is the $(n - 1)$ -dimensional Hausdorff measure in \mathbb{R}^n .

Functionals as in (1.1) are commonly used in applications where the physical quantity described by u can exhibit discontinuities, e.g. in variational models of fracture mechanics, in the theory of computer vision and image segmentation, and in problems involving phase transformations.

We are interested in determining the *almost sure* limit behaviour of E_ε as $\varepsilon \rightarrow 0+$, when f and g satisfy *linear* growth and coercivity conditions in the gradient and in the jump, respectively. The linear growth of the volume energy sets the limit problem naturally in the space BV of functions of bounded variation. Indeed, in this setting, limits of sequences of displacements with bounded energy can develop a Cantor component in the distributional gradient.

This is in contrast with our previous work [16] (in the deterministic case) and [17] (in the stochastic case), where, under the assumption of superlinear growth for f , the limit problem was naturally set in the space SBV of special functions with bounded variation.

The two main results of this paper are a deterministic homogenisation result for functionals of the type (1.1), when ω is fixed and f and g are not necessarily periodic, and a stochastic homogenisation result, obtained for P -a.e. $\omega \in \Omega$, under the additional assumption of *stationarity* of f and g .

1.1 The deterministic result

For the deterministic result we consider ω as fixed in (1.1) and write $E_\varepsilon(u, A)$ instead of $E_\varepsilon(\omega)(u, A)$.

We study the limit behaviour of the functionals $E_\varepsilon(\cdot, A)$, for every $A \in \mathcal{A}$, as $\varepsilon \rightarrow 0+$, under the assumption that the energy densities f and g belong to suitable classes \mathcal{F} and \mathcal{G} of admissible volume and surface densities (see Definition 3.1). As announced above, a key requirement for the class \mathcal{F} is that f satisfies *linear* upper

and lower bounds in the gradient variable. Additionally, we require that the recession function f^∞ of f is defined at every point. For the class \mathcal{G} , we require that g is bounded from below and from above by the amplitude of the jump, and that its directional derivative g_0 in the jump variable at $[u] = 0$ exists and is finite. The functions f^∞ and g_0 will play an important role in determining the limiting densities.

We stress here that we *do not* require any periodicity in the spatial variable x for the volume and surface densities; moreover, we *do not* require any continuity in the spatial variable either, since it would be unnatural for applications.

Under these general assumptions, using the so-called localisation method of Γ -convergence [18], we can prove that there exists a subsequence (ε_k) such that, for every $A \in \mathcal{A}$, $(E_{\varepsilon_k}(\cdot, A))$ Γ -converges to an abstract functional $\widehat{E}(\cdot, A)$, that $\widehat{E}(\cdot, A)$ is finite only in BV , and that, for every $u \in BV_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, the set function $\widehat{E}(u, \cdot)$ is the restriction to \mathcal{A} of a Borel measure (see Theorem 5.1).

Note that, without any additional assumptions, one cannot expect that $\widehat{E}(\cdot, A)$ can be written in an integral form. In particular, since there is no guarantee that $z \mapsto \widehat{E}(u(\cdot - z), A + z)$ is continuous (which is instead automatically satisfied in the periodic case), we cannot directly apply the integral representation result in BV [10]. Our integral representation result is hence obtained under some additional assumptions, which are though more general than periodicity. We require that the limits of some rescaled minimisation problems, defined in terms of f, g, f^∞ , and g_0 , exist and are independent of the spatial variable. These limits will then define the densities of \widehat{E} .

More precisely, for $A \in \mathcal{A}, w \in SBV(A, \mathbb{R}^m), f \in \mathcal{F}$, and $g \in \mathcal{G}$ we set

$$m^{f,g}(w, A) := \inf \left\{ \int_A f(x, \nabla u) dx + \int_{S_u \cap A} g(x, [u], \nu_u) d\mathcal{H}^{n-1} : u \in SBV(A, \mathbb{R}^m), u = w \text{ near } \partial A \right\}, \tag{1.2}$$

and we *assume* that for every $\xi \in \mathbb{R}^{m \times n}$ the limit

$$\lim_{r \rightarrow +\infty} \frac{m^{f,g_0}(\ell_\xi, Q_r(rx))}{r^n} =: f_{\text{hom}}(\xi) \tag{1.3}$$

exists and is independent of $x \in \mathbb{R}^n$, and that for every $\zeta \in \mathbb{R}^m$ and $\nu \in \mathbb{S}^{n-1}$ the limit

$$\lim_{r \rightarrow +\infty} \frac{m^{f^\infty,g}(u_{rx,\zeta,\nu}, Q_r^\nu(rx))}{r^{n-1}} =: g_{\text{hom}}(\zeta, \nu) \tag{1.4}$$

exists and is independent of $x \in \mathbb{R}^n$. In (1.3), ℓ_ξ denotes the linear function with gradient ξ ; in (1.4), $Q_r^\nu(rx) := R_\nu \left((-\frac{r}{2}, \frac{r}{2})^n \right) + rx$, where R_ν is an orthogonal $n \times n$ matrix such that $R_\nu e_n = \nu$, and

$$u_{rx,\zeta,\nu}(y) := \begin{cases} \zeta & \text{if } (y - rx) \cdot \nu \geq 0, \\ 0 & \text{if } (y - rx) \cdot \nu < 0. \end{cases}$$

The limits (1.3) and (1.4) are the counterpart of the asymptotic cell-formulas in the classical periodic homogenisation [10]. In that case, periodicity in the spatial variable x ensures the existence of the limits and their homogeneity in x , that here we have to postulate. Our assumptions are however weaker than periodicity: notably, they are fulfilled in the case of *stationary* integrands, as we show in the present work.

In line with the result in [10], the functional being minimised in (1.3) (respectively in (1.4)) has densities f and g_0 (respectively f^∞ and g) rather than f and g . We note that, if the density f satisfied a superlinear growth, then $f^\infty(\cdot, \xi) = +\infty$ for $\xi \neq 0$. Since one always has $f^\infty(\cdot, 0) = 0$, in the superlinear case there would be the following changes in (1.4): on the one hand the minimisation would be done over functions with $\nabla u = 0$; on the other hand the functional to be minimised would reduce to just the surface term. This is indeed the situation in [16], where we assume a superlinear growth for f . Moreover, in [16] we work under different growth conditions for g as well, which in particular satisfies $g \geq c$, for a given fixed positive constant. In that case $g_0 \equiv +\infty$ (see (3.3)), and hence formally the minimisation in (1.3) would be over Sobolev functions, and the functional to be minimised would reduce to just the bulk term. Again, this is exactly what happens in [16] (see also [11, 28]).

In Lemmas 4.2 and 4.5 we show that $f_{\text{hom}} \in \mathcal{F}$ and $g_{\text{hom}} \in \mathcal{G}$; the fact that $f_{\text{hom}} \in \mathcal{F}$ guarantees in particular the existence of its recession function f_{hom}^∞ . In Propositions 6.2, 7.2, and 8.3, we show that the functions f_{hom} , g_{hom} and f_{hom}^∞ are the densities of the volume, surface, and Cantor terms of \widehat{E} , respectively. To do so we use the blow-up technique of Fonseca and Müller [27] (see also [12]), extended to the *BV*-setting by Bouchitté, Fonseca, and Mascarenhas [10]. More precisely, thanks to (1.3) and (1.4), we prove that the following identities hold true for every $A \in \mathcal{A}$ and for every $u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ with $u|_A \in BV(A, \mathbb{R}^m)$:

$$\begin{aligned} \frac{d\widehat{E}(u, \cdot)}{d\mathcal{L}^n}(x) &= f_{\text{hom}}(\nabla u(x)) \quad \text{for } \mathcal{L}^n\text{-a.e. } x \in A, \\ \frac{d\widehat{E}(u, \cdot)}{d\mathcal{H}^{n-1} \llcorner S_u}(x) &= g_{\text{hom}}([u](x), \nu_u(x)) \quad \text{for } \mathcal{H}^{n-1}\text{-a.e. } x \in S_u \cap A, \end{aligned}$$

and

$$\frac{d\widehat{E}(u, \cdot)}{d|C(u)|}(x) = f_{\text{hom}}^\infty\left(\frac{dC(u)}{d|C(u)|}(x)\right) \quad \text{for } |C(u)|\text{-a.e. } x \in A,$$

where \mathcal{L}^n is the Lebesgue measure in \mathbb{R}^n , $\mathcal{H}^{n-1} \llcorner S_u$ is the measure defined by $(\mathcal{H}^{n-1} \llcorner S_u)(B) := \mathcal{H}^{n-1}(S_u \cap B)$ for every Borel set $B \subset \mathbb{R}^n$, and $C(u)$ is the Cantor part of the distributional derivative of u .

In particular, since the right-hand sides of the previous equalities do not depend on (ε_k) , the homogenisation result holds true for the whole sequence (E_ε) . Moreover, the fact that $f_{\text{hom}} \in \mathcal{F}$ and $g_{\text{hom}} \in \mathcal{G}$ implies that the classes \mathcal{F} and \mathcal{G} are closed under homogenisation, and that on *SBV* the functionals E_ε and their Γ -limit \widehat{E} are free-discontinuity functionals of the same type.

As observed before, f_{hom} and g_{hom} depend on both the volume and the surface densities of the functional E_ε . Indeed, the minimisation problems in (1.3) and (1.4)

involve both f (or f^∞) and g (or g_0). In other words, volume and surface term *do interact* in the limit, which is a typical feature of the linear-growth setting. This is in contrast with the case of superlinear growth considered in [16], in which the limit volume density only depends on the volume density of E_ε , and similarly the limit surface density only depends on the surface density of E_ε . The volume-surface decoupling is typical of the *SBV*-setting in presence of superlinear growth conditions on f [11, 16, 28].

Note however that, even in the superlinear case, if f and g satisfy “degenerate” coercivity conditions, due for instance to the presence of perforations or “weak” inclusions in the domain, the situation is more involved. Indeed, while in [7, 14, 15, 25, 33] the volume and surface terms do not interact in the homogenised limit, in [6, 8, 20, 32, 34, 35] they do interact and produce rather complex limit effects.

1.2 The stochastic result

In Section 9 we prove the almost sure Γ -convergence of the sequence of random functionals $E_\varepsilon(\omega)$ in (1.1) to a random homogenised integral functional, under the assumption that the volume and surface integrands f and g are stationary (see Definition 3.12 and Remark 3.16).

In the random setting stationarity is the natural counterpart of periodicity, since it implies that f and g are “statistically” translation-invariant, or “periodic in law”.

The application of the deterministic result Theorem 4.1, at ω fixed, ensures that $E_\varepsilon(\omega)$ Γ -converges to the free-discontinuity functional

$$E_{\text{hom}}(\omega)(u, A) := \int_A f_{\text{hom}}(\omega, \nabla u) \, dx + \int_{S_u \cap A} g_{\text{hom}}(\omega, [u], \nu_u) \, d\mathcal{H}^{n-1} + \int_A f_{\text{hom}}^\infty\left(\omega, \frac{dC(u)}{d|C(u)|}\right) d|C(u)|,$$

with

$$f_{\text{hom}}(\omega, \xi) := \lim_{r \rightarrow +\infty} \frac{m^{f(\omega), g_0(\omega)}(\ell_\xi, Q_r(rx))}{r^n}, \tag{1.5}$$

for every $\xi \in \mathbb{R}^{m \times n}$, and

$$g_{\text{hom}}(\omega, \zeta, \nu) := \lim_{r \rightarrow +\infty} \frac{m^{f^\infty(\omega), g(\omega)}(u_{rx, \zeta, \nu}, Q_r^v(rx))}{r^{n-1}}, \tag{1.6}$$

for every $\zeta \in \mathbb{R}^m$ and $\nu \in \mathbb{S}^{n-1}$, provided the limits in (1.5) and (1.6) (which are the same as (1.3) and (1.4), modulo the additional dependence on the random parameter ω) exist and are independent of $x \in \mathbb{R}^n$. Therefore, to show that the Γ -convergence of $E_\varepsilon(\omega)$ towards $E_{\text{hom}}(\omega)$ actually holds true for P -a.e. $\omega \in \Omega$ it is necessary to show that the limits in (1.5) and (1.6) exist and are independent of $x \in \mathbb{R}^n$ for P -almost every realisation $\omega \in \Omega$. To do so, we follow the general strategy firstly introduced

in [19] in the Sobolev setting, and then extended to the SBV -setting in [17] (see also [3]).

This strategy relies on the Subadditive Ergodic Theorem by Akcoglu and Krengel [1] (see Theorem 3.15) and requires, among other things, to show that the minimisation problems in (1.5) and (1.6) define two *subadditive stochastic processes* (see Definition 3.13).

This task however poses a challenge even at the very first step: proving that $\omega \mapsto m^{f(\omega), g_0(\omega)}$ and $\omega \mapsto m^{f^\infty(\omega), g(\omega)}$ are *measurable*. Indeed, while both $m^{f(\omega), g_0(\omega)}$ and $m^{f^\infty(\omega), g(\omega)}$, by (1.2), involve the minimisation of measurable functionals in the random variable ω , such minimisation is performed over the space SBV , which is not separable. Since the infimum in (1.2) cannot be reduced to a countable set, the measurability of $\omega \mapsto m^{f(\omega), g_0(\omega)}$ and $\omega \mapsto m^{f^\infty(\omega), g(\omega)}$ cannot be inferred directly from the measurability in ω of f , f^∞ , g and g_0 (see the Appendix). Let us also observe that the situation here is substantially different from that treated in [16, 17], for a number of reasons. Indeed in [16, 17], as observed before, due to the different assumptions on f and g , the Γ -limit exhibits a “separation” of the volume and surface term. In particular, the limit volume density is obtained as the limit of some minimisation problems similar to (1.2), but where the minimisation is done over the space of Sobolev functions. Hence in that case the limit volume density is ω -measurable, due to the separability of the space (see also [19]). On the other hand, the measurability of the minimisation problems defining the limit surface density was delicate also in [17], since the minimum was taken over Caccioppoli partitions. However, in [17] the minimisation involved only the surface term of the functional, which makes the proof much simpler than the one required now.

Once the measurability in ω is established (see Proposition A.12), we have to face yet another difficulty: determining the *dimension* of the stochastic processes. Indeed, using the competitors ℓ_ξ and $u_{r,x,\zeta,v}$ in the minimisation problems $m^{f(\omega), g_0(\omega)}$ and $m^{f^\infty(\omega), g(\omega)}$, respectively, suggests the rescalings in (1.5) and (1.6). Hence it suggests that $m^{f(\omega), g_0(\omega)}$ should define an n -dimensional process, while $m^{f^\infty(\omega), g(\omega)}$ should define an $(n - 1)$ -dimensional process. On the other hand, in both cases the functionals appearing in the minimisation problems, if seen as set functions, are defined on n -dimensional sets.

To solve this problem we proceed as in [3, 13] and [17], where similar issues arise in the study of pure surface energies of spin systems, and in the case of free-discontinuity problems with superlinear growth, respectively.

Finally, the last difficulty consists in showing that, as in [3, 17], the limits in (1.5) and (1.6) do not depend on x . This is particularly delicate for (1.6), due to the presence of an x -dependent boundary condition.

We conclude by observing that our analysis also shows that, if f and g are ergodic, then the homogenised integrands f_{hom} and g_{hom} are ω -independent, and hence the limit E_{hom} is deterministic.

1.3 Outline of the paper.

This paper is organised as follows. In Section 2 we introduce some notation. Section 3 consists of two parts: in Section 3.1 we introduce the stochastic free-discontinuity functionals and recall the Ergodic Subadditive Theorem; in Section 3.2 we state the main results of the paper.

Sections 4–8 focus on the deterministic results. More in detail, in Section 4 we state the deterministic Γ -convergence results and prove some properties of the limit densities; Section 5 is devoted to the abstract Γ -convergence result; the volume, surface and Cantor terms of the abstract Γ -limit are then identified in Sections 6, 7 and 8, respectively.

Finally, Section 9 focuses on the stochastic homogenisation result, while the proof of the measurability of $\omega \mapsto m^{f(\omega),g_0(\omega)}$ and $\omega \mapsto m^{f^\infty(\omega),g(\omega)}$ is postponed to the Appendix.

2 Notation

We introduce now some notation that will be used throughout the paper.

- (a) m and n are fixed positive integers, with $n \geq 2$, \mathbb{R} is the set of real numbers, while \mathbb{Q} is the set of rational numbers. The canonical basis of \mathbb{R}^n is denoted by e_1, \dots, e_n . For $a, b \in \mathbb{R}^n$, $a \cdot b$ denotes the Euclidean scalar product between a and b , and $|\cdot|$ denotes the absolute value in \mathbb{R} or the Euclidean norm in \mathbb{R}^n , \mathbb{R}^m , or $\mathbb{R}^{m \times n}$ (the space of $m \times n$ matrices with real entries), depending on the context. If $v \in \mathbb{R}^m$ and $w \in \mathbb{R}^n$, the symbol $v \otimes w$ stands for the matrix in $\mathbb{R}^{m \times n}$ whose entries are $(v \otimes w)_{ij} = v_i w_j$, for $i = 1, \dots, m$ and $j = 1, \dots, n$.
- (b) $\mathbb{S}^{m-1} := \{\zeta = (\zeta_1, \dots, \zeta_m) \in \mathbb{R}^m : \zeta_1^2 + \dots + \zeta_m^2 = 1\}$, $\mathbb{S}^{n-1} := \{x = (x_1, \dots, x_n) \in \mathbb{R}^n : x_1^2 + \dots + x_n^2 = 1\}$, and $\widehat{\mathbb{S}}_\pm^{n-1} := \{x \in \mathbb{S}^{n-1} : \pm x_{i(x)} > 0\}$, where $i(x)$ is the largest $i \in \{1, \dots, n\}$ such that $x_i \neq 0$. Note that $\mathbb{S}^{n-1} = \widehat{\mathbb{S}}_+^{n-1} \cup \widehat{\mathbb{S}}_-^{n-1}$.
- (c) \mathcal{L}^n denotes the Lebesgue measure on \mathbb{R}^n and \mathcal{H}^{n-1} the $(n - 1)$ -dimensional Hausdorff measure on \mathbb{R}^n .
- (d) \mathcal{A} denotes the collection of all bounded open subsets of \mathbb{R}^n ; if $A, B \in \mathcal{A}$, by $A \subset\subset B$ we mean that A is relatively compact in B .
- (e) For $u \in BV(A, \mathbb{R}^m)$, with $A \in \mathcal{A}$, the jump of u across the jump set S_u is defined by $[u] := u^+ - u^-$, while ν_u denotes the (generalised) normal to S_u (see [5, Definition 3.67]).
- (f) For every $u \in BV(A, \mathbb{R}^m)$, with $A \in \mathcal{A}$, the distributional gradient, denoted by Du , is an $\mathbb{R}^{m \times n}$ -valued Radon measure on A , whose absolutely continuous part with respect to \mathcal{L}^n , denoted by $D^a u$, has a density $\nabla u \in L^1(A, \mathbb{R}^{m \times n})$ (which coincides with the approximate gradient of u), while the singular part $D^s u$ can be decomposed as $D^s u = D^j u + C(u)$, where the jump part $D^j u$ is given by

$$D^j u(B) = \int_{B \cap S_u} [u] \otimes \nu_u d\mathcal{H}^{n-1} \quad \text{for every Borel set } B \subset A,$$

and the Cantor part $C(u)$ is an $\mathbb{R}^{m \times n}$ -valued Radon measure on A which vanishes on all Borel sets $B \subset A$ with $\mathcal{H}^{n-1}(B) < +\infty$.

(g) For $x \in \mathbb{R}^n$ and $\rho > 0$ we define

$$B_\rho(x) := \{y \in \mathbb{R}^n : |y - x| < \rho\},$$

$$Q_\rho(x) := \{y \in \mathbb{R}^n : |(y - x) \cdot e_i| < \frac{\rho}{2} \text{ for } i = 1, \dots, n\}.$$

We omit the subscript ρ when $\rho = 1$.

(h) For every $v \in \mathbb{S}^{n-1}$ let R_v be an orthogonal $n \times n$ matrix such that $R_v e_n = v$; we assume that the restrictions of the function $v \mapsto R_v$ to the sets $\widehat{\mathbb{S}}_{\pm}^{n-1}$ defined in (b) are continuous and that $R_{-v} Q(0) = R_v Q(0)$ for every $v \in \mathbb{S}^{n-1}$; moreover, we assume that $R_v \in O(n) \cap \mathbb{Q}^{n \times n}$ for every $v \in \mathbb{Q}^n \cap \mathbb{S}^{n-1}$, where $O(n)$ denotes the set of orthogonal $n \times n$ matrices. A map $v \mapsto R_v$ satisfying these properties is provided in [16, Example A.1 and Remark A.2].

(i) For $x \in \mathbb{R}^n$, $\rho > 0$, and $v \in \mathbb{S}^{n-1}$ we set

$$Q_\rho^v(x) := R_v Q_\rho(0) + x.$$

For $k \in \mathbb{R}$, with $k > 0$, we also define the rectangle

$$Q_\rho^{v,k}(x) := Q_\rho^{v,k}(0) + x$$

where $Q_\rho^{v,k}(0)$ is obtained from $Q_\rho^v(0)$ by a dilation of amplitude k in the directions orthogonal to v ; *i.e.*,

$$Q_\rho^{v,k}(0) := R_v \left(\left(-\frac{k\rho}{2}, \frac{k\rho}{2} \right)^{n-1} \times \left(-\frac{\rho}{2}, \frac{\rho}{2} \right) \right).$$

We set

$$\partial_v^\perp Q_\rho^{v,k}(x) := \partial Q_\rho^{v,k}(x) \cap R_v \left(\left(-\frac{k\rho}{2}, \frac{k\rho}{2} \right)^{n-1} \times \mathbb{R} \right),$$

$$\partial_v^\parallel Q_\rho^{v,k}(x) := \partial Q_\rho^{v,k}(x) \cap R_v \left(\mathbb{R}^{n-1} \times \left(-\frac{\rho}{2}, \frac{\rho}{2} \right) \right),$$

namely the union of the faces of $Q_\rho^{v,k}(x)$ that are orthogonal and parallel to v , respectively.

(j) Let μ and λ be two Radon measures on $A \in \mathcal{A}$, with values in a finite dimensional Hilbert space X and in $[0, +\infty]$, respectively; for every $x \in A$ the Radon-Nikodym derivative of μ with respect to λ is defined as

$$\frac{d\mu}{d\lambda}(x) := \lim_{r \rightarrow 0^+} \frac{\mu(x + rC)}{\lambda(x + rC)}$$

whenever the limit exists in X and is independent of the choice of the bounded, closed set C containing the origin in its interior (see [26, Definition 1.156]); according to the Besicovich differentiation theorem $\frac{d\mu}{d\lambda}(x)$ exists for λ -a.e. $x \in A$ and

$\mu = \frac{d\mu}{d\lambda}\lambda + \mu^s$, where μ^s is the singular part of μ with respect to λ (see [26, Theorem 1.155]).

- (k) For $\xi \in \mathbb{R}^{m \times n}$, the linear function from \mathbb{R}^n to \mathbb{R}^m with gradient ξ is denoted by ℓ_ξ ; i.e., $\ell_\xi(x) := \xi x$, where x is considered as an $n \times 1$ matrix.
- (l) For $x \in \mathbb{R}^n$, $\zeta \in \mathbb{R}^m$, and $\nu \in \mathbb{S}^{n-1}$ we define the function $u_{x,\zeta,\nu}$ as

$$u_{x,\zeta,\nu}(y) := \begin{cases} \zeta & \text{if } (y - x) \cdot \nu \geq 0, \\ 0 & \text{if } (y - x) \cdot \nu < 0. \end{cases}$$

- (m) For $x \in \mathbb{R}^n$ and $\nu \in \mathbb{S}^{n-1}$, we set

$$\Pi_0^\nu := \{y \in \mathbb{R}^n : y \cdot \nu = 0\} \quad \text{and} \quad \Pi_x^\nu := \{y \in \mathbb{R}^n : (y - x) \cdot \nu = 0\}.$$

- (n) For a given topological space X , $\mathcal{B}(X)$ denotes its Borel σ -algebra. For every integer $k \geq 1$, \mathcal{B}^k is the Borel σ -algebra of \mathbb{R}^k , while \mathcal{B}_ζ^n stands for the Borel σ -algebra of \mathbb{S}^{n-1} .

3 Setting of the problem and statements of the main results

This section consists of two parts: in Section 3.1 we introduce the stochastic free-discontinuity functionals and recall the Ergodic Subadditive Theorem; in Section 3.2 we state the main results of the paper.

3.1 Setting of the problem

Throughout the paper we fix the following constants: $c_1, c_2, c_3, c_4, c_5 \in [0, +\infty)$, with $0 < c_2 \leq c_3$, and $\alpha \in (0, 1)$. Moreover, we fix two nondecreasing continuous functions $\sigma_1, \sigma_2 : [0, +\infty) \rightarrow [0, +\infty)$ such that $\sigma_1(0) = \sigma_2(0) = 0$.

Definition 3.1 (*Volume and surface integrands*) Let $\mathcal{F} = \mathcal{F}(c_1, c_2, c_3, c_4, c_5, \alpha, \sigma_1)$ be the collection of all functions $f : \mathbb{R}^n \times \mathbb{R}^{m \times n} \rightarrow [0, +\infty)$ satisfying the following conditions:

- (f1) (measurability) f is Borel measurable on $\mathbb{R}^n \times \mathbb{R}^{m \times n}$;
- (f2) (continuity in ξ) for every $x \in \mathbb{R}^n$ we have

$$|f(x, \xi_1) - f(x, \xi_2)| \leq \sigma_1(|\xi_1 - \xi_2|)(f(x, \xi_1) + f(x, \xi_2)) + c_1|\xi_1 - \xi_2|$$

for every $\xi_1, \xi_2 \in \mathbb{R}^{m \times n}$;

- (f3) (lower bound) for every $x \in \mathbb{R}^n$ and every $\xi \in \mathbb{R}^{m \times n}$

$$c_2|\xi| \leq f(x, \xi);$$

- (f4) (upper bound) for every $x \in \mathbb{R}^n$ and every $\xi \in \mathbb{R}^{m \times n}$

$$f(x, \xi) \leq c_3|\xi| + c_4;$$

(f_5) (recession function) for every $x \in \mathbb{R}^n$ and every $\xi \in \mathbb{R}^{m \times n}$ the limit

$$f^\infty(x, \xi) := \lim_{t \rightarrow +\infty} \frac{1}{t} f(x, t\xi), \tag{3.1}$$

which defines the recession function of f , exists and is finite; moreover, f^∞ satisfies the inequality

$$|f^\infty(x, \xi) - \frac{1}{t} f(x, t\xi)| \leq \frac{c_5}{t} + \frac{c_5}{t} f(x, t\xi)^{1-\alpha} \tag{3.2}$$

for every $x \in \mathbb{R}^n$, every $\xi \in \mathbb{R}^{m \times n}$, and every $t > 0$.

Let $\mathcal{G} = \mathcal{G}(c_2, c_3, \sigma_2)$ be the collection of all functions $g: \mathbb{R}^n \times \mathbb{R}^m \times \mathbb{S}^{n-1} \rightarrow [0, +\infty)$ satisfying the following conditions:

- (g1) (measurability) g is Borel measurable on $\mathbb{R}^n \times \mathbb{R}^m \times \mathbb{S}^{n-1}$;
- (g2) (continuity in ζ) for every $x \in \mathbb{R}^n$ and every $v \in \mathbb{S}^{n-1}$ we have

$$|g(x, \zeta_2, v) - g(x, \zeta_1, v)| \leq \sigma_2(|\zeta_1 - \zeta_2|)(g(x, \zeta_1, v) + g(x, \zeta_2, v))$$

for every $\zeta_1, \zeta_2 \in \mathbb{R}^m$;

- (g3) (lower bound) for every $x \in \mathbb{R}^n, \zeta \in \mathbb{R}^m$, and $v \in \mathbb{S}^{n-1}$

$$c_2|\zeta| \leq g(x, \zeta, v);$$

- (g4) (upper bound) for every $x \in \mathbb{R}^n, \zeta \in \mathbb{R}^m$, and $v \in \mathbb{S}^{n-1}$

$$g(x, \zeta, v) \leq c_3|\zeta|;$$

- (g5) (directional derivative at 0) for every $x \in \mathbb{R}^n, \zeta \in \mathbb{R}^m$, and $v \in \mathbb{S}^{n-1}$ the limit

$$g_0(x, \zeta, v) := \lim_{t \rightarrow 0^+} \frac{1}{t} g(x, t\zeta, v) \tag{3.3}$$

exists, is finite, and is uniform with respect to $x \in \mathbb{R}^n, \zeta \in \mathbb{S}^{m-1}$, and $v \in \mathbb{S}^{n-1}$.

- (g6) (symmetry) for every $x \in \mathbb{R}^n, \zeta \in \mathbb{R}^m$, and $v \in \mathbb{S}^{n-1}$

$$g(x, \zeta, v) = g(x, -\zeta, -v).$$

For every $f \in \mathcal{F}, g \in \mathcal{G}, A \in \mathcal{A}$, and $u \in SBV(A, \mathbb{R}^m)$ we set

$$E^{f,g}(u, A) := \int_A f(x, \nabla u) dx + \int_{S_u \cap A} g(x, [u], \nu_u) d\mathcal{H}^{n-1},$$

and for every $w \in SBV(A, \mathbb{R}^m)$ we set

$$m^{f,g}(w, A) := \inf\{E^{f,g}(u, A) : u \in SBV(A, \mathbb{R}^m), u = w \text{ near } \partial A\}. \tag{3.4}$$

The expression “ $u = w$ near ∂A ” means that there exists a neighbourhood U of ∂A such that $u = w$ \mathcal{L}^n -a.e. in $U \cap A$. More in general, if $\Lambda \subset \partial A$ is a relatively open subset of ∂A , the expression “ $u = w$ near Λ ” means that there exists a neighbourhood U of Λ in \mathbb{R}^n such that $u = w$ \mathcal{L}^n -a.e. in $U \cap A$.

For technical reasons, related to the details of the statement of the Subadditive Ergodic Theorem, it is convenient to extend this definition to an arbitrary bounded subset A of \mathbb{R}^n , by setting $m^{f,g}(w, A) := m^{f,g}(w, \text{int}A)$, where $\text{int}A$ denotes the interior of A .

Remark 3.2 If $f \in \mathcal{F}$ and $g \in \mathcal{G}$, then $f^\infty \in \mathcal{F}$ and $g_0 \in \mathcal{G}$. Moreover, the lower bounds (f3) and (g3) imply that

$$c_2|Du|(A) \leq \min\{E^{f,g}(u, A), E^{f,g_0}(u, A), E^{f^\infty,g}(u, A), E^{f^\infty,g_0}(u, A)\}$$

for every $A \in \mathcal{A}$ and $u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, with $u|_A \in SBV(A, \mathbb{R}^m)$.

Remark 3.3 From (f4) and (f5) it follows that for every $L > 0$ there exists $M > 0$, depending on c_3, c_4, c_5 , and L , such that

$$|f^\infty(x, \xi) - \frac{1}{t}f(x, t\xi)| \leq \frac{M}{t^\alpha} \text{ for every } x \in \mathbb{R}^n, \xi \in \mathbb{R}^{m \times n} \text{ with } |\xi| = 1, \text{ and } t \geq L. \tag{3.5}$$

Conversely, if the limit in (3.1) exists and f satisfies (f3), (f4), and (3.5) for some $L > 0$ and $M > 0$, then for every $x \in \mathbb{R}^n, \xi \in \mathbb{R}^{m \times n}$, and $t > 0$ we have

$$\begin{cases} |f^\infty(x, \xi) - \frac{1}{t}f(x, t\xi)| \leq \frac{M}{t^\alpha}|\xi|^{1-\alpha} = \frac{M}{t}|t\xi|^{1-\alpha} \leq \frac{M}{t^{c_2^{1-\alpha}}}f(x, t\xi)^{1-\alpha} & \text{if } t|\xi| \geq L, \\ |f^\infty(x, \xi) - \frac{1}{t}f(x, t\xi)| \leq \frac{2c_3L}{t} + \frac{c_4}{t} & \text{if } t|\xi| < L, \end{cases}$$

where in the last inequality we used the fact that that $f^\infty(x, \xi) \leq c_3|\xi|$ for every $x \in \mathbb{R}^n$ and $\xi \in \mathbb{R}^{m \times n}$. This implies that f satisfies (3.2) for a suitable constant c_5 , depending only on c_2, c_3, c_4, L , and M .

Remark 3.4 If (f4) holds, then (f5) is equivalent to the fact that

$$|\frac{1}{s}f(x, s\xi) - \frac{1}{t}f(x, t\xi)| \leq \frac{c_5}{s} + \frac{c_5}{s}f(x, s\xi)^{1-\alpha} + \frac{c_5}{t} + \frac{c_5}{t}f(x, t\xi)^{1-\alpha} \tag{3.6}$$

for every $x \in \mathbb{R}^n$, every $\xi \in \mathbb{R}^{m \times n}$, and every $s, t > 0$. Indeed, using the triangle inequality we obtain (3.6) from (3.2) for s and t . Conversely, if (f4) holds, then $\frac{c_5}{t} + \frac{c_5}{t}f(x, t\xi)^{1-\alpha} \rightarrow 0$ as $t \rightarrow +\infty$. Therefore (3.6) implies that $t \mapsto \frac{1}{t}f(x, t\xi)$ satisfies the Cauchy condition as $t \rightarrow +\infty$, hence the limit in (3.1) exists, while (3.2) follows from (3.6) by taking the limit as $s \rightarrow +\infty$.

Remark 3.5 Assume that $g: \mathbb{R}^n \times \mathbb{R}^m \times \mathbb{S}^{n-1} \rightarrow [0, +\infty)$ satisfies (g5) and let $\lambda: [0, +\infty) \rightarrow [0, +\infty)$ be defined by

$$\lambda(t) := \sup\{|g_0(x, \zeta, \nu) - \frac{1}{t}g(x, t\zeta, \nu)| : x \in \mathbb{R}^n, \zeta \in \mathbb{S}^{m-1}, \nu \in \mathbb{S}^{n-1}, t \in (0, 1]\}. \tag{3.7}$$

Then λ is nondecreasing and

$$\lim_{t \rightarrow 0+} \lambda(t) = 0, \tag{3.8}$$

$$|g_0(x, \zeta, \nu) - \frac{1}{t}g(x, t\zeta, \nu)| \leq |\zeta|\lambda(t|\zeta|), \tag{3.9}$$

for every $x \in \mathbb{R}^n, \zeta \in \mathbb{R}^m$, and $\nu \in \mathbb{S}^{n-1}$. If g satisfies also (g3), then (3.9) gives

$$|g_0(x, \zeta, \nu) - \frac{1}{t}g(x, t\zeta, \nu)| \leq \frac{1}{c_2}\lambda(t|\zeta|)\frac{1}{t}g(x, t\zeta, \nu). \tag{3.10}$$

Conversely, if the limit in (3.3) exists (even with no uniformity assumptions) and g satisfies (g4) and (3.10), then it satisfies (3.9) with λ replaced by $\frac{c_3}{c_2}\lambda$, which implies that g satisfies (g5).

Remark 3.6 If $g : \mathbb{R}^n \times \mathbb{R}^m \times \mathbb{S}^{n-1} \rightarrow [0, +\infty)$ satisfies (g3), (g4), and (g5), then by (3.10) and by the triangle inequality we get

$$|\frac{1}{s}g(x, s\zeta, \nu) - \frac{1}{t}g(x, t\zeta, \nu)| \leq \frac{1}{c_2}\lambda(s|\zeta|)\frac{1}{s}g(x, s\zeta, \nu) + \frac{1}{c_2}\lambda(t|\zeta|)\frac{1}{t}g(x, t\zeta, \nu) \tag{3.11}$$

for every $s, t > 0, x \in \mathbb{R}^n, \zeta \in \mathbb{R}^m$, and $\nu \in \mathbb{S}^{n-1}$. Conversely, if (g4) and (3.11) hold, with some function λ satisfying (3.8), then $\lambda(t|\zeta|)\frac{1}{t}g(x, t\zeta, \nu) \rightarrow 0$ as $t \rightarrow 0+$ and hence, using (3.11), we deduce that the function $t \mapsto \frac{1}{t}g(x, t\zeta, \nu)$ satisfies the Cauchy condition as $t \rightarrow 0+$. This implies that the limit in (3.3) exists and is finite. Moreover, passing to the limit as $s \rightarrow 0+$ from (3.11) we obtain (3.10), which, in turn, yields (g5).

We are now ready to introduce the probabilistic setting of our problem. In what follows (Ω, \mathcal{T}, P) denotes a fixed probability space.

Definition 3.7 (Random integrands) A function $f : \Omega \times \mathbb{R}^n \times \mathbb{R}^{m \times n} \rightarrow [0, +\infty)$ is called a *random volume integrand* if

- (a1) f is $\mathcal{T} \otimes \mathcal{B}^n \otimes \mathcal{B}^{m \times n}$ -measurable;
- (b1) $f(\omega, \cdot, \cdot) \in \mathcal{F}$ for every $\omega \in \Omega$.

A function $g : \Omega \times \mathbb{R}^n \times \mathbb{R}^m \times \mathbb{S}^{n-1} \rightarrow [0, +\infty)$ is called a *random surface integrand* if

- (a2) g is $\mathcal{T} \otimes \mathcal{B}^n \otimes \mathcal{B}^m \otimes \mathcal{B}_S^{n-1}$ -measurable;
- (b2) $g(\omega, \cdot, \cdot, \cdot) \in \mathcal{G}$ for every $\omega \in \Omega$.

Let f be a random volume integrand and let g be a random surface integrand. For every $\omega \in \Omega$ and every $\varepsilon > 0$ we consider the free-discontinuity functional $E_\varepsilon(\omega) : L^1_{loc}(\mathbb{R}^n, \mathbb{R}^m) \times \mathcal{A} \rightarrow [0, +\infty]$ defined by

$$\begin{aligned}
 & E_\varepsilon(\omega)(u, A) \\
 & := \begin{cases} \int_A f(\omega, \frac{x}{\varepsilon}, \nabla u) dx + \int_{S_u \cap A} g(\omega, \frac{x}{\varepsilon}, [u], \nu_u) d\mathcal{H}^{n-1} & \text{if } u|_A \in SBV(A, \mathbb{R}^m), \\ +\infty & \text{otherwise in } L^1_{loc}(\mathbb{R}^n, \mathbb{R}^m). \end{cases}
 \end{aligned}
 \tag{3.12}$$

Definition 3.8 If f is a random volume integrand, we define $f^\infty : \Omega \times \mathbb{R}^n \times \mathbb{R}^{m \times n} \rightarrow [0, +\infty)$ by

$$f^\infty(\omega, x, \xi) := \lim_{t \rightarrow +\infty} \frac{1}{t} f(\omega, x, t\xi).
 \tag{3.13}$$

If g is a random surface integrand, we define $g_0 : \Omega \times \mathbb{R}^n \times \mathbb{R}^m \times \mathbb{S}^{n-1} \rightarrow [0, +\infty)$ by

$$g_0(\omega, x, \zeta, \nu) := \lim_{t \rightarrow 0^+} \frac{1}{t} g(\omega, x, t\zeta, \nu).
 \tag{3.14}$$

Remark 3.9 The existence of the limit in (3.13) follows from (b1) in Definition 3.7 and from (f5). Since for every $t > 0$ the functions $(\omega, x, \xi) \mapsto \frac{1}{t} f(\omega, x, t\xi)$ are $\mathcal{T} \otimes \mathcal{B}^n \otimes \mathcal{B}^{m \times n}$ -measurable by (a1), the same property holds for f^∞ . Moreover, from Remark 3.2 and from (b1) we deduce that $f^\infty(\omega, \cdot, \cdot) \in \mathcal{F}$ for every $\omega \in \Omega$. We conclude that f^∞ is a random volume integrand.

The existence of the limit in (3.14) follows from (b2) in Definition 3.7 and from (g5). Since for every $t > 0$ the functions $(\omega, x, \zeta, \nu) \mapsto \frac{1}{t} g(\omega, x, t\zeta, \nu)$ are $\mathcal{T} \otimes \mathcal{B}^n \otimes \mathcal{B}^m \otimes \mathcal{B}^n_{\mathbb{S}}$ -measurable by (a2), the same property holds for g_0 . Moreover, from Remark 3.2 and from (b2) we deduce that $g_0(\omega, \cdot, \cdot, \cdot) \in \mathcal{G}$ for every $\omega \in \Omega$. We conclude that g_0 is a random surface integrand.

In the study of stochastic homogenisation an important role is played by the notions introduced by the following definitions.

Definition 3.10 (*P-preserving transformation*) A P -preserving transformation on (Ω, \mathcal{T}, P) is a map $T : \Omega \rightarrow \Omega$ satisfying the following properties:

- (a) (measurability) T is \mathcal{T} -measurable;
- (b) (bijectivity) T is bijective;
- (c) (invariance) $P(T(E)) = P(E)$, for every $E \in \mathcal{T}$.

If, in addition, every set $E \in \mathcal{T}$ which satisfies $T(E) = E$ (called T -invariant set) has probability 0 or 1, then T is called *ergodic*.

Definition 3.11 (*Group of P-preserving transformations*) Let d be a positive integer. A group of P -preserving transformations on (Ω, \mathcal{T}, P) is a family $(\tau_z)_{z \in \mathbb{Z}^d}$ of mappings $\tau_z : \Omega \rightarrow \Omega$ satisfying the following properties:

- (a) (measurability) τ_z is \mathcal{T} -measurable for every $z \in \mathbb{Z}^d$;
- (b) (bijectivity) τ_z is bijective for every $z \in \mathbb{Z}^d$;
- (c) (invariance) $P(\tau_z(E)) = P(E)$, for every $E \in \mathcal{T}$ and every $z \in \mathbb{Z}^d$;

(d) (group property) $\tau_0 = id_\Omega$ (the identity map on Ω) and $\tau_{z+z'} = \tau_z \circ \tau_{z'}$ for every $z, z' \in \mathbb{Z}^d$.

If, in addition, every set $E \in \mathcal{T}$ which satisfies $\tau_z(E) = E$ for every $z \in \mathbb{Z}^d$ has probability 0 or 1, then $(\tau_z)_{z \in \mathbb{Z}^d}$ is called *ergodic*.

We are now in a position to define the notion of stationary random integrand.

Definition 3.12 (*Stationary random integrand*) A random volume integrand f is *stationary* with respect to a group $(\tau_z)_{z \in \mathbb{Z}^n}$ of P -preserving transformations on (Ω, \mathcal{T}, P) if

$$f(\omega, x + z, \xi) = f(\tau_z(\omega), x, \xi)$$

for every $\omega \in \Omega, x \in \mathbb{R}^n, z \in \mathbb{Z}^n$, and $\xi \in \mathbb{R}^{m \times n}$.

Similarly, a random surface integrand g is *stationary* with respect to $(\tau_z)_{z \in \mathbb{Z}^n}$ if

$$g(\omega, x + z, \zeta, \nu) = g(\tau_z(\omega), x, \zeta, \nu)$$

for every $\omega \in \Omega, x \in \mathbb{R}^n, z \in \mathbb{Z}^n, \zeta \in \mathbb{R}^m$, and $\nu \in \mathbb{S}^{n-1}$.

We now recall the notion of subadditive stochastic process as well as the Subadditive Ergodic Theorem by Akcoglu and Krengel [1, Theorem 2.7].

Let d be a positive integer. For every $a, b \in \mathbb{R}^d$, with $a_i < b_i$ for $i = 1, \dots, d$, we define

$$[a, b) := \{x \in \mathbb{R}^d : a_i \leq x_i < b_i \text{ for } i = 1, \dots, d\},$$

and we set

$$\mathcal{I}_d := \{[a, b) : a, b \in \mathbb{R}^d, a_i < b_i \text{ for } i = 1, \dots, d\}. \tag{3.15}$$

Definition 3.13 (*Subadditive process*) A d -dimensional *subadditive process* with respect to a group $(\tau_z)_{z \in \mathbb{Z}^d}, d \geq 1$, of P -preserving transformations on (Ω, \mathcal{T}, P) is a function $\mu : \Omega \times \mathcal{I}_d \rightarrow \mathbb{R}$ satisfying the following properties:

- (a) (measurability) for every $A \in \mathcal{I}_d$ the function $\omega \mapsto \mu(\omega, A)$ is \mathcal{T} -measurable;
- (b) (covariance) for every $\omega \in \Omega, A \in \mathcal{I}_d$, and $z \in \mathbb{Z}^d$ we have $\mu(\omega, A + z) = \mu(\tau_z(\omega), A)$;
- (c) (subadditivity) for every $A \in \mathcal{I}_d$ and for every *finite* family $(A_i)_{i \in I} \subset \mathcal{I}_d$ of pairwise disjoint sets such that $A = \cup_{i \in I} A_i$, we have

$$\mu(\omega, A) \leq \sum_{i \in I} \mu(\omega, A_i) \quad \text{for every } \omega \in \Omega;$$

- (d) (boundedness) there exists $c > 0$ such that $0 \leq \mu(\omega, A) \leq c\mathcal{L}^d(A)$ for every $\omega \in \Omega$ and every $A \in \mathcal{I}_d$.

Definition 3.14 (*Regular family of sets*) A family of sets $(A_t)_{t>0}$ in \mathcal{I}_d is called *regular* (with constant $C > 0$) if there exists another family of sets $(A'_t)_{t>0} \subset \mathcal{I}_d$ such that:

- (a) $A_t \subset A'_t$ for every $t > 0$;
- (b) $A'_s \subset A'_t$ whenever $0 < s < t$;
- (c) $0 < \mathcal{L}^d(A'_t) \leq C\mathcal{L}^d(A_t)$ for every $t > 0$.

If the family $(A'_t)_{t>0}$ can be chosen in a way such that $\mathbb{R}^d = \bigcup_{t>0} A'_t$, then we write $\lim_{t \rightarrow +\infty} A_t = \mathbb{R}^d$.

We now state a variant of the pointwise ergodic Theorem [1, Theorem 2.7 and Remark p. 59] which is suitable for our purposes. This variant can be found in [30, Theorem 4.1].

Theorem 3.15 (Subadditive Ergodic Theorem) *Let $d \in \{n - 1, n\}$ and let $(\tau_z)_{z \in \mathbb{Z}^d}$ be a group of P -preserving transformations on (Ω, \mathcal{T}, P) . Let $\mu: \Omega \times \mathcal{I}_d \rightarrow \mathbb{R}$ be a subadditive process with respect to $(\tau_z)_{z \in \mathbb{Z}^d}$. Then there exist a \mathcal{T} -measurable function $\varphi: \Omega \rightarrow [0, +\infty)$ and a set $\Omega' \in \mathcal{T}$ with $P(\Omega')=1$ such that*

$$\lim_{t \rightarrow +\infty} \frac{\mu(\omega, A_t)}{\mathcal{L}^d(A_t)} = \varphi(\omega),$$

for every regular family of sets $(A_t)_{t>0} \subset \mathcal{I}_d$ with $\lim_{t \rightarrow +\infty} A_t = \mathbb{R}^d$ and for every $\omega \in \Omega'$. If in addition $(\tau_z)_{z \in \mathbb{Z}^d}$ is ergodic, then φ is constant P -a.e.

Remark 3.16 (Covariance with respect to a continuous group $(\tau_z)_{z \in \mathbb{R}^d}$) Definitions 3.11, 3.12, 3.13 and Theorem 3.15 can be adapted also to the case of a continuous group $(\tau_z)_{z \in \mathbb{R}^d}$, see for instance [17, Section 3.1].

3.2 Statement of the main results

In this section we state the main result of the paper, Theorem 3.18, which provides a Γ -convergence result for the random functionals $(E_\varepsilon(\omega))_{\varepsilon>0}$ introduced in (3.12), under the assumption that the volume and surface integrands f and g are stationary.

The next theorem proves the existence of the limits in the asymptotic cell formulas that will be used in the statement of the main result.

When f and g are random integrands it is convenient to introduce the following shorthand notation

$$\begin{aligned} m_\omega^{f,g_0} &:= m^{f(\omega, \cdot, \cdot), g_0(\omega, \cdot, \cdot)}, & m_\omega^{f^\infty, g} &:= m^{f^\infty(\omega, \cdot, \cdot), g(\omega, \cdot, \cdot)}, \\ m_\omega^{f^\infty, g_0} &:= m^{f^\infty(\omega, \cdot, \cdot), g_0(\omega, \cdot, \cdot)}, \end{aligned} \tag{3.16}$$

where m^{f,g_0} , $m^{f^\infty, g}$, and m^{f^∞, g_0} are defined as in (3.4), with (f, g) replaced by (f, g_0) , (f^∞, g) , and (f^∞, g_0) , respectively.

Theorem 3.17 (Homogenisation formulas) *Let f be a stationary random volume integrand and let g be a stationary random surface integrand with respect to a group $(\tau_z)_{z \in \mathbb{Z}^n}$ of P -preserving transformations on (Ω, \mathcal{T}, P) . Then there exists $\Omega' \in \mathcal{T}$, with $P(\Omega') = 1$, such that*

(a) for every $\omega \in \Omega', x \in \mathbb{R}^n, \xi \in \mathbb{R}^{m \times n}, v \in \mathbb{S}^{n-1}$, and $k \in \mathbb{N}$ the limit

$$\lim_{r \rightarrow +\infty} \frac{m_{\omega}^{f, g_0}(\ell_{\xi}, Q_r^{v, k}(rx))}{k^{n-1}r^n}$$

exists and is independent of x, v , and k ;

(b) for every $\omega \in \Omega', x \in \mathbb{R}^n, \zeta \in \mathbb{R}^m, v \in \mathbb{S}^{n-1}$ the limit

$$\lim_{r \rightarrow +\infty} \frac{m_{\omega}^{f^{\infty}, g}(u_{rx, \zeta, v}, Q_r^v(rx))}{r^{n-1}}$$

exists and is independent of x .

More precisely, there exist a random volume integrand $f_{\text{hom}}: \Omega \times \mathbb{R}^{m \times n} \rightarrow [0, +\infty)$, and a random surface integrand $g_{\text{hom}}: \Omega \times \mathbb{R}^m \times \mathbb{S}^{n-1} \rightarrow [0, +\infty)$ such that for every $\omega \in \Omega', x \in \mathbb{R}^n, \xi \in \mathbb{R}^{m \times n}, \zeta \in \mathbb{R}^m$, and $v \in \mathbb{S}^{n-1}$

$$f_{\text{hom}}(\omega, \xi) = \lim_{r \rightarrow +\infty} \frac{m_{\omega}^{f, g_0}(\ell_{\xi}, Q_r^{v, k}(rx))}{k^{n-1}r^n} = \lim_{r \rightarrow +\infty} \frac{m_{\omega}^{f, g_0}(\ell_{\xi}, Q_r)}{r^n}, \tag{3.17}$$

$$g_{\text{hom}}(\omega, \zeta, v) = \lim_{r \rightarrow +\infty} \frac{m_{\omega}^{f^{\infty}, g}(u_{rx, \zeta, v}, Q_r^v(rx))}{r^{n-1}} = \lim_{r \rightarrow +\infty} \frac{m_{\omega}^{f^{\infty}, g}(u_{0, \zeta, v}, Q_r^v)}{r^{n-1}}, \tag{3.18}$$

where $Q_r := Q_r(0)$ and $Q_r^v = Q_r^v(0)$.

For every $\omega \in \Omega'$ and $\xi \in \mathbb{R}^{m \times n}$ let

$$f_{\text{hom}}^{\infty}(\omega, \xi) := \lim_{t \rightarrow +\infty} \frac{f_{\text{hom}}(\omega, t\xi)}{t}$$

(since $f_{\text{hom}}(\omega, \cdot) \in \mathcal{F}$, the existence of the limit is guaranteed by (f5)). Then for every $\omega \in \Omega', x \in \mathbb{R}^n, \xi \in \mathbb{R}^{m \times n}, v \in \mathbb{S}^{n-1}$, and $k \in \mathbb{N}$ we have

$$f_{\text{hom}}^{\infty}(\omega, \xi) = \lim_{r \rightarrow +\infty} \frac{m_{\omega}^{f^{\infty}, g_0}(\ell_{\xi}, Q_r^{v, k}(rx))}{k^{n-1}r^n} = \lim_{r \rightarrow +\infty} \frac{m_{\omega}^{f^{\infty}, g_0}(\ell_{\xi}, Q_r)}{r^n}. \tag{3.19}$$

If, in addition, $(\tau_z)_{z \in \mathbb{Z}^n}$ is ergodic, then f_{hom} and g_{hom} are independent of ω and

$$f_{\text{hom}}(\xi) = \lim_{r \rightarrow +\infty} \frac{1}{r^n} \int_{\Omega} m_{\omega}^{f, g_0}(\ell_{\xi}, Q_r) dP(\omega), \tag{3.20}$$

$$g_{\text{hom}}(\zeta, v) = \lim_{r \rightarrow +\infty} \frac{1}{r^{n-1}} \int_{\Omega} m_{\omega}^{f^{\infty}, g}(u_{0, \zeta, v}, Q_r^v) dP(\omega), \tag{3.21}$$

$$f_{\text{hom}}^{\infty}(\xi) = \lim_{r \rightarrow +\infty} \frac{1}{r^n} \int_{\Omega} m_{\omega}^{f^{\infty}, g_0}(\ell_{\xi}, Q_r) dP(\omega). \tag{3.22}$$

We are now ready to state the main result of this paper, namely the almost sure Γ -convergence of the sequence of random functionals $(E_{\varepsilon}(\omega))_{\varepsilon > 0}$ introduced in (3.12).

Theorem 3.18 (Almost sure Γ -convergence) *Let f and g be stationary random volume and surface integrands with respect to a group $(\tau_z)_{z \in \mathbb{Z}^n}$ of P -preserving transformations on (Ω, \mathcal{T}, P) , and for every $\varepsilon > 0$ and $\omega \in \Omega$ let $E_\varepsilon(\omega)$ be as in (3.12). Let $\Omega' \in \mathcal{T}$ (with $P(\Omega') = 1$), $f_{\text{hom}}, f_{\text{hom}}^\infty$ and g_{hom} be as in Theorem 3.17, and for every $\omega \in \Omega$ let $E_{\text{hom}}(\omega) : L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m) \times \mathcal{A} \rightarrow [0, +\infty]$ be the functional defined by*

$$E_{\text{hom}}(\omega)(u, A) := \int_A f_{\text{hom}}(\omega, \nabla u) \, dx + \int_{S_u \cap A} g_{\text{hom}}(\omega, [u], \nu_u) \, d\mathcal{H}^{n-1} + \int_A f_{\text{hom}}^\infty\left(\omega, \frac{dC(u)}{d|C(u)|}\right) d|C(u)|,$$

if $u|_A \in BV(A, \mathbb{R}^m)$, and by $E_{\text{hom}}(\omega)(u, A) := +\infty$, if $u|_A \notin BV(A, \mathbb{R}^m)$. Then for every $\omega \in \Omega'$ and every $A \in \mathcal{A}$ the functionals $E_\varepsilon(\omega)(\cdot, A)$ Γ -converge to $E_{\text{hom}}(\omega)(\cdot, A)$ in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, as $\varepsilon \rightarrow 0+$.

If, in addition, $(\tau_z)_{z \in \mathbb{Z}^n}$ is ergodic, then E_{hom} is a deterministic functional; i.e., it does not depend on ω .

Thanks to Theorem 3.18 we can also characterise the asymptotic behaviour of some minimisation problems involving $E_\varepsilon(\omega)$. An example is shown in the corollary below. Since for every $A \in \mathcal{A}$ the values of $E_\varepsilon(\omega)(u, A)$ and $E_{\text{hom}}(\omega)(u, A)$ depend only on the restriction of u to A , in the corollary we regard $E_\varepsilon(\omega)(u, \cdot)$ and $E_{\text{hom}}(\omega)(u, \cdot)$ as functionals defined on $L^1(A, \mathbb{R}^m)$.

Corollary 3.19 (Convergence of minima and minimisers) *Let f and g be stationary random volume and surface integrands with respect to a group $(\tau_z)_{z \in \mathbb{Z}^n}$ of P -preserving transformations on (Ω, \mathcal{T}, P) , and for every $\varepsilon > 0$ and $\omega \in \Omega$ let $E_\varepsilon(\omega)$ be as in (3.12). Let $\Omega' \in \mathcal{T}$ (with $P(\Omega') = 1$) be as in Theorem 3.17, and let $E_{\text{hom}}(\omega)$ be as in Theorem 3.18. Given $\omega \in \Omega', A \in \mathcal{A}$, and $h \in L^1(A, \mathbb{R}^m)$, we have*

$$\inf_{u \in SBV(A, \mathbb{R}^m)} (E_\varepsilon(\omega)(u, A) + \|u - h\|_{L^1(A, \mathbb{R}^m)}) \rightarrow \min_{u \in BV(A, \mathbb{R}^m)} (E_{\text{hom}}(\omega)(u, A) + \|u - h\|_{L^1(A, \mathbb{R}^m)}) \tag{3.23}$$

as $\varepsilon \rightarrow 0+$. Moreover, if $(u_\varepsilon) \subset SBV(A, \mathbb{R}^m)$ is a sequence such that

$$E_\varepsilon(\omega)(u_\varepsilon, A) + \|u_\varepsilon - h\|_{L^1(A, \mathbb{R}^m)} \leq \inf_{u \in SBV(A, \mathbb{R}^m)} (E_\varepsilon(\omega)(u, A) + \|u - h\|_{L^1(A, \mathbb{R}^m)}) + \eta_\varepsilon \tag{3.24}$$

for some $\eta_\varepsilon \rightarrow 0+$, then there exists a sequence $\varepsilon_j \rightarrow 0+$ such that $(u_{\varepsilon_j})_{j \in \mathbb{N}}$ converges in $L^1(A, \mathbb{R}^m)$, as $j \rightarrow +\infty$, to a solution of the minimisation problem

$$\min_{u \in BV(A, \mathbb{R}^m)} (E_{\text{hom}}(\omega)(u, A) + \|u - h\|_{L^1(A, \mathbb{R}^m)}). \tag{3.25}$$

Proof If A has a Lipschitz boundary, then the functionals $E_\varepsilon(\omega)(\cdot, A) + \|\cdot - h\|_{L^1(A, \mathbb{R}^m)}$ are equi-coercive in $L^1(A, \mathbb{R}^m)$ thanks to Remark 3.2. Since we have Γ -convergence in $L^1(A, \mathbb{R}^m)$ by virtue of Theorem 3.18, the proof readily follows from the fundamental property of Γ -convergence (see, e.g., [18, Corollary 7.20]).

We now show that the convergence of minimum values and of minimisers can be obtained even if ∂A is not regular. Let us fix $\omega \in \Omega'$, $A \in \mathcal{A}$, and $h \in L^1(A, \mathbb{R}^m)$. By Theorem 3.18 for every $A' \in \mathcal{A}$ the functional $E_{\text{hom}}(\omega)(\cdot, A')$ is a Γ -limit in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, hence it is lower semicontinuous in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ (see [18, Proposition 6.8]). This implies that $E_{\text{hom}}(\omega)(\cdot, A')$, considered as a functional on $L^1(A', \mathbb{R}^m)$, is lower semicontinuous. Since

$$E_{\text{hom}}(\omega)(\cdot, A) = \sup\{E_{\text{hom}}(\omega)(\cdot, A') : A' \in \mathcal{A}, A' \subset\subset A\},$$

the functional $E_{\text{hom}}(\omega)(\cdot, A)$, defined on $L^1(A, \mathbb{R}^m)$, is lower semicontinuous with respect to the convergence in $L^1_{\text{loc}}(A, \mathbb{R}^m)$.

Since $f_{\text{hom}}(\omega, \cdot)$ and $f_{\text{hom}}^\infty(\omega, \cdot)$ belong to \mathcal{F} , while $g_{\text{hom}}(\omega, \cdot, \cdot)$ belongs to \mathcal{G} , it follows from the definition of $E_{\text{hom}}(\omega)(\cdot, A)$ that $c_2|Du|(A) + \|u\|_{L^1(A, \mathbb{R}^m)} \leq E_{\text{hom}}(\omega)(u, A) + \|u - h\|_{L^1(A, \mathbb{R}^m)} + \|h\|_{L^1(A, \mathbb{R}^m)}$ for every $u \in BV(A, \mathbb{R}^m)$. This shows that the functional $u \mapsto E_{\text{hom}}(\omega)(u, A) + \|u - h\|_{L^1(A, \mathbb{R}^m)}$ is coercive in $BV(A, \mathbb{R}^m)$ with respect to the convergence in $L^1_{\text{loc}}(A, \mathbb{R}^m)$. Therefore it attains a minimum value in $BV(A, \mathbb{R}^m)$, which we denote by μ_0 .

Let u_0 be a minimum point in $BV(A, \mathbb{R}^m)$. We extend u_0 to a function of $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, still denoted by u_0 . By Γ -convergence, for every sequence (ε_j) of positive numbers converging to 0 there exists a sequence u_j converging to u_0 in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ such that $E_{\varepsilon_j}(\omega)(u_j, A) \rightarrow E_{\text{hom}}(\omega)(u_0, A) < +\infty$. By the definition of $E_{\varepsilon_j}(\omega)$ we have $u_j \in SBV(A, \mathbb{R}^m)$ for j large enough, hence

$$\inf_{u \in SBV(A, \mathbb{R}^m)} (E_{\varepsilon_j}(\omega)(u, A) + \|u - h\|_{L^1(A, \mathbb{R}^m)}) \leq E_{\varepsilon_j}(\omega)(u_j, A) + \|u_j - h\|_{L^1(A, \mathbb{R}^m)}.$$

This implies that

$$\limsup_{j \rightarrow +\infty} \inf_{u \in SBV(A, \mathbb{R}^m)} (E_{\varepsilon_j}(\omega)(u, A) + \|u - h\|_{L^1(A, \mathbb{R}^m)}) \leq \mu_0.$$

Since the sequence $\varepsilon_j \rightarrow 0$ is arbitrary, we obtain

$$\limsup_{\varepsilon \rightarrow 0^+} \inf_{u \in SBV(A, \mathbb{R}^m)} (E_\varepsilon(\omega)(u, A) + \|u - h\|_{L^1(A, \mathbb{R}^m)}) \leq \mu_0. \tag{3.26}$$

To prove the opposite inequality for the liminf, as well as the last statement of the corollary, we fix a sequence $(u_\varepsilon) \subset SBV(A, \mathbb{R}^m)$ satisfying (3.24). For every sequence (ε_j) of positive numbers converging to 0, by Remark 3.2 and by (3.26) the sequence (u_{ε_j}) is bounded in $BV(A, \mathbb{R}^m)$. Therefore a subsequence, not relabelled, converges in $L^1_{\text{loc}}(A, \mathbb{R}^m)$ to a function $u_* \in BV(A, \mathbb{R}^m)$.

Given $A' \in \mathcal{A}$, with $A' \subset\subset A$, we can consider the functions v_j , defined by $v_j := u_{\varepsilon_j}$ in A' and $v_j := 0$ in $\mathbb{R}^n \setminus A'$, which converge in $L^1(\mathbb{R}^n, \mathbb{R}^m)$ to the function v_* , defined by $v_* := u_*$ in A' and $v_* := 0$ in $\mathbb{R}^n \setminus A'$. Since $E_{\varepsilon_j}(\omega)(\cdot, A')$ Γ -converges to $E_{\text{hom}}(\omega)(\cdot, A')$ in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, we have

$$\begin{aligned}
 E_{\text{hom}}(\omega)(u_*, A') &= E_{\text{hom}}(\omega)(v_*, A') \leq \liminf_{j \rightarrow +\infty} E_{\varepsilon_j}(\omega)(v_j, A') \\
 &\leq \liminf_{j \rightarrow +\infty} E_{\varepsilon_j}(\omega)(u_{\varepsilon_j}, A),
 \end{aligned}$$

which implies that

$$\begin{aligned}
 E_{\text{hom}}(\omega)(u_*, A') + \|u_* - h\|_{L^1(A', \mathbb{R}^m)} \\
 \leq \liminf_{j \rightarrow +\infty} (E_{\varepsilon_j}(\omega)(u_{\varepsilon_j}, A) + \|u_{\varepsilon_j} - h\|_{L^1(A, \mathbb{R}^m)}).
 \end{aligned}$$

Taking the supremum for $A' \subset\subset A$ in the previous inequalities we obtain

$$E_{\text{hom}}(\omega)(u_*, A) \leq \liminf_{j \rightarrow +\infty} E_{\varepsilon_j}(\omega)(u_{\varepsilon_j}, A), \tag{3.27}$$

and

$$\begin{aligned}
 \mu_0 &\leq E_{\text{hom}}(\omega)(u_*, A) + \|u_* - h\|_{L^1(A, \mathbb{R}^m)} \\
 &\leq \liminf_{j \rightarrow +\infty} (E_{\varepsilon_j}(\omega)(u_{\varepsilon_j}, A) + \|u_{\varepsilon_j} - h\|_{L^1(A, \mathbb{R}^m)}) \\
 &\leq \liminf_{j \rightarrow +\infty} \inf_{u \in SBV(A, \mathbb{R}^m)} (E_{\varepsilon_j}(\omega)(u, A) + \|u - h\|_{L^1(A, \mathbb{R}^m)}). \tag{3.28}
 \end{aligned}$$

By the arbitrariness of the sequence $\varepsilon_j \rightarrow 0+$, this chain of inequalities, together with (3.26), gives (3.23) and shows that u_* is a solution of the minimisation problem (3.25).

In turn, (3.23), (3.27), and (3.28) imply that $\|u_{\varepsilon_j} - h\|_{L^1(A, \mathbb{R}^m)} \rightarrow \|u_* - h\|_{L^1(A, \mathbb{R}^m)}$. Since u_{ε_j} converges to u_* in $L^1_{\text{loc}}(A, \mathbb{R}^m)$, from the general version of the Dominated Convergence Theorem we obtain that u_{ε_j} converges to u_* in $L^1(A, \mathbb{R}^m)$. This concludes the proof of the last statement of the corollary. \square

4 Deterministic homogenisation: properties of the homogenised integrands

Let $f \in \mathcal{F}$ and $g \in \mathcal{G}$. For $\varepsilon > 0$ consider the functionals $E_\varepsilon : L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m) \times \mathcal{A} \rightarrow [0, +\infty]$ defined by

$$\begin{aligned}
 E_\varepsilon(u, A) \\
 := \begin{cases} \int_A f(\frac{x}{\varepsilon}, \nabla u) dx + \int_{S_u \cap A} g(\frac{x}{\varepsilon}, [u], \nu_u) d\mathcal{H}^{n-1} & \text{if } u|_A \in SBV(A, \mathbb{R}^m), \\ +\infty & \text{otherwise in } L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m). \end{cases} \tag{4.1}
 \end{aligned}$$

In this section we prove the Γ -convergence of E_ε under suitable assumptions on f and g , which are more general than the periodicity with respect to x .

The main result of this section is the following theorem.

Theorem 4.1 (Homogenisation) *Let $f \in \mathcal{F}$, $g \in \mathcal{G}$, and let m^{f,g_0} and $m^{f^\infty,g}$ be defined as in (3.4) with (f, g) replaced by (f, g_0) and (f^∞, g) , respectively. Assume that*

(a) *for every $x \in \mathbb{R}^n$, $\xi \in \mathbb{R}^{m \times n}$, $v \in \mathbb{S}^{n-1}$, and $k \in \mathbb{N}$ the limit*

$$\lim_{r \rightarrow +\infty} \frac{m^{f,g_0}(\ell_\xi, Q_r^{v,k}(rx))}{k^{n-1}r^n} =: f_{\text{hom}}(\xi) \tag{4.2}$$

exists and is independent of x , v , and k ;

(b) *for every $x \in \mathbb{R}^n$, $\zeta \in \mathbb{R}^m$, and $v \in \mathbb{S}^{n-1}$ the limit*

$$\lim_{r \rightarrow +\infty} \frac{m^{f^\infty,g}(u_{rx,\zeta,v}, Q_r^v(rx))}{r^{n-1}} =: g_{\text{hom}}(\zeta, v) \tag{4.3}$$

exists and is independent of x .

Then $f_{\text{hom}} \in \mathcal{F}$ and $g_{\text{hom}} \in \mathcal{G}$. Let f_{hom}^∞ be the recession function of f_{hom} and let $E_{\text{hom}} : L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m) \times \mathcal{A} \rightarrow [0, +\infty]$ be the functional defined by

$$\begin{aligned} E_{\text{hom}}(u, A) &:= \int_A f_{\text{hom}}(\nabla u) \, dx + \int_{S_u \cap A} g_{\text{hom}}([u], v_u) \, d\mathcal{H}^{n-1} \\ &+ \int_A f_{\text{hom}}^\infty\left(\frac{dC(u)}{|dC(u)|}\right) \, d|C(u)| \end{aligned} \tag{4.4}$$

if $u|_A \in BV(A, \mathbb{R}^m)$, while $E_{\text{hom}}(u, A) := +\infty$ if $u|_A \notin BV(A, \mathbb{R}^m)$. Then, for every $A \in \mathcal{A}$ the functionals $E_\varepsilon(\cdot, A)$ defined as in (4.1) Γ -converge to $E_{\text{hom}}(\cdot, A)$ in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, as $\varepsilon \rightarrow 0+$, meaning that for every sequence (ε_j) of positive numbers converging to zero the sequence $(E_{\varepsilon_j}(\cdot, A))$ Γ -converges to $E_{\text{hom}}(\cdot, A)$ in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$.

The proof of the homogenisation result Theorem 4.1 will be carried out in three main steps. In the first step (Lemmas 4.2 and 4.5) we show that $f_{\text{hom}} \in \mathcal{F}$ and $g_{\text{hom}} \in \mathcal{G}$. In the second step (Theorem 5.1) we prove that, up to subsequences, for every $A \in \mathcal{A}$ the functionals $E_\varepsilon(\cdot, A)$ Γ -converge to some functional $\widehat{E}(\cdot, A)$, whose domain is $BV(A, \mathbb{R}^m)$. Further, we prove that \widehat{E} satisfies some suitable properties both as a functional and as a set-function. In particular $\widehat{E}(u, \cdot)$ is the restriction to \mathcal{A} of a Borel measure.

In the third and last step we show that (4.2) and (4.3) imply, respectively, that the following identities hold true for every $A \in \mathcal{A}$ and for every $u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ with $u|_A \in BV(A, \mathbb{R}^m)$:

$$\frac{d\widehat{E}(u, \cdot)}{d\mathcal{L}^n}(x) = f_{\text{hom}}(\nabla u(x)) \quad \text{for } \mathcal{L}^n\text{-a.e. } x \in A, \tag{4.5}$$

$$\frac{d\widehat{E}(u, \cdot)}{d\mathcal{H}^{n-1} \llcorner S_u}(x) = g_{\text{hom}}([u](x), v_u(x)) \quad \text{for } \mathcal{H}^{n-1}\text{-a.e. } x \in S_u \cap A, \tag{4.6}$$

$$\frac{d\widehat{E}(u, \cdot)}{d|C(u)|}(x) = f_{\text{hom}}^\infty\left(\frac{dC(u)}{d|C(u)|}(x)\right) \quad \text{for } |C(u)|\text{-a.e. } x \in A \tag{4.7}$$

(see Propositions 6.2, 7.2, and 8.3). Moreover, thanks to (4.5)-(4.7) we deduce that \widehat{E} coincides with the functional E_{hom} defined in (4.4); as a consequence, the Γ -convergence result proved in the second step actually holds true for the whole sequence (E_ε) .

In the next lemmas we prove that the homogenised integrands f_{hom} and g_{hom} belong to the classes \mathcal{F} and \mathcal{G} , respectively.

Lemma 4.2 *Let $f \in \mathcal{F}$ and $g \in \mathcal{G}$. Assume that hypothesis (a) of Theorem 4.1 is satisfied and let f_{hom} be defined as in (4.2). Then $f_{\text{hom}} \in \mathcal{F}$.*

Proof To prove (f2) we fix $\xi_1, \xi_2 \in \mathbb{R}^{m \times n}$ and set $\xi := \xi_2 - \xi_1$. We claim that for every $r > 0$

$$\begin{aligned} & |m^{f, g_0}(\ell_{\xi_1}, Q_r) - m^{f, g_0}(\ell_{\xi_2}, Q_r)| \\ & \leq \sigma_1(|\xi|)(m^{f, g_0}(\ell_{\xi_1}, Q_r) + m^{f, g_0}(\ell_{\xi_2}, Q_r)) + c_1|\xi|r^n, \end{aligned} \tag{4.8}$$

where $Q_r := Q_r(0)$. Indeed, by (f2), for every $u \in SBV(Q_r, \mathbb{R}^m)$ we have

$$\begin{aligned} E^{f, g_0}(u + \ell_\xi, Q_r) & \leq E^{f, g_0}(u, Q_r) + \sigma_1(|\xi|)(E^{f, g_0}(u + \ell_\xi, Q_r) \\ & \quad + E^{f, g_0}(u, Q_r)) + c_1|\xi|r^n. \end{aligned}$$

By rearranging the terms we get

$$(1 - \sigma_1(|\xi|))E^{f, g_0}(u + \ell_\xi, Q_r) \leq (1 + \sigma_1(|\xi|))E^{f, g_0}(u, Q_r) + c_1|\xi|r^n.$$

If $\sigma_1(|\xi|) < 1$, we minimise over all functions $u \in SBV(Q_r, \mathbb{R}^m)$ such that $u = \ell_{\xi_1}$ near ∂Q_r and, using (3.4), we obtain

$$(1 - \sigma_1(|\xi|))m^{f, g_0}(\ell_{\xi_2}, Q_r) \leq (1 + \sigma_1(|\xi|))m^{f, g_0}(\ell_{\xi_1}, Q_r) + c_1|\xi|r^n.$$

This inequality is trivial if $\sigma_1(|\xi|) \geq 1$. Exchanging the roles of ξ_1 and ξ_2 we obtain (4.8). We now divide both sides of this inequality by r^n and, passing to the limit as $r \rightarrow +\infty$, from (4.2) we obtain that f_{hom} satisfies (f2).

Property (f1) for f_{hom} follows from the continuity estimate (f2), since f_{hom} does not depend on x . The lower bound (f3) for f_{hom} follows from the lower bound in Remark 3.2, which gives

$$c_2 \inf |Du|(Q_r) \leq m^{f, g_0}(\ell_\xi, Q_r)$$

for every $\xi \in \mathbb{R}^{m \times n}$, where the infimum is over all functions $u \in SBV(Q_r, \mathbb{R}^m)$, and such that $u = \ell_\xi$ near ∂Q_r . By Jensen’s inequality the left-hand side is equal to $c_2|\xi|r^n$. Using (4.2) we conclude that that f_{hom} satisfies (f3).

Property (f4) for f_{hom} follows from the fact that for every $\xi \in \mathbb{R}^{m \times n}$ we have

$$\frac{1}{r^n} m^{f, g_0}(\ell_\xi, Q_r) \leq \frac{1}{r^n} E^{f, g_0}(\ell_\xi, Q_r) = \frac{1}{r^n} \int_{Q_r} f(x, \xi) dx \leq c_3 |\xi| + c_4.$$

Passing to the limit as $r \rightarrow +\infty$, from (4.2) we obtain that f_{hom} satisfies (f4).

We now prove that f_{hom} satisfies (f5). Fix $\xi \in \mathbb{R}^{m \times n}$, $s > 0, t > 0$, and $\eta \in (0, 1)$. By (3.4) for every $r > 0$ there exists $u_r \in SBV(Q_r, \mathbb{R}^m)$, with $u_r = \ell_\xi$ near ∂Q_r , such that

$$\int_{Q_r} f(x, t \nabla u_r) dx + \int_{S_{u_r} \cap Q_r} g_0(x, t[u_r], v_{u_r}) d\mathcal{H}^{n-1} \leq m^{f, g_0}(\ell_{t\xi}, Q_r) + \eta r^n. \tag{4.9}$$

By (3.6) for every $x \in \mathbb{R}^n$ and $\xi \in \mathbb{R}^{m \times n}$ we have

$$\frac{1}{s} f(x, s\xi) - \frac{c_5}{s} - \frac{c_5}{s} f(x, s\xi)^{1-\alpha} \leq \frac{1}{t} f(x, t\xi) + \frac{c_5}{t} + \frac{c_5}{t} f(x, t\xi)^{1-\alpha},$$

hence, using the positive 1-homogeneity of g_0 ,

$$\begin{aligned} & \frac{1}{s} \frac{1}{r^n} \int_{Q_r} f(x, s \nabla u_r) dx - \frac{c_5}{s} - \frac{c_5}{s} \frac{1}{r^n} \int_{Q_r} f(x, s \nabla u_r)^{1-\alpha} dx \\ & \quad + \frac{1}{s} \frac{1}{r^n} \int_{S_{u_r} \cap Q_r} g_0(x, s[u_r], v_{u_r}) d\mathcal{H}^{n-1} \\ & \leq \frac{1}{t} \frac{1}{r^n} \int_{Q_r} f(x, t \nabla u_r) dx + \frac{c_5}{t} + \frac{c_5}{t} \frac{1}{r^n} \int_{Q_r} f(x, t \nabla u_r)^{1-\alpha} dx \\ & \quad + \frac{1}{t} \frac{1}{r^n} \int_{S_{u_r} \cap Q_r} g_0(x, t[u_r], v_{u_r}) d\mathcal{H}^{n-1}. \end{aligned}$$

By Hölder’s inequality we obtain

$$\begin{aligned} & \frac{1}{s} \frac{1}{r^n} \int_{Q_r} f(x, s \nabla u_r) dx + \frac{1}{s} \frac{1}{r^n} \int_{S_{u_r} \cap Q_r} g_0(x, s[u_r], v_{u_r}) d\mathcal{H}^{n-1} - \frac{c_5}{s} \\ & \quad - \frac{c_5}{s} \left(\frac{1}{r^n} \int_{Q_r} f(x, s \nabla u_r) dx \right)^{1-\alpha} \\ & \leq \frac{1}{t} \frac{1}{r^n} \int_{Q_r} f(x, t \nabla u_r) dx + \frac{1}{t} \frac{1}{r^n} \int_{S_{u_r} \cap Q_r} g_0(x, t[u_r], v_{u_r}) d\mathcal{H}^{n-1} + \frac{c_5}{t} \\ & \quad + \frac{c_5}{t} \left(\frac{1}{r^n} \int_{Q_r} f(x, t \nabla u_r) dx \right)^{1-\alpha}. \end{aligned}$$

By (4.9) this inequality implies that

$$\frac{1}{s} \left(\frac{1}{r^n} \int_{Q_r} f(x, s \nabla u_r) dx + \frac{1}{r^n} \int_{S_{u_r} \cap Q_r} g_0(x, s[u_r], v_{u_r}) d\mathcal{H}^{n-1} \right) - \frac{c_5}{s}$$

$$\begin{aligned}
 & -\frac{c_5}{s} \left(\frac{1}{r^n} \int_{Q_r} f(x, s\nabla u_r) dx + \frac{1}{r^n} \int_{S_{u_r} \cap Q_r} g_0(x, s[u_r], v_{u_r}) d\mathcal{H}^{n-1} \right)^{1-\alpha} \\
 & \leq \frac{1}{t} \left(\frac{1}{r^n} m^{f, g_0}(\ell_{t\xi}, Q_r) + \eta \right) + \frac{c_5}{t} + \frac{c_5}{t} \left(\frac{1}{r^n} m^{f, g_0}(\ell_{t\xi}, Q_r) + \eta \right)^{1-\alpha}. \tag{4.10}
 \end{aligned}$$

If

$$\frac{1}{r^n} m^{f, g_0}(\ell_{s\xi}, Q_r) - c_5 \left(\frac{1}{r^n} m^{f, g_0}(\ell_{s\xi}, Q_r) \right)^{1-\alpha} \leq 0, \tag{4.11}$$

then we have

$$\begin{aligned}
 & \frac{1}{s} \frac{1}{r^n} m^{f, g_0}(\ell_{s\xi}, Q_r) - \frac{c_5}{s} - \frac{c_5}{s} \left(\frac{1}{r^n} m^{f, g_0}(\ell_{s\xi}, Q_r) \right)^{1-\alpha} \\
 & \leq \frac{1}{t} \left(\frac{1}{r^n} m^{f, g_0}(\ell_{t\xi}, Q_r) + \eta \right) + \frac{c_5}{t} + \frac{c_5}{t} \left(\frac{1}{r^n} m^{f, g_0}(\ell_{t\xi}, Q_r) + \eta \right)^{1-\alpha}, \tag{4.12}
 \end{aligned}$$

just because the left-hand side is negative and the right-hand side is positive. Since the function $\tau \mapsto \tau - c_5\tau^{1-\alpha}$, defined for $\tau > 0$, is increasing in the half-line where it is positive, from the inequality

$$m^{f, g_0}(\ell_{s\xi}, Q_r) \leq \int_{Q_r} f(x, s\nabla u_r) dx + \int_{S_{u_r} \cap Q_r} g_0(x, s[u_r], v_{u_r}) d\mathcal{H}^{n-1}$$

and from (4.10) we deduce that (4.12) is satisfied even if (4.11) is not.

Passing to the limit first as $r \rightarrow +\infty$ and then as $\eta \rightarrow 0+$, from (4.2) and (4.12) we obtain

$$\frac{1}{s} f_{\text{hom}}(s\xi) - \frac{c_5}{s} - \frac{c_5}{s} f_{\text{hom}}(s\xi)^{1-\alpha} \leq \frac{1}{t} f_{\text{hom}}(t\xi) + \frac{c_5}{t} + \frac{c_5}{t} f_{\text{hom}}(t\xi)^{1-\alpha}.$$

By exchanging the roles of s and t we obtain (3.6). Recalling that f_{hom} satisfies (f4), we can apply Remark 3.4 and we obtain that f_{hom} satisfies (f5). □

To prove that $g_{\text{hom}} \in \mathcal{G}$ we need the truncation result given by the following lemma, which will be used several times in this paper. The proof is given in [9, Lemma 3.7] (see also [10, Lemma 2.8]).

Lemma 4.3 *Let $C_1 > 0$, $C_2 > 0$, and $\eta > 0$. Then there exists a constant $M = M(C_1, C_2, \eta) > 0$ such that for every $f \in \mathcal{F}$ and $g \in \mathcal{G}$, for every $A \in \mathcal{A}$, for every $w \in SBV(A, \mathbb{R}^m) \cap L^\infty(A, \mathbb{R}^m)$, with $\|w\|_{L^\infty(A, \mathbb{R}^m)} \leq C_1$, and for every $u \in BV(A, \mathbb{R}^m)$, with $\|u\|_{L^1(A, \mathbb{R}^m)} + |Du|(A) \leq C_2$ and $u = w$ near ∂A , there exists $\tilde{u} \in SBV(A, \mathbb{R}^m) \cap L^\infty(A, \mathbb{R}^m)$ such that*

- (a) $\|\tilde{u}\|_{L^\infty(A, \mathbb{R}^m)} \leq M$,
- (b) $E^{f, g}(\tilde{u}, A) \leq E^{f, g}(u, A) + \eta$,
- (c) $\|\tilde{u} - w\|_{L^1(A, \mathbb{R}^m)} \leq \|u - w\|_{L^1(A, \mathbb{R}^m)}$,
- (d) $\tilde{u} = w$ near ∂A .

Remark 4.4 A careful inspection of the proof of [9, Lemma 3.7] shows that the lemma also applies if u only attains the boundary conditions on a subset of ∂A , as defined after (3.4). More precisely, if $\Lambda \subset \partial A$ is a relatively open subset of ∂A , U is a neighbourhood of Λ in \mathbb{R}^n , $w \in SBV(A, \mathbb{R}^m) \cap L^\infty(U \cap A, \mathbb{R}^m)$, and $u = w$ \mathcal{L}^n -a.e. in $U \cap A$, then the conclusion still holds true, with (d) replaced by

$$(d') \quad \tilde{u} = w \text{ } \mathcal{L}^n\text{-a.e. in } U \cap A,$$

and in this case $M = M(\tilde{C}_1, C_2, \eta) > 0$, where $\|w\|_{L^\infty(U \cap A, \mathbb{R}^m)} \leq \tilde{C}_1$.

We are now ready to prove that $g_{\text{hom}} \in \mathcal{G}$.

Lemma 4.5 *Let $f \in \mathcal{F}$ and $g \in \mathcal{G}$. Assume that hypothesis (b) of Theorem 4.1 is satisfied and let g_{hom} be defined as in (4.3). Then $g_{\text{hom}} \in \mathcal{G}$.*

Proof To prove (g2) we fix $\zeta_1, \zeta_2 \in \mathbb{R}^m$ and $\nu \in \mathbb{S}^{n-1}$, and we set $\zeta := \zeta_2 - \zeta_1$. We claim that for every $r > 0$

$$\begin{aligned} & |m^{f^\infty, g}(u_{0, \zeta_1, \nu}, Q_r^\nu) - m^{f^\infty, g}(u_{0, \zeta_2, \nu}, Q_r^\nu)| \\ & \leq \sigma_2(|\zeta|)(m^{f^\infty, g}(u_{0, \zeta_1, \nu}, Q_r^\nu) + m^{f^\infty, g}(u_{0, \zeta_2, \nu}, Q_r^\nu)), \end{aligned} \tag{4.13}$$

where $Q_r^\nu := Q_r^\nu(0)$. Indeed, for every $u \in SBV(Q_r^\nu, \mathbb{R}^m)$, by (g2) we have

$$\begin{aligned} & E^{f^\infty, g}(u + u_{0, \zeta, \nu}, Q_r^\nu) \\ & \leq E^{f^\infty, g}(u, Q_r^\nu) + \sigma_2(|\zeta|)(E^{f^\infty, g}(u + u_{0, \zeta, \nu}, Q_r^\nu) + E^{f^\infty, g}(u, Q_r^\nu)). \end{aligned}$$

By rearranging the terms we get

$$(1 - \sigma_2(|\zeta|))E^{f^\infty, g}(u + u_{0, \zeta, \nu}, Q_r^\nu) \leq (1 + \sigma_2(|\zeta|))E^{f^\infty, g}(u, Q_r^\nu).$$

If $\sigma_2(|\zeta|) < 1$, we minimise over all functions $u \in SBV(Q_r^\nu, \mathbb{R}^m)$ such that $u = u_{0, \zeta_1, \nu}$ near ∂Q_r^ν and by (3.4) we obtain

$$(1 - \sigma_2(|\zeta|))m^{f^\infty, g}(u_{0, \zeta_2, \nu}, Q_r^\nu) \leq (1 + \sigma_2(|\zeta|))m^{f^\infty, g}(u_{0, \zeta_1, \nu}, Q_r^\nu).$$

This inequality is trivial if $\sigma_2(|\zeta|) \geq 1$. Exchanging the roles of ζ_1 and ζ_2 we obtain (4.13). We now divide both sides of this inequality by r^{n-1} and, passing to the limit as $r \rightarrow +\infty$, from (4.3) we obtain that g_{hom} satisfies (g2).

In view of (g2), to prove (g1) for g_{hom} it is enough to show that for every $\zeta \in \mathbb{R}^m$ the restriction of the function $\nu \mapsto g_{\text{hom}}(\zeta, \nu)$ to the sets $\widehat{\mathbb{S}}_+^{n-1}$ and $\widehat{\mathbb{S}}_-^{n-1}$ is continuous. We only prove this property for $\widehat{\mathbb{S}}_+^{n-1}$, the other case being analogous. To this end, let us fix $\zeta \in \mathbb{R}^m$, $\nu \in \widehat{\mathbb{S}}_+^{n-1}$, and a sequence $(\nu_j) \subset \widehat{\mathbb{S}}_+^{n-1}$ such that $\nu_j \rightarrow \nu$ as $j \rightarrow +\infty$. Since the function $\nu \mapsto R_\nu$ is continuous on $\widehat{\mathbb{S}}_+^{n-1}$, for every $\delta \in (0, \frac{1}{2})$ there exists an integer j_δ such that

$$|\nu_j - \nu| < \delta \quad \text{and} \quad Q_{(1-\delta)r}^{\nu_j} \subset\subset Q_r^\nu \subset\subset Q_{(1+\delta)r}^{\nu_j}, \tag{4.14}$$

for every $j \geq j_\delta$ and every $r > 0$. Fix $j \geq j_\delta, r > 0$, and $\eta > 0$. By (3.4) there exists $u \in SBV(Q_r^v, \mathbb{R}^m)$, with $u = u_{0,\zeta,v}$ near ∂Q_r^v , such that

$$\int_{Q_r^v} f^\infty(x, \nabla u) \, dx + \int_{S_u \cap Q_r^v} g(x, [u], v_u) \, d\mathcal{H}^{n-1} \leq m^{f^\infty, g}(u_{0,\zeta,v}, Q_r^v) + \eta. \tag{4.15}$$

We define $v \in SBV_{loc}(Q_{(1+\delta)r}^{v_j}, \mathbb{R}^m)$ as

$$v(x) := \begin{cases} u(x) & \text{if } x \in Q_r^v, \\ u_{0,\zeta,v_j}(x) & \text{if } x \in Q_{(1+\delta)r}^{v_j} \setminus Q_r^v. \end{cases}$$

Then $v = u_{0,\zeta,v_j}$ near $\partial Q_{(1+\delta)r}^{v_j}$ and $S_v \subset S_u \cup \Sigma$, where

$$\Sigma := \{x \in \partial Q_r^v : (x \cdot \nu)(x \cdot \nu_j) < 0\} \cup (\Pi_0^{v_j} \cap (Q_{(1+\delta)r}^{v_j} \setminus Q_r^v)).$$

By (4.14) there exists $\zeta(\delta) > 0$, independent of j and r , with $\zeta(\delta) \rightarrow 0$ as $\delta \rightarrow 0+$, such that $\mathcal{H}^{n-1}(\Sigma) \leq \zeta(\delta)r^{n-1}$. Thanks to (g4), (3.4) and (4.15) we then have

$$\begin{aligned} m^{f^\infty, g}(u_{0,\zeta,v_j}, Q_{(1+\delta)r}^{v_j}) &\leq \int_{Q_{(1+\delta)r}^{v_j}} f^\infty(x, \nabla v) \, dx + \int_{S_v \cap Q_{(1+\delta)r}^{v_j}} g(x, [v], v_v) \, d\mathcal{H}^{n-1} \\ &\leq \int_{Q_r^v} f^\infty(x, \nabla u) \, dx + \int_{S_u \cap Q_r^v} g(x, [u], v_u) \, d\mathcal{H}^{n-1} + c_3|\zeta|\zeta(\delta)r^{n-1} \\ &\leq m^{f^\infty, g}(u_{0,\zeta,v}, Q_r^v) + \eta + c_3|\zeta|\zeta(\delta)r^{n-1}, \end{aligned}$$

where we used the fact that $f^\infty(x, 0) = 0$ for every $x \in \mathbb{R}^n$. Dividing by r^{n-1} and passing to the limit as $r \rightarrow +\infty$, recalling (4.3) we obtain

$$g_{\text{hom}}(\zeta, v_j)(1 + \delta)^{n-1} \leq g_{\text{hom}}(\zeta, v) + c_3|\zeta|\zeta(\delta).$$

Letting $j \rightarrow +\infty$ and then $\delta \rightarrow 0+$ we deduce that

$$\limsup_{j \rightarrow +\infty} g_{\text{hom}}(\zeta, v_j) \leq g_{\text{hom}}(\zeta, v).$$

An analogous argument, now using the cubes $Q_{(1-\delta)r}^{v_j}$, implies that

$$g_{\text{hom}}(\zeta, v) \leq \liminf_{j \rightarrow +\infty} g_{\text{hom}}(\zeta, v_j),$$

hence the restriction of the function $v \mapsto g_{\text{hom}}(\zeta, v)$ to $\widehat{\mathbb{S}}_+^{n-1}$ is continuous. This concludes the proof of (g1) for g_{hom} .

The lower bound (g3) for g_{hom} can be obtained from the lower bound in Remark 3.2, which gives

$$c_2 \inf |Du|(Q_r^v) \leq m^{f^\infty, g}(u_{0, \zeta, v}, Q_r^v)$$

for every $\zeta \in \mathbb{R}^m$ and $v \in \mathbb{S}^{n-1}$, where the infimum is over all functions $u \in SBV(Q_r^v, \mathbb{R}^m)$ and such that $u = u_{0, \zeta, v}$ near ∂Q_r^v . In turn, this infimum is larger than or equal to

$$c_2 \inf |Dv|(Q_r^v), \tag{4.16}$$

where the infimum is now over all scalar functions $v \in SBV(Q_r^v)$ and such that $v = u_{0, |\zeta|, v}$ near ∂Q_r^v . Using (4.3), property (g3) for g_{hom} follows from these inequalities and from the fact that the value of (4.16) is $c_2|\zeta|r^{n-1}$. This is a well known fact, which can be proved, for instance, using a slicing argument based on [5, Theorem 3.103].

Property (g4) for g_{hom} follows from the fact that for every $\zeta \in \mathbb{R}^m$ and $v \in \mathbb{S}^{n-1}$ we have

$$\begin{aligned} \frac{1}{r^{n-1}} m^{f^\infty, g}(u_{0, \zeta, v}, Q_r^v) &\leq \frac{1}{r^{n-1}} E^{f^\infty, g}(u_{0, \zeta, v}, Q_r^v) \\ &= \frac{1}{r^{n-1}} \int_{\Pi_0^v \cap Q_r^v} g(x, \zeta, v) dx \leq c_3 |\zeta|. \end{aligned}$$

Passing to the limit as $r \rightarrow +\infty$, from (4.3) we obtain that g_{hom} satisfies (g4).

We now prove that g_{hom} satisfies (g5). Fix $\zeta \in \mathbb{S}^{m-1}$, $v \in \mathbb{S}^{n-1}$, $s > 0$, $t > 0$, and $\eta \in (0, 1)$. By (3.4) for every $r > 0$ there exists $v_r \in SBV(Q_r^v, \mathbb{R}^m)$, with $v_r = u_{0, \zeta, v}$ near ∂Q_r^v , such that

$$\int_{Q_r^v} f^\infty(x, t\nabla v_r) dx + \int_{S_{v_r} \cap Q_r^v} g(x, t[v_r], v_{v_r}) d\mathcal{H}^{n-1} \leq m^{f^\infty, g}(u_{0, t\zeta, v}, Q_r^v) + \eta t r^{n-1},$$

hence

$$\begin{aligned} \int_{Q_r^v} f^\infty(x, \nabla v_r) dx + \frac{1}{t} \int_{S_{v_r} \cap Q_r^v} g(x, t[v_r], v_{v_r}) d\mathcal{H}^{n-1} \\ \leq \frac{1}{t} m^{f^\infty, g}(u_{0, t\zeta, v}, Q_r^v) + \eta r^{n-1}, \end{aligned}$$

where we used the positive 1-homogeneity of $f^\infty(x, \cdot)$.

Let $Q^v := Q_1^v(0)$ and let $w_r \in SBV(Q^v, \mathbb{R}^m)$ be the rescaled function, defined by $w_r(x) := v_r(rx)$ for every $x \in Q^v$. Then $w_r = u_{0, \zeta, v}$ near ∂Q^v and, by a change of variables,

$$\begin{aligned} \int_{Q^v} f^\infty(rx, \nabla w_r) dx + \frac{1}{t} \int_{S_{w_r} \cap Q^v} g(rx, t[w_r], v_{w_r}) d\mathcal{H}^{n-1} \\ \leq \frac{1}{t} \frac{1}{r^{n-1}} m^{f^\infty, g}(u_{0, t\zeta, v}, Q_r^v) + \eta, \end{aligned} \tag{4.17}$$

where we used the positive 1-homogeneity of $f^\infty(x, \cdot)$. Since the function $g_{r,t}$ defined by $g_{r,t}(x, \zeta, \nu) := \frac{1}{t}g(rx, t\zeta, \nu)$ satisfies (g3) with the constant c_2 independent of r and t , and by (g4)

$$\begin{aligned} \frac{1}{tr^{n-1}}m^{f^\infty, g}(u_{0,t\zeta, \nu}, Q_r^\nu) &\leq \frac{1}{tr^{n-1}}E^{f^\infty, g}(u_{0,t\zeta, \nu}, Q_r^\nu) \\ &= \frac{1}{tr^{n-1}} \int_{\Pi_0^\nu \cap Q_r^\nu} g(x, t\zeta, \nu) d\mathcal{H}^{n-1} \\ &\leq c_3|\zeta| = c_3, \end{aligned}$$

from Remark 3.2 and (4.17) we deduce that there exists a constant C such that $|Dw_r|(Q^\nu) \leq C$, for every $r > 0, t > 0$, and $\eta \in (0, 1)$. In addition, since w_r coincides with $u_{0,\zeta, \nu}$ near ∂Q^ν , we can apply Poincaré’s inequality and from the bound on its total variation we deduce that w_r is bounded in $BV(Q^\nu, \mathbb{R}^m)$, uniformly with respect to r , by a constant that we still denote with C ; namely, $\|w_r\|_{L^1(Q^\nu, \mathbb{R}^m)} + |Dw_r|(Q^\nu) \leq C$.

By Lemma 4.3 for every $\eta \in (0, 1)$ there exists a constant $M_\eta > 0$, depending on C but not on $t > 0, r > 0, \zeta \in \mathbb{S}^{m-1}$, and $\nu \in \mathbb{S}^{n-1}$, such that for every $r > 0$ there exists a function $\tilde{w}_r \in SBV(Q^\nu, \mathbb{R}^m) \cap L^\infty(Q^\nu, \mathbb{R}^m)$ with the following properties: $\tilde{w}_r = u_{0,\zeta, \nu}$ near ∂Q^ν , $\|\tilde{w}_r\|_{L^\infty(Q^\nu, \mathbb{R}^m)} \leq M_\eta$, and

$$\begin{aligned} &\int_{Q^\nu} f^\infty(rx, \nabla \tilde{w}_r) dx + \frac{1}{t} \int_{S_{\tilde{w}_r} \cap Q^\nu} g(rx, t[\tilde{w}_r], \nu_{\tilde{w}_r}) d\mathcal{H}^{n-1} \\ &\leq \int_{Q^\nu} f^\infty(rx, \nabla w_r) dx + \frac{1}{t} \int_{S_{w_r} \cap Q^\nu} g(rx, t[w_r], \nu_{w_r}) d\mathcal{H}^{n-1} + \eta, \end{aligned}$$

where we used the fact that $f^\infty \in \mathcal{F}$ and $g_{r,t} \in \mathcal{G}$. By (4.17) this implies that

$$\begin{aligned} &\int_{Q^\nu} f^\infty(rx, \nabla \tilde{w}_r) dx + \frac{1}{t} \int_{S_{\tilde{w}_r} \cap Q^\nu} g(rx, t[\tilde{w}_r], \nu_{\tilde{w}_r}) d\mathcal{H}^{n-1} \\ &\leq \frac{1}{t} \frac{1}{r^{n-1}} m^{f^\infty, g}(u_{0,t\zeta, \nu}, Q_r^\nu) + 2\eta. \end{aligned} \tag{4.18}$$

Let $\tilde{v}_r \in SBV(Q_r^\nu, \mathbb{R}^m) \cap L^\infty(Q_r^\nu, \mathbb{R}^m)$ be the function defined by $\tilde{v}_r(x) := \tilde{w}_r(\frac{x}{r})$ for every $x \in Q_r^\nu$. Then $\tilde{v}_r = u_{0,\zeta, \nu}$ near ∂Q_r^ν , $\|\tilde{v}_r\|_{L^\infty(Q_r^\nu, \mathbb{R}^m)} \leq M_\eta$, and, by a change of variables,

$$\begin{aligned} &\int_{Q_r^\nu} f^\infty(x, \nabla \tilde{v}_r) dx + \frac{1}{t} \int_{S_{\tilde{v}_r} \cap Q_r^\nu} g(x, t[\tilde{v}_r], \nu_{\tilde{v}_r}) d\mathcal{H}^{n-1} \\ &\leq \frac{1}{t} m^{f^\infty, g}(u_{0,t\zeta, \nu}, Q_r^\nu) + 2\eta r^{n-1}, \end{aligned} \tag{4.19}$$

where we used (4.18) and the positive 1-homogeneity of $f^\infty(x, \cdot)$. Since $\|\tilde{v}_r\|_{L^\infty(Q_r^\nu, \mathbb{R}^m)} \leq M_\eta$, by (3.11) we have

$$\left(1 - \frac{1}{c_2} \lambda(2sM_\eta)\right) \left(\int_{Q_r^\nu} f^\infty(x, \nabla \tilde{v}_r) dx + \frac{1}{s} \int_{S_{\tilde{v}_r} \cap Q_r^\nu} g(x, s[\tilde{v}_r], \nu_{\tilde{v}_r}) d\mathcal{H}^{n-1}\right)$$

$$\leq \left(1 + \frac{1}{c_2} \lambda(2tM_\eta)\right) \left(\int_{Q_r^v} f^\infty(x, \nabla \tilde{v}_r) dx + \frac{1}{t} \int_{S_{\tilde{v}_r} \cap Q_r^v} g(x, t[\tilde{v}_r], v_{\tilde{v}_r}) d\mathcal{H}^{n-1}\right). \tag{4.20}$$

Assume that

$$1 - \frac{1}{c_2} \lambda(2sM_\eta) > 0. \tag{4.21}$$

Since $s\tilde{v}_r = u_{0,s\zeta,v}$ near ∂Q_r^v , using the positive 1-homogeneity of f^∞ we obtain that

$$\frac{1}{s} m^{f^\infty, g}(u_{0,s\zeta,v}, Q_r^v) \leq \int_{Q_r^v} f^\infty(x, \nabla \tilde{v}_r) dx + \frac{1}{s} \int_{S_{\tilde{v}_r} \cap Q_r^v} g(x, s[\tilde{v}_r], v_{\tilde{v}_r}) d\mathcal{H}^{n-1}.$$

Hence from (4.19) and (4.20) we have

$$\begin{aligned} & \left(1 - \frac{1}{c_2} \lambda(2sM_\eta)\right) \frac{1}{s} m^{f^\infty, g}(u_{0,s\zeta,v}, Q_r^v) \\ & \leq \left(1 + \frac{1}{c_2} \lambda(2tM_\eta)\right) \left(\frac{1}{t} m^{f^\infty, g}(u_{0,t\zeta,v}, Q_r^v) + 2\eta r^{n-1}\right). \end{aligned} \tag{4.22}$$

This inequality holds also when (4.21) is not satisfied, since in that case the left-hand side is nonpositive.

Since M_η does not depend on $t, s,$ and $r,$ we can divide (4.22) by r^{n-1} and, passing to the limit as $r \rightarrow +\infty,$ by (4.3) we obtain

$$\left(1 - \frac{1}{c_2} \lambda(2sM_\eta)\right) \frac{1}{s} g_{\text{hom}}(s\zeta, v) \leq \left(1 + \frac{1}{c_2} \lambda(2tM_\eta)\right) \left(\frac{1}{t} g_{\text{hom}}(t\zeta, v) + 2\eta\right),$$

which gives

$$\begin{aligned} \frac{1}{s} g_{\text{hom}}(s\zeta, v) - \frac{1}{t} g_{\text{hom}}(t\zeta, v) & \leq \frac{1}{c_2} \lambda(2sM_\eta) \frac{1}{s} g_{\text{hom}}(s\zeta, v) + \frac{1}{c_2} \lambda(2tM_\eta) \frac{1}{t} g_{\text{hom}}(t\zeta, v) \\ & \quad + 2\eta \left(1 + \frac{1}{c_2} \lambda(2tM_\eta)\right). \end{aligned}$$

Since g_{hom} satisfies (g4) and $|\zeta| = 1,$ from the previous inequality we deduce that

$$\frac{1}{s} g_{\text{hom}}(s\zeta, v) - \frac{1}{t} g_{\text{hom}}(t\zeta, v) \leq \frac{c_3}{c_2} \lambda(2sM_\eta) + \frac{c_3}{c_2} \lambda(2tM_\eta) + 2\eta \left(1 + \frac{1}{c_2} \lambda(2tM_\eta)\right).$$

Exchanging the roles of s and t we obtain

$$\begin{aligned} \left| \frac{1}{s} g_{\text{hom}}(s\zeta, v) - \frac{1}{t} g_{\text{hom}}(t\zeta, v) \right| & \leq \frac{c_3}{c_2} \lambda(2sM_\eta) + \frac{c_3}{c_2} \lambda(2tM_\eta) + 2\eta \left(2 + \frac{1}{c_2} \lambda(2sM_\eta) \right. \\ & \quad \left. + \frac{1}{c_2} \lambda(2tM_\eta)\right) \end{aligned} \tag{4.23}$$

for every $s > 0, t > 0, \zeta \in \mathbb{S}^{m-1},$ and $v \in \mathbb{S}^{n-1}.$

Given $\tau > 0,$ we fix $\eta > 0$ such that $4\eta < \frac{\tau}{5}.$ Then, using (3.8), we find $\delta > 0$ such that for every $t \in (0, \delta)$ we have $\frac{c_3}{c_2} \lambda(2tM_\eta) < \frac{\tau}{5}$ and $2\eta \frac{1}{c_2} \lambda(2tM_\eta) < \frac{\tau}{5}.$ From (4.23) we obtain

$$|\frac{1}{s}g_{\text{hom}}(s\zeta, \nu) - \frac{1}{t}g_{\text{hom}}(t\zeta, \nu)| \leq \tau \tag{4.24}$$

for every $s, t \in (0, \delta)$, $\zeta \in \mathbb{S}^{m-1}$, and $\nu \in \mathbb{S}^{n-1}$. This shows that the function $t \mapsto \frac{1}{t}g_{\text{hom}}(t\zeta, \nu)$ satisfies the Cauchy condition for $t \rightarrow 0+$, hence the limit

$$g_{\text{hom},0}(\zeta, \nu) := \lim_{t \rightarrow 0+} \frac{1}{t}g_{\text{hom}}(t\zeta, \nu)$$

exists and is finite. This limit is uniform with respect to $\zeta \in \mathbb{S}^{m-1}$ and $\nu \in \mathbb{S}^{n-1}$ thanks to (4.24). This concludes the proof of (g5).

Property (g6) for g_{hom} follows from (4.3) and from the fact that $u_{0,-\zeta,-\nu} = u_{0,\zeta,\nu} - \zeta$ and $Q_r^\nu = Q_r^{-\nu}$ (see (h) in Section 2). □

5 Γ -convergence of a subsequence of (E_ε)

In this short section we show that, up to a subsequence, the functionals E_ε defined in (4.1) Γ -converge to some functional \widehat{E} as $\varepsilon \rightarrow 0+$, and study the main properties of this functional.

Theorem 5.1 (Properties of the Γ -limit) *Let $f \in \mathcal{F}$, let $g \in \mathcal{G}$, and for $\varepsilon > 0$ let $E_\varepsilon : L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m) \times \mathcal{A} \rightarrow [0, +\infty]$ be the functionals defined in (4.1). Then, for every sequence of positive numbers converging to zero, there exist a subsequence (ε_j) and a functional $\widehat{E} : L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m) \times \mathcal{A} \rightarrow [0, +\infty]$ such that for every $A \in \mathcal{A}$ the functionals $E_{\varepsilon_j}(\cdot, A)$ Γ -converge to $\widehat{E}(\cdot, A)$ in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, as $j \rightarrow +\infty$. Moreover, \widehat{E} satisfies the following properties:*

- (a) (locality) \widehat{E} is local; i.e., $\widehat{E}(u, A) = \widehat{E}(v, A)$ for every $A \in \mathcal{A}$ and every $u, v \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ such that $u = v$ \mathcal{L}^n -a.e. in A ;
- (b) (semicontinuity) for every $A \in \mathcal{A}$ the functional $\widehat{E}(\cdot, A)$ is lower semicontinuous in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$;
- (c) (bounds) for every $A \in \mathcal{A}$ and every $u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ we have

$$c_2|Du|(A) \leq \widehat{E}(u, A) \leq c_3|Du|(A) + c_4\mathcal{L}^n(A) \quad \text{if } u|_A \in BV(A, \mathbb{R}^m),$$

$$\widehat{E}(u, A) = +\infty \quad \text{otherwise;}$$

- (d) (measure property) for every $u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ the set function $\widehat{E}(u, \cdot)$ is the restriction to \mathcal{A} of a Borel measure defined on \mathcal{B}^n , which we still denote by $\widehat{E}(u, \cdot)$;
- (e) (translation invariance in u) for every $A \in \mathcal{A}$ and every $u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ we have

$$\widehat{E}(u + s, A) = \widehat{E}(u, A) \quad \text{for every } s \in \mathbb{R}^m.$$

Proof Given an infinitesimal sequence (ε_j) of positive numbers, let $\widehat{E}', \widehat{E}'': L^1_{loc}(\mathbb{R}^n, \mathbb{R}^m) \times \mathcal{A} \rightarrow [0, +\infty]$ be the functionals defined as

$$\widehat{E}'(\cdot, A) := \Gamma\text{-}\liminf_{j \rightarrow +\infty} E_{\varepsilon_j}(\cdot, A) \quad \text{and} \quad \widehat{E}''(\cdot, A) := \Gamma\text{-}\limsup_{j \rightarrow +\infty} E_{\varepsilon_j}(\cdot, A).$$

In view of the bounds (f3), (f4), (g3), and (g4) satisfied by f and g , it immediately follows that

$$\begin{aligned} c_2|Du|(A) &\leq \widehat{E}'(u, A) \\ &\leq \widehat{E}''(u, A) \leq c_3|Du|(A) + c_4\mathcal{L}^n(A) \quad \text{if } u|_A \in BV(A, \mathbb{R}^m), \end{aligned} \tag{5.1}$$

$$\widehat{E}'(u, A) = \widehat{E}''(u, A) = +\infty \quad \text{if } u|_A \notin BV(A, \mathbb{R}^m). \tag{5.2}$$

By the definition of E_{ε_j} and the general properties of Γ -convergence, we can also deduce that the functionals \widehat{E}' and \widehat{E}'' are local [18, Proposition 16.15], lower semicontinuous (in u) [18, Proposition 6.8], and increasing (in A) [18, Proposition 6.7]. Moreover \widehat{E}' is superadditive (in A) [18, Proposition 16.12]. Since it is not obvious that \widehat{E}' and \widehat{E}'' are inner regular (in A), at this stage of the proof we consider their inner regular envelopes; i.e., the functionals $\widehat{E}'_-, \widehat{E}''_-: L^1_{loc}(\mathbb{R}^n, \mathbb{R}^m) \times \mathcal{A} \rightarrow [0, +\infty]$ defined as

$$\widehat{E}'_-(u, A) := \sup_{\substack{A' \subset\subset A \\ A' \in \mathcal{A}}} \widehat{E}'(u, A') \quad \text{and} \quad \widehat{E}''_-(u, A) := \sup_{\substack{A' \subset\subset A \\ A' \in \mathcal{A}}} \widehat{E}''(u, A').$$

Also \widehat{E}'_- and \widehat{E}''_- are increasing, lower semicontinuous [18, Remark 15.10], and local [18, Remark 15.25]. Moreover, by [18, Theorem 16.9] we can find a subsequence of (ε_j) (not relabelled) such that the corresponding functionals \widehat{E}' and \widehat{E}'' satisfy

$$\widehat{E}'_- = \widehat{E}''_- =: \widehat{E}. \tag{5.3}$$

The functional \widehat{E} defined in (5.3) is inner regular [18, Remark 15.10] and superadditive [18, Proposition 16.12].

By virtue of [11, Proposition 3.1] applied with $p = 1$ we can immediately deduce that the functionals E_ε satisfy the so-called fundamental estimate uniformly in ε . Therefore [18, Proposition 18.4] yields the subadditivity of $\widehat{E}(u, \cdot)$. Therefore, invoking the measure-property criterion of De Giorgi and Letta [18, Theorem 14.23], we can deduce that, for every $u \in L^1_{loc}(\mathbb{R}^n, \mathbb{R}^m)$, the set function $\widehat{E}(u, \cdot)$ is the restriction to \mathcal{A} of a Borel measure defined on \mathcal{B}^n .

Moreover [18, Proposition 18.6] and (5.1) imply that $\widehat{E}(u, A) = \widehat{E}'(u, A) = \widehat{E}''(u, A)$ whenever $u|_A \in BV(A, \mathbb{R}^m)$. Finally, it follows from (5.1) and (5.2) that $\widehat{E}(u, A) = \widehat{E}'(u, A) = \widehat{E}''(u, A) = +\infty$ if $u|_A \notin BV(A, \mathbb{R}^m)$. We then conclude that $\widehat{E} = \widehat{E}' = \widehat{E}''$ in $L^1_{loc}(\mathbb{R}^n, \mathbb{R}^m) \times \mathcal{A}$, hence that $\widehat{E}(\cdot, A)$ is the Γ -limit of $E_{\varepsilon_j}(\cdot, A)$ in $L^1_{loc}(\mathbb{R}^n, \mathbb{R}^m)$ for every $A \in \mathcal{A}$.

Eventually, the translation invariance in u of $\widehat{E}(\cdot, A)$ can be easily checked arguing as in [11, Lemma 3.7]. □

For later use we need to introduce the following notation. Let $A \in \mathcal{A}$ and $w \in BV(A, \mathbb{R}^m)$, we set

$$m_{\widehat{E}}(w; A) := \inf\{\widehat{E}(u, A) : u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m), u|_A \in BV(A, \mathbb{R}^m), u = w \text{ near } \partial A\}.$$

6 Identification of the volume term

In Proposition 6.2 below we characterise the derivative of \widehat{E} with respect to the Lebesgue measure \mathcal{L}^n . In order to prove this result we need the estimate established in the following lemma, whose proof is an immediate consequence of (3.9).

Lemma 6.1 *Let $g \in \mathcal{G}$, $A \in \mathcal{A}$, and $u \in BV(A, \mathbb{R}^m) \cap L^\infty(A, \mathbb{R}^m)$. Then for every $t > 0$*

$$\begin{aligned} & \int_{S_u \cap A} |g_0(x, [u], \nu_u) - \frac{1}{t}g(x, t[u], \nu_u)| \, d\mathcal{H}^{n-1} \\ & \leq \lambda(t\|[u]\|_{L^\infty(S_u \cap A, \mathbb{R}^m)}) \int_{S_u \cap A} |[u]| \, d\mathcal{H}^{n-1}, \end{aligned}$$

where λ is the function defined in (3.7).

Proposition 6.2 (Homogenised volume integrand) *Let $f, g, E_\varepsilon, (\varepsilon_j)$, and \widehat{E} be as in Theorem 5.1. Assume that (a) of Theorem 4.1 holds, and let f_{hom} be as in (4.2). Then for every $A \in \mathcal{A}$ and every $u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, with $u|_A \in BV(A, \mathbb{R}^m)$, we have that*

$$\frac{d\widehat{E}(u, \cdot)}{d\mathcal{L}^n}(x) = f_{\text{hom}}(\nabla u(x)) \text{ for } \mathcal{L}^n\text{-a.e. } x \in A.$$

Proof Let us fix $A \in \mathcal{A}$ and $u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, with $u|_A \in BV(A, \mathbb{R}^m)$. We divide the proof into two steps.

Step 1: We claim that

$$\frac{d\widehat{E}(u, \cdot)}{d\mathcal{L}^n}(x) \leq f_{\text{hom}}(\nabla u(x)) \text{ for } \mathcal{L}^n\text{-a.e. } x \in A. \tag{6.1}$$

By (a)-(e) of Theorem 5.1 and by [10, Lemmas 3.1 and 3.5], arguing as in the proof of (3.16) in [10, Theorem 3.7], for \mathcal{L}^n -a.e. $x \in A$ we have

$$\frac{d\widehat{E}(u, \cdot)}{d\mathcal{L}^n}(x) = \lim_{\rho \rightarrow 0^+} \frac{m_{\widehat{E}}(\ell_{\xi(x)}, Q_\rho(x))}{\rho^n}, \tag{6.2}$$

where $\xi(x) := \nabla u(x)$. Fix $x \in A$ such that (6.2) holds and set $\xi := \xi(x) = \nabla u(x)$.

For every $\rho > 0$ we have

$$f_{\text{hom}}(\xi) = \lim_{r \rightarrow +\infty} \frac{m^{f, g_0}(\ell_\xi, Q_r(\frac{rx}{\rho}))}{r^n}, \tag{6.3}$$

since the above identity directly follows from (4.2) by replacing x with $\frac{x}{\rho}$.

Let us fix $\eta \in (0, 1)$. By (3.4) for every $\rho > 0$ and $r > 0$ there exists $v_r^\rho \in SBV(Q_r(\frac{rx}{\rho}), \mathbb{R}^m)$, with $v_r^\rho = \ell_\xi$ near $\partial Q_r(\frac{rx}{\rho})$, such that

$$\begin{aligned} E^{f, g_0}(v_r^\rho, Q_r(\frac{rx}{\rho})) &\leq m^{f, g_0}(\ell_\xi, Q_r(\frac{rx}{\rho})) + \eta r^n \leq \int_{Q_r(\frac{rx}{\rho})} f(y, \xi) dy + \eta r^n \\ &\leq (c_3|\xi| + c_4 + 1)r^n, \end{aligned} \tag{6.4}$$

where we used that $g_0(\cdot, 0, \cdot) = 0$ and (f4). We extend v_r^ρ to \mathbb{R}^n by setting $v_r^\rho(y) = \ell_\xi(y)$ for every $y \in \mathbb{R}^n \setminus Q_r(\frac{rx}{\rho})$.

For every $y \in \mathbb{R}^n$ let $w_r^\rho(y) := \frac{1}{r}v_r^\rho(\frac{rx}{\rho} + ry) - \frac{1}{\rho}\ell_\xi(x)$. Clearly $w_r^\rho \in SBV_{loc}(\mathbb{R}^n, \mathbb{R}^m)$ and $w_r^\rho = \ell_\xi$ near ∂Q and in $\mathbb{R}^n \setminus Q$, where $Q := Q_1(0)$. Moreover, by a change of variables we obtain

$$\begin{aligned} &\int_Q f(\frac{rx}{\rho} + ry, \nabla w_r^\rho(y)) dy + \int_{S_{w_r^\rho} \cap Q} g_0(\frac{rx}{\rho} + ry, [w_r^\rho](y), v_{w_r^\rho}(y)) d\mathcal{H}^{n-1}(y) \\ &= \frac{1}{r^n} E^{f, g_0}(v_r^\rho, Q_r(\frac{rx}{\rho})), \end{aligned} \tag{6.5}$$

where we used the 1-homogeneity of g_0 in the second variable. By the lower bounds (f3) and (g3), from (6.4) and (6.5) we deduce that there exists a constant K , depending on $|\xi|$, such that $|Dw_r^\rho|(Q) \leq K$ for every $\rho > 0$ and $r > 0$. In addition, since w_r^ρ coincides with ℓ_ξ in $\mathbb{R}^n \setminus Q$, we can apply Poincaré’s inequality and from the bound on its total variation we deduce that the sequence (w_r^ρ) is bounded in $BV_{loc}(\mathbb{R}^n, \mathbb{R}^m)$. In particular it is bounded in $BV(Q, \mathbb{R}^m)$, uniformly with respect to ρ and r , by a constant that we still denote with K .

By Lemma 4.3 and by (6.4) and (6.5) for every $\eta \in (0, 1)$ there exists a constant M_η , depending also on $|\xi|$ and K , such that for every $\rho > 0$ and $r > 0$ there exists $\tilde{w}_r^\rho \in SBV(Q, \mathbb{R}^m) \cap L^\infty(Q, \mathbb{R}^m)$ with the following properties: $\tilde{w}_r^\rho = \ell_\xi$ near ∂Q , $\|\tilde{w}_r^\rho\|_{L^\infty(Q, \mathbb{R}^m)} \leq M_\eta$, and

$$\begin{aligned} &\int_Q f(\frac{rx}{\rho} + ry, \nabla \tilde{w}_r^\rho(y)) dy + \int_{S_{\tilde{w}_r^\rho} \cap Q} g_0(\frac{rx}{\rho} + ry, [\tilde{w}_r^\rho](y), v_{\tilde{w}_r^\rho}(y)) d\mathcal{H}^{n-1}(y) \\ &\leq \frac{m^{f, g_0}(\ell_\xi, Q_r(\frac{rx}{\rho}))}{r^n} + 2\eta. \end{aligned}$$

Let $\tilde{v}_r^\rho \in SBV_{loc}(\mathbb{R}^n, \mathbb{R}^m)$ be defined by $\tilde{v}_r^\rho(y) := r\tilde{w}_r^\rho(\frac{y}{r} - \frac{x}{\rho}) + \frac{r}{\rho}\ell_\xi(x)$. Then $\tilde{v}_r^\rho = \ell_\xi$ near $\partial Q_r(\frac{rx}{\rho})$ and, by a change of variables,

$$\|[\tilde{v}_r^\rho]\|_{L^\infty(S_{\tilde{v}_r^\rho} \cap Q_r(\frac{rx}{\rho}), \mathbb{R}^m)} \leq 2M_\eta r, \tag{6.6}$$

$$E^{f, g_0}(\tilde{v}_r^\rho, Q_r(\frac{rx}{\rho})) \leq m^{f, g_0}(\ell_\xi, Q_r(\frac{rx}{\rho})) + 2\eta r^n. \tag{6.7}$$

Moreover, by combining (6.4) and (6.7) with the lower bound (g3) we immediately deduce the existence of a constant $C > 0$, depending on $|\xi|$, such that

$$\frac{1}{r^n} \int_{S_{\tilde{v}_r^\rho} \cap Q_r(\frac{rx}{\rho})} |[\tilde{v}_r^\rho]| d\mathcal{H}^{n-1} \leq C. \tag{6.8}$$

By Lemma 6.1, applied with $t = \rho/r$, using (6.6) and (6.8) we obtain

$$\frac{1}{r^n} \int_{S_{\tilde{v}_r^\rho} \cap Q_r(\frac{rx}{\rho})} \left| g_0(y, [\tilde{v}_r^\rho], v_{\tilde{v}_r^\rho} - \frac{r}{\rho} g(y, \frac{\rho}{r} [\tilde{v}_r^\rho], v_{\tilde{v}_r^\rho}) \right| d\mathcal{H}^{n-1} \leq C\lambda(2\rho M_\eta).$$

This inequality, together with (6.3) and (6.7), gives

$$\begin{aligned} & \limsup_{r \rightarrow +\infty} \frac{1}{r^n} \left(\int_{Q_r(\frac{rx}{\rho})} f(y, \nabla \tilde{v}_r^\rho) dy + \int_{S_{\tilde{v}_r^\rho} \cap Q_r(\frac{rx}{\rho})} \frac{r}{\rho} g(y, \frac{\rho}{r} [\tilde{v}_r^\rho], v_{\tilde{v}_r^\rho}) d\mathcal{H}^{n-1} \right) \\ & \leq f_{\text{hom}}(\xi) + 2\eta + C\lambda(2\rho M_\eta). \end{aligned}$$

Given $\varepsilon > 0$ and $\rho > 0$, we choose $r = \rho/\varepsilon$ and for every $y \in \mathbb{R}^n$ we define $u_\varepsilon^\rho(y) := \varepsilon \tilde{v}_r^\rho(\frac{y}{\varepsilon}) = \frac{\rho}{r} \tilde{v}_r^\rho(\frac{ry}{\rho})$. Then $u_\varepsilon^\rho \in SBV_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, $u_\varepsilon^\rho = \ell_\xi$ near $\partial Q_\rho(x)$ and in $\mathbb{R}^n \setminus Q_\rho(x)$. By a change of variables, from the previous inequality we get that for every $\rho > 0$

$$\limsup_{\varepsilon \rightarrow 0^+} \frac{E_\varepsilon(u_\varepsilon^\rho, Q_\rho(x))}{\rho^n} \leq f_{\text{hom}}(\xi) + 2\eta + C\lambda(2\rho M_\eta). \tag{6.9}$$

Since the functions u_ε^ρ coincide with ℓ_ξ in $Q_{(1+\eta)\rho}(x) \setminus Q_{(1-\delta_\varepsilon)\rho}(x)$ for some $\delta_\varepsilon \in (0, 1)$, by (f4) we have $E_\varepsilon(u_\varepsilon^\rho, Q_{(1+\eta)\rho}(x)) \leq E_\varepsilon(u_\varepsilon^\rho, Q_\rho(x)) + (c_3|\xi| + c_4)2^n \rho^n \eta$, which, together with (6.9), gives

$$\limsup_{\varepsilon \rightarrow 0^+} \frac{E_\varepsilon(u_\varepsilon^\rho, Q_{(1+\eta)\rho}(x))}{\rho^n} \leq f_{\text{hom}}(\xi) + C\lambda(2\rho M_\eta) + \tilde{K}\eta, \tag{6.10}$$

where $\tilde{K} := 2 + (c_3|\xi| + c_4)2^n$. Since u_ε^ρ coincides with ℓ_ξ in $\mathbb{R}^n \setminus Q_\rho(x)$, using Poincaré’s inequality and the lower bounds (f3) and (g3) we deduce from (6.10) that for every $\rho > 0$ there exists $\varepsilon(\rho) > 0$ such that the sequence (u_ε^ρ) is bounded in $BV_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$. In particular it is bounded in $BV(Q_\rho(x), \mathbb{R}^m)$ uniformly with respect to $\varepsilon \in (0, \varepsilon(\rho))$. Note that the bound on the L^1 -norm can be obtained by e.g. [5, Theorem 3.47]. Then there exists a subsequence, not relabelled, of the sequence (ε_j) considered in Theorem 5.1, such that $(u_{\varepsilon_j}^\rho)$ converges in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ to some $u^\rho \in BV_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ with $u^\rho = \ell_\xi$ in $Q_{(1+\eta)\rho}(x) \setminus Q_\rho(x)$. As a consequence of the Γ -convergence of $E_{\varepsilon_j}(\cdot, Q_{(1+\eta)\rho}(x))$ to $\tilde{E}(\cdot, Q_{(1+\eta)\rho}(x))$, from (6.10) we obtain

$$\begin{aligned} \frac{m_{\widehat{E}}(\ell_{\xi}, Q_{(1+\eta)\rho}(x))}{\rho^n} &\leq \frac{\widehat{E}(u^\rho, Q_{(1+\eta)\rho}(x))}{\rho^n} \leq \limsup_{j \rightarrow +\infty} \frac{E_{\varepsilon_j}(u_{\varepsilon_j}^\rho, Q_{(1+\eta)\rho}(x))}{\rho^n} \\ &\leq f_{\text{hom}}(\xi) + C\lambda(2\rho M_\eta) + \widetilde{K}\eta. \end{aligned}$$

Finally, passing to the limit as $\rho \rightarrow 0+$, from (3.8) and (6.2) we get

$$(1 + \eta)^n \frac{d\widehat{E}(u, \cdot)}{d\mathcal{L}^n}(x) \leq f_{\text{hom}}(\xi) + \widetilde{K}\eta.$$

Since $\xi := \nabla u(x)$, this gives (6.1) by the arbitrariness of $\eta > 0$.

Step 2: We claim that

$$\frac{d\widehat{E}(u, \cdot)}{d\mathcal{L}^n}(x) \geq f_{\text{hom}}(\nabla u(x)) \quad \text{for } \mathcal{L}^n\text{-a.e. } x \in A. \tag{6.11}$$

We extend u to \mathbb{R}^n by setting $u = 0$ on $\mathbb{R}^n \setminus A$. By Γ -convergence there exists $(u_\varepsilon) \subset L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, with $u_\varepsilon|_A \in SBV(A, \mathbb{R}^m)$, such that

$$u_\varepsilon \rightarrow u \quad \text{in } L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m) \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0+} E_\varepsilon(u_\varepsilon, A) = \widehat{E}(u, A), \tag{6.12}$$

along the sequence (ε_j) considered in Theorem 5.1. Passing to a further subsequence we have

$$\lim_{\varepsilon \rightarrow 0+} u_\varepsilon(x) = u(x) \tag{6.13}$$

for \mathcal{L}^n -a.e. $x \in A$. In the rest of the proof ε will always be an element of this sequence.

By (j) of Section 2 and by [5, Definition 3.70], for \mathcal{L}^n -a.e. $x \in A$ we have

$$\lim_{\rho \rightarrow 0+} \frac{1}{\rho^n} |Du|(Q_\rho(x)) = |\nabla u(x)| < +\infty, \tag{6.14}$$

$$\lim_{\rho \rightarrow 0+} \frac{1}{\rho^{n+1}} \int_{Q_\rho(x)} |u(y) - u(x) - \nabla u(x)(y - x)| dx = 0, \tag{6.15}$$

$$\lim_{\rho \rightarrow 0+} \frac{\widehat{E}(u, Q_\rho(x))}{\rho^n} = \frac{d\widehat{E}(u, \cdot)}{d\mathcal{L}^n}(x). \tag{6.16}$$

Let us fix $x \in A$ such that (6.13)-(6.16) hold true.

Recalling (5.1) we have that $\widehat{E}(u, Q_\rho(x)) \leq c_3|Du|(Q_\rho(x)) + c_4\rho^n$, hence by (6.14) and (6.16) there exists $\rho_0 \in (0, 1)$ such that

$$Q_\rho(x) \subset\subset A \quad \text{and} \quad \frac{\widehat{E}(u, Q_\rho(x))}{\rho^n} \leq c_3|\nabla u(x)| + c_4 + 1 \tag{6.17}$$

for every $\rho \in (0, \rho_0)$. Since $\widehat{E}(u, \cdot)$ is a Radon measure, there exists a set $B \subset (0, \rho_0)$, with $(0, \rho_0) \setminus B$ at most countable, such that $\widehat{E}(u, \partial Q_\rho(x)) = 0$ for every $\rho \in B$. Then we have

$$\widehat{E}(u, A) = \widehat{E}(u, Q_\rho(x)) + \widehat{E}(u, A \setminus \overline{Q}_\rho(x))$$

for every $\rho \in B$. By Γ -convergence we also have

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0^+} E_\varepsilon(u_\varepsilon, Q_\rho(x)) &\geq \widehat{E}(u, Q_\rho(x)), \\ \liminf_{\varepsilon \rightarrow 0^+} E_\varepsilon(u_\varepsilon, A \setminus \overline{Q}_\rho(x)) &\geq \widehat{E}(u, A \setminus \overline{Q}_\rho(x)), \end{aligned}$$

so that by (6.12) it follows that for every $\rho \in B$

$$\lim_{\varepsilon \rightarrow 0^+} E_\varepsilon(u_\varepsilon, Q_\rho(x)) = \widehat{E}(u, Q_\rho(x)). \tag{6.18}$$

Note that for every $\rho \in B$ there exists $\varepsilon(\rho) > 0$ such that for every $\varepsilon \in (0, \varepsilon(\rho))$

$$\frac{E_\varepsilon(u_\varepsilon, Q_\rho(x))}{\rho^n} \leq \frac{\widehat{E}(u, Q_\rho(x))}{\rho^n} + \rho \leq c_3|\nabla u(x)| + c_4 + 2, \tag{6.19}$$

where in the last inequality we used (6.17).

The rest of this proof is devoted to modifying u_ε in order to construct a competitor for the minimisation problems appearing in (4.2), which defines f_{hom} . To this end, for every $\rho \in B$ and $\varepsilon > 0$ we consider the blow-up functions defined for $y \in Q := Q_1(0)$ by

$$w_\varepsilon^\rho(y) := \frac{u_\varepsilon(x + \rho y) - u_\varepsilon(x)}{\rho} \quad \text{and} \quad w^\rho(y) := \frac{u(x + \rho y) - u(x)}{\rho}.$$

Then $w_\varepsilon^\rho \in SBV(Q, \mathbb{R}^m)$ and $w^\rho \in BV(Q, \mathbb{R}^m)$. Since $u_\varepsilon \rightarrow u$ in $L^1(Q_\rho(x), \mathbb{R}^m)$ by (6.12), using (6.13) for every $\rho \in B$ we obtain

$$w_\varepsilon^\rho \rightarrow w^\rho \quad \text{in } L^1(Q, \mathbb{R}^m) \text{ as } \varepsilon \rightarrow 0^+. \tag{6.20}$$

Moreover, from (6.15) we can deduce that

$$w^\rho \rightarrow \ell_\xi \quad \text{in } L^1(Q, \mathbb{R}^m) \text{ as } \rho \rightarrow 0^+, \tag{6.21}$$

where we set $\xi := \nabla u(x)$. Therefore, by possibly reducing the values of ρ_0 and $\varepsilon(\rho)$, we may assume that

$$\|w_\varepsilon^\rho - \ell_\xi\|_{L^1(Q, \mathbb{R}^m)} \leq 1, \tag{6.22}$$

for every $\rho \in B$ and $\varepsilon \in (0, \varepsilon(\rho))$. By the definition of w_ε^ρ , a change of variables gives

$$\frac{E_\varepsilon(u_\varepsilon, Q_\rho(x))}{\rho^n} = E_\varepsilon^\rho(w_\varepsilon^\rho, Q), \tag{6.23}$$

where E_ε^ρ is the functional corresponding to the integrands $f_\varepsilon^\rho(y, \xi) := f(\frac{x+\rho y}{\varepsilon}, \xi)$ and $g_\varepsilon^\rho(y, \zeta, \nu) := \frac{1}{\rho}g(\frac{x+\rho y}{\varepsilon}, \rho\zeta, \nu)$; i.e.,

$$E_\varepsilon^\rho(w, Q) := \int_Q f(\frac{x+\rho y}{\varepsilon}, \nabla w(y)) dy + \int_{S_w \cap Q} \frac{1}{\rho}g(\frac{x+\rho y}{\varepsilon}, \rho[w](y), \nu_w(y))d\mathcal{H}^{n-1}(y)$$

for every $w \in SBV(Q, \mathbb{R}^m)$. Note that f_ε^ρ satisfies (f3) and (f4), while g_ε^ρ satisfies (g3) and (g4).

We now modify w_ε^ρ in a way such that it attains the linear boundary datum ℓ_ξ near ∂Q . To this end we apply the Fundamental Estimate [11, Proposition 3.1] to the functionals E_ε^ρ . Thus for $\eta \in (0, \frac{1}{2})$ fixed there exist a constant $L_\eta > 0$ with the following property: for every $\rho \in B$ and $\varepsilon \in (0, \varepsilon(\rho))$ there exists a cut-off function $\varphi_\varepsilon^\rho \in C_c^\infty(Q)$, with $0 \leq \varphi_\varepsilon^\rho \leq 1$ in Q , $\text{supp}(\varphi_\varepsilon^\rho) \subset Q_{1-\eta} := Q_{1-\eta}(0)$, and $\varphi_\varepsilon^\rho = 1$ in $Q_{1-2\eta} := Q_{1-2\eta}(0)$, such that, setting $\hat{w}_\varepsilon^\rho := \varphi_\varepsilon^\rho w_\varepsilon^\rho + (1 - \varphi_\varepsilon^\rho)\ell_\xi$, we have

$$E_\varepsilon^\rho(\hat{w}_\varepsilon^\rho, Q) \leq (1 + \eta)(E_\varepsilon^\rho(w_\varepsilon^\rho, Q) + E_\varepsilon^\rho(\ell_\xi, Q \setminus \overline{Q}_{1-2\eta})) + L_\eta \|w_\varepsilon^\rho - \ell_\xi\|_{L^1(Q, \mathbb{R}^m)}. \tag{6.24}$$

We note that $\hat{w}_\varepsilon^\rho = \ell_\xi$ in $Q \setminus Q_{1-\eta}$, as desired. Moreover in view of (f4) we have

$$E_\varepsilon^\rho(\ell_\xi, Q \setminus \overline{Q}_{1-2\eta}) \leq (c_3|\xi| + c_4)\mathcal{L}^n(Q \setminus Q_{1-2\eta}) \leq (c_3|\xi| + c_4)2n\eta. \tag{6.25}$$

From (6.19), (6.22), (6.23), (6.24) and (6.25) we obtain

$$E_\varepsilon^\rho(\hat{w}_\varepsilon^\rho, Q) \leq \frac{3}{2}(c_3|\xi| + c_4 + 2 + (c_3|\xi| + c_4)n) + L_\eta \tag{6.26}$$

for every $\rho \in B$ and $\varepsilon \in (0, \varepsilon(\rho))$. By the lower bounds (f3) and (g3), we deduce that the total variation of \hat{w}_ε^ρ is bounded uniformly with respect to $\rho \in B$ and $\varepsilon \in (0, \varepsilon(\rho))$. Note moreover that also the L^1 -norm of \hat{w}_ε^ρ is bounded uniformly, by (6.22). In conclusion, the sequence $(\hat{w}_\varepsilon^\rho)$ is bounded in $BV(Q, \mathbb{R}^m)$ uniformly with respect to $\rho \in B$ and $\varepsilon \in (0, \varepsilon(\rho))$.

By Lemma 4.3 there exist a constant $M_\eta > 0$ with the following property: for every $\rho \in B$ and $\varepsilon \in (0, \varepsilon(\rho))$ there exists $\tilde{w}_\varepsilon^\rho \in SBV(Q, \mathbb{R}^m) \cap L^\infty(Q, \mathbb{R}^m)$, with $\tilde{w}_\varepsilon^\rho = \ell_\xi$ near ∂Q , such that

$$\|\tilde{w}_\varepsilon^\rho\|_{L^\infty(Q, \mathbb{R}^m)} \leq M_\eta, \quad \|\tilde{w}_\varepsilon^\rho - \ell_\xi\|_{L^1(Q, \mathbb{R}^m)} \leq \|\hat{w}_\varepsilon^\rho - \ell_\xi\|_{L^1(Q, \mathbb{R}^m)}, \tag{6.27}$$

$$E_\varepsilon^\rho(\tilde{w}_\varepsilon^\rho, Q) \leq E_\varepsilon^\rho(\hat{w}_\varepsilon^\rho, Q) + \eta.$$

We now set $r := \frac{\rho}{\varepsilon}$ and $v_\varepsilon^\rho(y) := r\tilde{w}_\varepsilon^\rho(\frac{y}{r} - \frac{x}{\rho}) + \frac{r}{\rho}\ell_\xi(x)$; clearly $v_\varepsilon^\rho \in SBV(Q_r(\frac{rx}{\rho}), \mathbb{R}^m)$, $v_\varepsilon^\rho = \ell_\xi$ near $\partial Q_r(\frac{rx}{\rho})$, and, by a change of variables

$$\| [v_\varepsilon^\rho] \|_{L^\infty(S_{v_\varepsilon^\rho} \cap Q_r(\frac{rx}{\rho}), \mathbb{R}^m)} \leq 2M_\eta r, \tag{6.28}$$

$$\frac{1}{r^n} \left(\int_{Q_r(\frac{rx}{\rho})} f(y, \nabla v_\varepsilon^\rho) dy + \int_{S_{v_\varepsilon^\rho} \cap Q_r(\frac{rx}{\rho})} \frac{r}{\rho} g(y, \frac{\rho}{r}[v_\varepsilon^\rho], v_{v_\varepsilon^\rho}) d\mathcal{H}^{n-1} \right) = E_\varepsilon^\rho(\tilde{w}_\varepsilon^\rho, Q). \tag{6.29}$$

Moreover, by combining (6.26), (6.27), and (6.29) with the lower bound (g3) we immediately deduce the existence of a constant $C > 0$, depending on $|\xi|$, such that

$$\frac{1}{r^n} \int_{S_{v_\varepsilon^\rho} \cap Q_r(\frac{rx}{\rho})} |[v_\varepsilon^\rho]| d\mathcal{H}^{n-1} \leq C. \tag{6.30}$$

By Lemma 6.1, applied with $t = \rho/r$, using (6.28) and (6.30) we deduce that

$$\frac{1}{r^n} \int_{S_{v_\varepsilon^\rho} \cap Q_r(\frac{rx}{\rho})} \left| g_0(y, [v_\varepsilon^\rho], v_{v_\varepsilon^\rho}) - \frac{r}{\rho} g(y, \frac{\rho}{r}[v_\varepsilon^\rho], v_{v_\varepsilon^\rho}) \right| d\mathcal{H}^{n-1} \leq C\lambda(2\rho M_\eta)$$

for every $\rho \in B$ and $\varepsilon \in (0, \varepsilon(\rho))$. From this inequality and from (6.23), (6.24), (6.25), (6.27), and (6.29) we obtain

$$\begin{aligned} \frac{E^{f, g_0}(v_\varepsilon^\rho, Q_r(\frac{rx}{\rho}))}{r^n} &\leq (1 + \eta) \frac{E_\varepsilon(u_\varepsilon, Q_\rho(x))}{\rho^n} + K\eta + C\lambda(2\rho M_\eta) \\ &\quad + L_\eta \|w_\varepsilon^\rho - \ell_\xi\|_{L^1(Q, \mathbb{R}^m)}, \end{aligned}$$

where $K := (c_3|\xi| + c_4)3n + 1$. Recalling that $v_\varepsilon^\rho = \ell_\xi$ near $\partial Q_r(\frac{rx}{\rho})$, we get

$$\begin{aligned} \frac{m^{f, g_0}(\ell_\xi, Q_r(\frac{rx}{\rho}))}{r^n} &\leq (1 + \eta) \frac{E_\varepsilon(u_\varepsilon, Q_\rho(x))}{\rho^n} + K\eta + C\lambda(2\rho M_\eta) \\ &\quad + L_\eta \|w_\varepsilon^\rho - \ell_\xi\|_{L^1(Q, \mathbb{R}^m)}. \end{aligned}$$

Since $r = \frac{\rho}{\varepsilon}$, by (4.2) with x replaced by $\frac{x}{\rho}$, the left-hand side of the previous inequality converges to $f_{\text{hom}}(\xi)$ as $\varepsilon \rightarrow 0+$. By (6.18) and (6.20) we can pass to the limit in the right-hand side as $\varepsilon \rightarrow 0+$ and we obtain

$$f_{\text{hom}}(\xi) \leq (1 + \eta) \frac{\widehat{E}(u, Q_\rho(x))}{\rho^n} + K\eta + C\lambda(2\rho M_\eta) + L_\eta \|w^\rho - \ell_\xi\|_{L^1(Q, \mathbb{R}^m)}.$$

By (3.8), (6.16), and (6.21), passing to the limit as $\rho \rightarrow 0+$ we get

$$f_{\text{hom}}(\xi) \leq (1 + \eta) \frac{d\widehat{E}(u, \cdot)}{d\mathcal{L}^n}(x) + K\eta.$$

Since $\xi = \nabla u(x)$, this inequality gives (6.11) by the arbitrariness of $\eta > 0$. □

7 Identification of the surface term

In Proposition 7.2 below we characterise the derivative of $\widehat{E}(u, \cdot)$ with respect to the measure $\mathcal{H}^{n-1} \llcorner S_u$ for a given BV -function u . In order to prove this result we need the estimate established in the following lemma.

Lemma 7.1 *Let $f \in \mathcal{F}$, $A \in \mathcal{A}$, $v \in BV(A, \mathbb{R}^m)$. Then for every $t > 0$*

$$\int_A |f^\infty(x, \nabla v) - \frac{1}{t} f(x, t\nabla v)| dx \leq \frac{1}{t} c_5 (1 + c_4^{1-\alpha}) \mathcal{L}^n(A) + \frac{1}{t^\alpha} c_5 c_3^{1-\alpha} (\mathcal{L}^n(A))^\alpha \|\nabla v\|_{L^1(A, \mathbb{R}^m)}^{1-\alpha}.$$

Proof Let $t > 0$. By (f4) and (3.2), using Hölder’s inequality, we obtain that

$$\begin{aligned} \int_A |f^\infty(x, \nabla v) - \frac{1}{t} f(x, t\nabla v)| dx &\leq \frac{c_5}{t} \mathcal{L}^n(A) + \frac{c_5}{t} \int_A f(x, t\nabla v)^{1-\alpha} dx \\ &\leq \frac{c_5}{t} \mathcal{L}^n(A) + \frac{c_5}{t} (\mathcal{L}^n(A))^\alpha \left(\int_A f(x, t\nabla v) dy \right)^{1-\alpha} \\ &\leq \frac{1}{t} c_5 (1 + c_4^{1-\alpha}) \mathcal{L}^n(A) + \frac{1}{t^\alpha} c_5 c_3^{1-\alpha} (\mathcal{L}^n(A))^\alpha \|\nabla v\|_{L^1(A, \mathbb{R}^m)}^{1-\alpha}. \end{aligned}$$

This concludes the proof. □

Proposition 7.2 (Homogenised surface integrand) *Let $f, g, E_\varepsilon, (\varepsilon_j)$, and \widehat{E} be as in Theorem 5.1. Assume that (b) of Theorem 4.1 holds, and let g_{hom} be as in (4.3). Then for every $A \in \mathcal{A}$ and every $u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, with $u|_A \in BV(A, \mathbb{R}^m)$, we have that*

$$\frac{d\widehat{E}(u, \cdot)}{d\mathcal{H}^{n-1} \llcorner S_u}(x) = g_{\text{hom}}([u](x), v_u(x)) \text{ for } \mathcal{H}^{n-1}\text{-a.e. } x \in S_u \cap A.$$

Proof Let us fix $A \in \mathcal{A}$ and $u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, with $u|_A \in BV(A, \mathbb{R}^m)$. We divide the proof into two steps.

Step 1: We claim that

$$\frac{d\widehat{E}(u, \cdot)}{d\mathcal{H}^{n-1} \llcorner S_u}(x) \leq g_{\text{hom}}([u](x), v_u(x)) \text{ for } \mathcal{H}^{n-1}\text{-a.e. } x \in S_u \cap A. \tag{7.1}$$

By (a)-(e) of Theorem 5.1 and by [10, Lemmas 3.1 and 3.5], arguing as in the proof of (3.17) in [10, Theorem 3.7], for \mathcal{H}^{n-1} -a.e. $x \in S_u \cap A$ we have

$$\frac{d\widehat{E}(u, \cdot)}{d\mathcal{H}^{n-1} \llcorner S_u}(x) = \lim_{\rho \rightarrow 0^+} \frac{m_{\widehat{E}}(u_{x, [u](x), v_u(x)}, Q_\rho^{v_u(x)}(x))}{\rho^{n-1}}. \tag{7.2}$$

Fix $x \in S_u \cap A$ such that (7.2) holds and set $\zeta := [u](x)$ and $v := v_u(x)$.

For every $\rho > 0$ we have

$$g_{\text{hom}}(\zeta, \nu) = \lim_{r \rightarrow +\infty} \frac{m^{f^\infty, g}(u_{\frac{rx}{\rho}, \zeta, \nu}, Q_r^v(\frac{rx}{\rho}))}{r^{n-1}}, \tag{7.3}$$

since the above identity directly follows from (4.3) by replacing x with $\frac{x}{\rho}$.

Let us fix $\eta \in (0, 1)$. By (3.4) for every $\rho \in (0, 1)$ and $r > 0$ there exists $v_r^\rho \in SBV(Q_r^v(\frac{rx}{\rho}), \mathbb{R}^m)$, with $v_r^\rho = u_{\frac{rx}{\rho}, \zeta, \nu}$ near $\partial Q_r^v(\frac{rx}{\rho})$, such that

$$\begin{aligned} E^{f^\infty, g}(v_r^\rho, Q_r^v(\frac{rx}{\rho})) &\leq m^{f^\infty, g}(u_{\frac{rx}{\rho}, \zeta, \nu}, Q_r^v(\frac{rx}{\rho})) + \eta r^{n-1} \\ &\leq \int_{\Pi_{\frac{rx}{\rho}}^v \cap Q_r^v(\frac{rx}{\rho})} g(y, \zeta, \nu) d\mathcal{H}^{n-1} + \eta r^{n-1} \leq (c_3|\zeta| + 1)r^{n-1}, \end{aligned} \tag{7.4}$$

where we used that $f^\infty(\cdot, 0) = 0$ and (g4). We extend v_r^ρ to \mathbb{R}^n by setting $v_r^\rho(y) = u_{\frac{rx}{\rho}, \zeta, \nu}(y)$ for every $y \in \mathbb{R}^n \setminus Q_r^v(\frac{rx}{\rho})$.

By combining (7.4) with the lower bound (f3) we immediately deduce the existence of a constant $C > 0$, depending on ζ , such that

$$\frac{1}{r^{n-1}} \int_{Q_r^v(\frac{rx}{\rho})} |\nabla v_r^\rho| dy \leq C. \tag{7.5}$$

By Lemma 7.1, applied with $t = r/\rho$, using (7.5), we deduce that

$$\frac{1}{r^{n-1}} \int_{Q_r^v(\frac{rx}{\rho})} |f^\infty(x, \nabla v_r^\rho) - \frac{\rho}{r} f(x, \frac{r}{\rho} \nabla v_r^\rho)| dy \leq c_5(1 + c_4^{1-\alpha})\rho + c_5c_3^{1-\alpha} \rho^\alpha C^{1-\alpha}.$$

This inequality, together with (7.3) and (7.4), gives

$$\begin{aligned} \limsup_{r \rightarrow +\infty} \frac{1}{r^{n-1}} &\left(\int_{Q_r^v(\frac{rx}{\rho})} \frac{\rho}{r} f(y, \frac{r}{\rho} \nabla v_r^\rho) dy + \int_{S_{v_r^\rho} \cap Q_r^v(\frac{rx}{\rho})} g(y, [v_r^\rho], \nu_{v_r^\rho}) d\mathcal{H}^{n-1} \right) \\ &\leq g_{\text{hom}}(\zeta, \nu) + \eta + K\rho^\alpha, \end{aligned}$$

where $K := c_5(1 + c_4^{1-\alpha}) + c_5c_3^{1-\alpha} C^{1-\alpha}$.

Given $\varepsilon > 0$ and $\rho \in (0, 1)$, for every $y \in \mathbb{R}^n$ we define $u_\varepsilon^\rho(y) := v_r^\rho(\frac{y}{\varepsilon}) = v_r^\rho(\frac{rx}{\rho})$, with $r := \rho/\varepsilon$. Then $u_\varepsilon^\rho \in SBV_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, $u_\varepsilon^\rho = u_{x, \zeta, \nu}$ near $\partial Q_\rho^v(x)$ and in $\mathbb{R}^n \setminus Q_\rho^v(x)$. By a change of variables, from the previous inequality we get that for every $\rho \in (0, 1)$

$$\limsup_{\varepsilon \rightarrow 0+} \frac{E_\varepsilon(u_\varepsilon^\rho, Q_\rho^v(x))}{\rho^{n-1}} \leq g_{\text{hom}}(\zeta, \nu) + \eta + K\rho^\alpha. \tag{7.6}$$

Since the functions u_ε^ρ coincide with $u_{x,\zeta,v}$ in $Q_{(1+\eta)\rho}^v(x) \setminus Q_{(1-\delta_\varepsilon)\rho}^v(x)$ for some $\delta_\varepsilon \in (0, 1)$, by (g4) we have $E_\varepsilon(u_\varepsilon^\rho, Q_{(1+\eta)\rho}^v(x)) \leq E_\varepsilon(u_\varepsilon^\rho, Q_\rho^v(x)) + c_3|\zeta|2^{n-1}\rho^{n-1}\eta$, which, together with (7.6), gives

$$\limsup_{\varepsilon \rightarrow 0+} \frac{E_\varepsilon(u_\varepsilon^\rho, Q_{(1+\eta)\rho}^v(x))}{\rho^{n-1}} \leq g_{\text{hom}}(\zeta, v) + K\rho^\alpha + \tilde{K}\eta, \tag{7.7}$$

where $\tilde{K} := 1 + c_3|\zeta|2^{n-1}$. Since u_ε^ρ coincides with $u_{x,\zeta,v}$ in $\mathbb{R}^n \setminus Q_\rho^v(x)$, using the lower bounds (f3) and (g3) and Poincaré’s inequality we deduce from (7.7) that for every $\rho > 0$ there exists $\varepsilon(\rho) > 0$ such that the functions u_ε^ρ are bounded in $BV_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ uniformly with respect to $\varepsilon \in (0, \varepsilon(\rho))$. Then there exists a subsequence, not relabelled, of the sequence (ε_j) considered in Theorem 5.1, such that $(u_{\varepsilon_k}^\rho)$ converges in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ to some $u^\rho \in BV_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ with $u^\rho = u_{x,\zeta,v}$ in $Q_{(1+\eta)\rho}^v(x) \setminus Q_\rho^v(x)$. As a consequence of the Γ -convergence of $E_{\varepsilon_j}(\cdot, Q_{(1+\eta)\rho}^v(x))$ to $\widehat{E}(\cdot, Q_{(1+\eta)\rho}^v(x))$, from (7.7) we obtain

$$\begin{aligned} \frac{m_{\widehat{E}}(u_{x,\zeta,v}, Q_{(1+\eta)\rho}^v(x))}{\rho^{n-1}} &\leq \frac{\widehat{E}(u^\rho, Q_{(1+\eta)\rho}^v(x))}{\rho^{n-1}} \leq \limsup_{j \rightarrow +\infty} \frac{E_{\varepsilon_j}(u_{\varepsilon_j}^\rho, Q_{(1+\eta)\rho}^v(x))}{\rho^{n-1}} \\ &\leq g_{\text{hom}}(\zeta, v) + K\rho^\alpha + \tilde{K}\eta. \end{aligned}$$

Finally, passing to the limit as $\rho \rightarrow 0+$, from (7.2) we get

$$(1 + \eta)^{n-1} \frac{d\widehat{E}(u, \cdot)}{d\mathcal{H}^{n-1} \llcorner S_u}(x) \leq g_{\text{hom}}(\zeta, v) + \tilde{K}\eta.$$

Since $\zeta := [u](x)$ and $v = v_u(x)$, this gives (7.1) by the arbitrariness of $\eta > 0$.

Step 2: We claim that

$$\frac{d\widehat{E}(u, \cdot)}{d\mathcal{H}^{n-1} \llcorner S_u}(x) \geq g_{\text{hom}}([u](x), v_u(x)) \quad \text{for } \mathcal{H}^{n-1}\text{-a.e. } x \in S_u \cap A. \tag{7.8}$$

We extend u to \mathbb{R}^n by setting $u = 0$ on $\mathbb{R}^n \setminus A$. By Γ -convergence there exists $(u_\varepsilon) \subset L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, with $u_\varepsilon|_A \in SBV(A, \mathbb{R}^m)$, such that

$$u_\varepsilon \rightarrow u \quad \text{in } L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m) \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0+} E_\varepsilon(u_\varepsilon, A) = \widehat{E}(u, A), \tag{7.9}$$

along the sequence (ε_k) considered in Theorem 5.1.

By [5, Definition 3.67 and Step 2 in the proof of Theorem 3.77] and thanks to a generalised version of the Besicovitch Differentiation Theorem (see [31] and [26, Sections 1.2.1-1.2.2]), for \mathcal{H}^{n-1} -a.e. $x \in S_u \cap A$ we have

$$\lim_{\rho \rightarrow 0+} \frac{1}{\rho^{n-1}} |Du|(Q_\rho^{v_u(x)}(x)) = |[u](x)| \neq 0, \tag{7.10}$$

$$\lim_{\rho \rightarrow 0^+} \frac{1}{\rho^n} \int_{Q_\rho^{v_u(x)}(x)} |u(y) - u_{x,[u](x),v_u(x)}(y)| dy = 0, \tag{7.11}$$

$$\lim_{\rho \rightarrow 0^+} \frac{\widehat{E}(u, Q_\rho^{v_u(x)}(x))}{\rho^{n-1}} = \frac{d\widehat{E}(u, \cdot)}{d\mathcal{H}^{n-1} \llcorner S_u}(x). \tag{7.12}$$

Let us fix $x \in S_u \cap A$ such that (7.10)-(7.12) are satisfied, and set $\zeta := [u](x)$ and $v := v_u(x)$.

Recalling (5.1) we have that $\widehat{E}(u, Q_\rho^v(x)) \leq c_3|Du|(Q_\rho^v(x)) + c_4\rho^n$, hence by (7.10) there exists $\rho_0 \in (0, 1)$ such that

$$Q_\rho^v(x) \subset\subset A \quad \text{and} \quad \frac{\widehat{E}(u, Q_\rho^v(x))}{\rho^{n-1}} \leq c_3|\zeta| + 1 \tag{7.13}$$

for every $\rho \in (0, \rho_0)$. Since $\widehat{E}(u, \cdot)$ is a Radon measure, there exists a set $B \subset (0, \rho_0)$, with $(0, \rho_0) \setminus B$ at most countable, such that $\widehat{E}(u, \partial Q_\rho^v(x)) = 0$ for every $\rho \in B$. Proceeding as in the proof of (6.18), by (7.9) we can show that for every $\rho \in B$

$$\lim_{\varepsilon \rightarrow 0^+} E_\varepsilon(u_\varepsilon, Q_\rho^v(x)) = \widehat{E}(u, Q_\rho^v(x)). \tag{7.14}$$

Hence, for every $\rho \in B$ there exists $\varepsilon(\rho) > 0$ such that for every $\varepsilon \in (0, \varepsilon(\rho))$

$$\frac{E_\varepsilon(u_\varepsilon, Q_\rho^v(x))}{\rho^{n-1}} \leq \frac{\widehat{E}(u, Q_\rho^v(x))}{\rho^{n-1}} + \rho \leq c_3|\zeta| + 2, \tag{7.15}$$

where in the last inequality we used (7.13).

The rest of this proof is devoted to modifying u_ε in order to construct a competitor for the minimisation problems appearing in (4.3), which defines g_{hom} . To this end, for every $\rho \in B$ and $\varepsilon > 0$ we consider the blow-up functions defined for $y \in Q^v := Q_1^v(0)$ by

$$w_\varepsilon^\rho(y) := u_\varepsilon(x + \rho y) \quad \text{and} \quad w^\rho(y) := u(x + \rho y).$$

Then $w_\varepsilon^\rho \in SBV(Q^v, \mathbb{R}^m)$ and $w^\rho \in BV(Q^v, \mathbb{R}^m)$. Since $u_\varepsilon \rightarrow u$ in $L^1(Q_\rho^v(x), \mathbb{R}^m)$ by (7.9), for every $\rho \in B$ we obtain

$$w_\varepsilon^\rho \rightarrow w^\rho \quad \text{in} \quad L^1(Q^v, \mathbb{R}^m) \quad \text{as} \quad \varepsilon \rightarrow 0^+; \tag{7.16}$$

moreover, from (7.11) we have

$$w^\rho \rightarrow u_{0,\zeta,v} \quad \text{in} \quad L^1(Q^v, \mathbb{R}^m) \quad \text{as} \quad \rho \rightarrow 0^+. \tag{7.17}$$

Up to possibly reducing the values of ρ_0 and $\varepsilon(\rho)$, we may then assume that

$$\|w_\varepsilon^\rho - u_{0,\zeta,v}\|_{L^1(Q^v, \mathbb{R}^m)} \leq 1, \tag{7.18}$$

for every $\rho \in B$ and $\varepsilon \in (0, \varepsilon(\rho))$. By a change of variables, we obtain the relation

$$\frac{E_\varepsilon(u_\varepsilon, Q_\rho^v(x))}{\rho^{n-1}} = E_\varepsilon^\rho(w_\varepsilon^\rho, Q^v), \tag{7.19}$$

where E_ε^ρ is the functional corresponding to the integrands $f_\varepsilon^\rho(y, \xi) := \rho f(\frac{x+\rho y}{\varepsilon}, \frac{\xi}{\rho})$ and $g_\varepsilon^\rho(y, \zeta, \nu) := g(\frac{x+\rho y}{\varepsilon}, \zeta, \nu)$; i.e.,

$$E_\varepsilon^\rho(w, Q^v) := \int_{Q^v} \rho f(\frac{x+\rho y}{\varepsilon}, \frac{1}{\rho} \nabla w(y)) dy + \int_{S_w \cap Q^v} g(\frac{x+\rho y}{\varepsilon}, [w](y), \nu_w(y)) d\mathcal{H}^{n-1}(y)$$

for every $w \in SBV(Q^v, \mathbb{R}^m)$. Note that f_ε^ρ satisfies (f3) and (f4) (recall that $\rho < 1$), while g_ε^ρ satisfies (g3) and (g4).

We now modify w_ε^ρ in a way such that it attains the boundary datum $u_{0,\zeta,\nu}$ near ∂Q^v . This will be done by applying the Fundamental Estimate [11, Proposition 3.1] to the functionals E_ε^ρ . Thus for $\eta \in (0, \frac{1}{2})$ fixed there exist a constant $L_\eta > 0$ with the following property: for every $\rho \in B$ and $\varepsilon \in (0, \varepsilon(\rho))$ there exists a cut-off function $\varphi_\varepsilon^\rho \in C_c^\infty(Q^v)$, with $0 \leq \varphi_\varepsilon^\rho \leq 1$ in Q^v , $\text{supp}(\varphi_\varepsilon^\rho) \subset Q_{1-\eta}^v := Q_{1-\eta}^v(0)$, and $\varphi_\varepsilon^\rho = 1$ in $Q_{1-2\eta}^v := Q_{1-2\eta}^v(0)$, such that, setting $\hat{w}_\varepsilon^\rho := \varphi_\varepsilon^\rho w_\varepsilon^\rho + (1 - \varphi_\varepsilon^\rho)u_{0,\zeta,\nu}$, we have

$$E_\varepsilon^\rho(\hat{w}_\varepsilon^\rho, Q^v) \leq (1 + \eta)(E_\varepsilon^\rho(w_\varepsilon^\rho, Q^v) + E_\varepsilon^\rho(u_{0,\zeta,\nu}, Q^v \setminus \overline{Q}_{1-2\eta}^v)) + L_\eta \|w_\varepsilon^\rho - u_{0,\zeta,\nu}\|_{L^1(Q^v, \mathbb{R}^m)}. \tag{7.20}$$

By definition we clearly have $\hat{w}_\varepsilon^\rho = u_{0,\zeta,\nu}$ in $Q^v \setminus Q_{1-\eta}^v$, as desired. Moreover, from (f4) and (g4) we obtain the bound

$$E_\varepsilon^\rho(u_{0,\zeta,\nu}, Q^v \setminus \overline{Q}_{1-2\eta}^v) \leq c_4 \mathcal{L}^n(Q^v \setminus Q_{1-2\eta}^v) + c_3 |\zeta| \mathcal{H}^{n-1}(\Pi_0^v \cap (Q^v \setminus \overline{Q}_{1-2\eta}^v)) \leq 2c_4 n \eta + 2c_3 |\zeta| (n - 1) \eta, \tag{7.21}$$

and hence, from (7.20) and (7.21) we have that

$$E_\varepsilon^\rho(\hat{w}_\varepsilon^\rho, Q^v) \leq (1 + \eta) E_\varepsilon^\rho(w_\varepsilon^\rho, Q^v) + K \eta + L_\eta \|w_\varepsilon^\rho - u_{0,\zeta,\nu}\|_{L^1(Q^v, \mathbb{R}^m)}, \tag{7.22}$$

for every $\rho \in B$ and every $\varepsilon \in (0, \varepsilon(\rho))$, where $K = 3(c_4 \rho n + c_3 |\zeta| (n - 1))$. Note that (f3), (g3), (7.15), (7.18), (7.19) and (7.22) imply that

$$c_2 \int_{Q^v} |\nabla \hat{w}_\varepsilon^\rho(y)| dy \leq (1 + \eta)(c_3 |\zeta| + 1) + K \eta + L_\eta, \tag{7.23}$$

for every $\rho \in B$ and $\varepsilon \in (0, \varepsilon(\rho))$.

We finally set $r := \frac{\rho}{\varepsilon}$ and $v_\varepsilon^\rho(y) := \hat{w}_\varepsilon^\rho(\frac{y}{r} - \frac{x}{\rho})$; clearly $v_\varepsilon^\rho \in SBV(Q_r^v(\frac{rx}{\rho}), \mathbb{R}^m)$, $v_\varepsilon^\rho = u_{\frac{rx}{\rho}, \zeta, \nu}$ near $\partial Q_r^v(\frac{rx}{\rho})$, and via a change of variables, using also (7.23), we have that

$$\frac{1}{r^{n-1}} \int_{Q_r^v(\frac{rx}{\rho})} |\nabla v_\varepsilon^\rho(y)| dy \leq C, \tag{7.24}$$

$$\begin{aligned} & \frac{1}{r^{n-1}} \left(\int_{Q_r^v(\frac{rx}{\rho})} \frac{\rho}{r} f(y, \frac{r}{\rho} \nabla v_\varepsilon^\rho) dy + \int_{S_{v_\varepsilon^\rho} \cap Q_r^v(\frac{rx}{\rho})} g(y, [v_r^\rho], v_{v_\varepsilon^\rho}) d\mathcal{H}^{n-1} \right) \\ & = E_\varepsilon^\rho(\hat{w}_\varepsilon^\rho, Q^v), \end{aligned} \tag{7.25}$$

where C depends only on $|\zeta|$. Hence by Lemma 7.1, applied with $t = r/\rho$, using (7.24), we have

$$\begin{aligned} \frac{1}{r^{n-1}} \int_{Q_r^v(\frac{rx}{\rho})} |f^\infty(y, \nabla v_\varepsilon^\rho) - \frac{\rho}{r} f(y, \frac{r}{\rho} \nabla v_\varepsilon^\rho)| dy & \leq c_5(1+c_4^{1-\alpha})\rho + c_5c_3^{1-\alpha} \rho^\alpha C^{1-\alpha} \\ & \leq \tilde{K} \rho^\alpha \end{aligned} \tag{7.26}$$

for every $\rho \in B$ and $\varepsilon \in (0, \varepsilon(\rho))$, where $\tilde{K} := c_5(1 + c_4^{1-\alpha}) + c_5c_3^{1-\alpha}C^{1-\alpha}$. From (7.19), (7.22), (7.25), and (7.26) we obtain

$$\begin{aligned} \frac{E^{f^\infty, g}(v_\varepsilon^\rho, Q_r^v(\frac{rx}{\rho}))}{r^{n-1}} & \leq (1 + \eta) \frac{E_\varepsilon(u_\varepsilon, Q_\rho^v(x))}{\rho^{n-1}} + K\eta + L_\eta \|w_\varepsilon^\rho - u_{0,\zeta,v}\|_{L^1(Q^v, \mathbb{R}^m)} \\ & \quad + \tilde{K} \rho^\alpha. \end{aligned}$$

Since $v_\varepsilon^\rho = u_{\frac{rx}{\rho}, \zeta, v}$ near $\partial Q_r^v(\frac{rx}{\rho})$, we have that

$$\begin{aligned} \frac{m^{f^\infty, g}(u_{\frac{rx}{\rho}, \zeta, v}, Q_r^v(\frac{rx}{\rho}))}{r^{n-1}} & \leq (1 + \eta) \frac{E_\varepsilon(u_\varepsilon, Q_\rho^v(x))}{\rho^{n-1}} + K\eta \\ & \quad + L_\eta \|w_\varepsilon^\rho - u_{0,\zeta,v}\|_{L^1(Q^v, \mathbb{R}^m)} + \tilde{K} \rho^\alpha. \end{aligned}$$

Since $r = \frac{\rho}{\varepsilon}$, by (4.3) with x replaced by $\frac{x}{\rho}$, the left-hand side converges to $g_{\text{hom}}(\xi)$ as $\varepsilon \rightarrow 0+$. By (7.14) and (7.16) we can pass to the limit in the right-hand side as $\varepsilon \rightarrow 0+$ and we obtain

$$g_{\text{hom}}(\zeta, v) \leq (1 + \eta) \frac{\widehat{E}(u, Q_\rho^v(x))}{\rho^{n-1}} + K\eta + L_\eta \|w^\rho - u_{0,\zeta,v}\|_{L^1(Q^v, \mathbb{R}^m)} + \tilde{K} \rho^\alpha.$$

By (7.12) and (7.17), passing to the limit as $\rho \rightarrow 0+$ we get

$$g_{\text{hom}}(\zeta, v) \leq (1 + \eta) \frac{d\widehat{E}(u, \cdot)}{d\mathcal{H}^{n-1} \llcorner S_u}(x) + K\eta.$$

Since $\zeta = [u](x)$ and $v = v_u(x)$, this inequality gives (7.8) by the arbitrariness of $\eta > 0$. □

8 Identification of the Cantor term

In Proposition 8.3 below we characterise the derivative of $\widehat{E}(u, \cdot)$ with respect to $|C(u)|$, the variaton of the Cantor part of the measure Du . At the end of this section we conclude the proof of Theorem 4.1.

The following proposition shows that the recession function of f_{hom} , which is defined in terms of the minimum values m^{f,g_0} , can be obtained from suitably rescaled limits of the minimum values m^{f^∞,g_0} , involving now the recession function f^∞ of f .

Proposition 8.1 *Let $f \in \mathcal{F}$ and $g \in \mathcal{G}$, and let m^{f,g_0} and m^{f^∞,g_0} be as in (3.4), with (f, g) replaced by (f, g_0) and (f^∞, g_0) , respectively. Assume that (a) of Theorem 4.1 holds, and let f_{hom} be as in (4.2). Let f_{hom}^∞ be the recession function of f_{hom} (whose existence is guaranteed by the fact that $f_{\text{hom}} \in \mathcal{F}$). Then*

$$f_{\text{hom}}^\infty(\xi) = \lim_{r \rightarrow +\infty} \frac{m^{f^\infty,g_0}(\ell_\xi, Q_r^{v,k}(rx))}{k^{n-1}r^n} \tag{8.1}$$

for every $x \in \mathbb{R}^n$, $\xi \in \mathbb{R}^{m \times n}$, $v \in \mathbb{S}^{n-1}$, and $k \in \mathbb{N}$.

Proof Let $x \in \mathbb{R}^n$, $\xi \in \mathbb{R}^{m \times n}$, $v \in \mathbb{S}^{n-1}$, $k \in \mathbb{N}$, and $\eta \in (0, 1)$ be fixed. By (3.4) for every $r > 0$ there exists $v_r \in SBV(Q_r^{v,k}(rx), \mathbb{R}^m)$, with $v_r = \ell_\xi$ near $\partial Q_r^{v,k}(rx)$, such that

$$E^{f^\infty,g_0}(v_r, Q_r^{v,k}(rx)) \leq m^{f^\infty,g_0}(\ell_\xi, Q_r^{v,k}(rx)) + \eta k^{n-1}r^n. \tag{8.2}$$

Note that, by (f3) and (f4), this implies that

$$c_2 \int_{Q_r^{v,k}(rx)} |\nabla v_r| dy \leq m^{f^\infty,g_0}(\ell_\xi, Q_r^{v,k}(rx)) + \eta k^{n-1}r^n \leq (c_3|\xi| + 1)k^{n-1}r^n, \tag{8.3}$$

where we used the fact that f^∞ satisfies (f4) with $c_4 = 0$. Let $t > 1$; by Lemma 7.1 and by (8.3), recalling that $\alpha \in (0, 1)$, we have

$$\begin{aligned} & \int_{Q_r^{v,k}(rx)} |f^\infty(y, \nabla v_r) - \frac{1}{t} f(y, t \nabla v_r)| dy \\ & \leq \frac{1}{t} c_5 (1 + c_4^{1-\alpha}) k^{n-1} r^n + \frac{1}{t^\alpha} c_5 c_3^{1-\alpha} (k^{n-1} r^n)^\alpha \|\nabla v_r\|_{L^1(Q_r^{v,k}(rx), \mathbb{R}^m)}^{1-\alpha} \leq \frac{1}{t^\alpha} K k^{n-1} r^n, \end{aligned}$$

where $K = c_5(1 + c_4^{1-\alpha}) + c_5(c_3/c_2)^{1-\alpha}(c_3|\xi| + 1)^{1-\alpha}$. Hence

$$E_{f_t,g_0}(v_r, Q_r^{v,k}(rx)) \leq E^{f^\infty,g_0}(v_r, Q_r^{v,k}(rx)) + \frac{1}{t^\alpha} K k^{n-1} r^n,$$

where $f_t(y, \xi) := \frac{1}{t} f(y, t\xi)$.

Since $v_r \in SBV(Q_r^{v,k}(rx), \mathbb{R}^m)$ and $v_r = \ell_\xi$ near $\partial Q_r^{v,k}(rx)$, the previous inequality, together with (3.4) and (8.2), gives

$$\frac{m^{f_t, g_0}(\ell_\xi, Q_r^{v,k}(rx))}{k^{n-1}r^n} \leq \frac{m^{f^\infty, g_0}(\ell_\xi, Q_r^{v,k}(rx))}{k^{n-1}r^n} + \eta + \frac{1}{t^\alpha}K. \tag{8.4}$$

We now let $r \rightarrow +\infty$ in the previous estimate. For the left-hand side we note that, by the definition of f_{hom} , (3.4), and the positive 1-homogeneity of g_0 with respect to its second variable, by a change of variables we have

$$\lim_{r \rightarrow +\infty} \frac{m^{f_t, g_0}(\ell_\xi, Q_r^{v,k}(rx))}{k^{n-1}r^n} = \lim_{r \rightarrow +\infty} \frac{m^{f, g_0}(\ell_{t\xi}, Q_r^{v,k}(rx))}{t k^{n-1}r^n} = \frac{f_{\text{hom}}(t\xi)}{t}. \tag{8.5}$$

Hence, from (8.4) and (8.5) we have that

$$\frac{f_{\text{hom}}(t\xi)}{t} \leq \liminf_{r \rightarrow +\infty} \frac{m^{f^\infty, g_0}(\ell_\xi, Q_r^{v,k}(rx))}{k^{n-1}r^n} + \eta + \frac{1}{t^\alpha}K.$$

By letting $t \rightarrow +\infty$, since $\eta \in (0, 1)$ is arbitrary, we obtain the inequality

$$f_{\text{hom}}^\infty(\xi) \leq \liminf_{r \rightarrow +\infty} \frac{m^{f^\infty, g_0}(\ell_\xi, Q_r^{v,k}(rx))}{k^{n-1}r^n}.$$

Exchanging the roles of f_t and f^∞ , an analogous argument yields the inequality

$$\limsup_{r \rightarrow +\infty} \frac{m^{f^\infty, g_0}(\ell_\xi, Q_r^{v,k}(rx))}{k^{n-1}r^n} \leq f_{\text{hom}}^\infty(\xi),$$

and hence (8.1) follows. □

For later purposes it is convenient to prove that f_{hom}^∞ can be equivalently expressed in terms of a (double) limit involving minimisation problems where the Dirichlet conditions are prescribed only on a part of the boundary. We recall that the definitions of $\partial_v^\perp Q_r^{v,k}(rx)$ and $\partial_v^\parallel Q_r^{v,k}(rx)$ are given in (i) in Section 2, while the meaning of the boundary condition on a part of the boundary is explained after (3.4).

Lemma 8.2 *Under the assumptions of Proposition 8.1 we have that*

$$\begin{aligned} f_{\text{hom}}^\infty(a \otimes v) &= \lim_{k \rightarrow +\infty} \liminf_{r \rightarrow +\infty} \frac{\tilde{m}^{f^\infty, g_0}(\ell_{a \otimes v}, Q_r^{v,k}(rx))}{k^{n-1}r^n} \\ &= \lim_{k \rightarrow +\infty} \limsup_{r \rightarrow +\infty} \frac{\tilde{m}^{f^\infty, g_0}(\ell_{a \otimes v}, Q_r^{v,k}(rx))}{k^{n-1}r^n} \end{aligned} \tag{8.6}$$

for every $x \in \mathbb{R}^n$, $a \in \mathbb{R}^m$, and $v \in \mathbb{S}^{n-1}$, where

$$\tilde{m}^{f^\infty, g_0}(\ell_{a \otimes v}, Q_r^{v,k}(rx)) := \inf\{E^{f^\infty, g_0}(v, Q_r^{v,k}(rx)) : v \in \mathcal{U}_{a,r}^{v,k}(x)\}, \tag{8.7}$$

with $U_{a,r}^{v,k}(x) := \{v \in SBV(Q_r^{v,k}(rx), \mathbb{R}^m) : v = \ell_{a \otimes v} \text{ near } \partial_v^\perp Q_r^{v,k}(rx)\}$.

Proof Let $x \in \mathbb{R}^n$, $a \in \mathbb{R}^m$, and $v \in \mathbb{S}^{n-1}$ be fixed, and for every $r > 0$ and $k > 0$ let $Q_{x,r}^{v,k} := Q_r^{v,k}(rx)$. Since $\partial_v^\perp Q_{x,r}^{v,k} \subset \partial Q_{x,r}^{v,k}$, we have $\tilde{m}^{f^\infty, g_0}(\ell_{a \otimes v}, Q_{x,r}^{v,k}) \leq m^{f^\infty, g_0}(\ell_{a \otimes v}, Q_{x,r}^{v,k})$. Due to (8.1), to obtain (8.6) we only need to prove the inequality

$$f_{\text{hom}}^\infty(a \otimes v) \leq \liminf_{k \rightarrow +\infty} \liminf_{r \rightarrow +\infty} \frac{\tilde{m}^{f^\infty, g_0}(\ell_{a \otimes v}, Q_{x,r}^{v,k})}{k^{n-1}r^n}. \tag{8.8}$$

To this aim, let us fix $h \in \mathbb{N}$. For every $r \geq 1$ and $k \in \mathbb{N}$, with $k \geq h$, there exists $v_r^k \in U_{a,r}^{v,k}(x)$ such that

$$E^{f^\infty, g_0}(v_r^k, Q_{x,r}^{v,k}) \leq \tilde{m}^{f^\infty, g_0}(\ell_{a \otimes v}, Q_{x,r}^{v,k}) + 1 \leq (c_3|a| + 1)k^{n-1}r^n, \tag{8.9}$$

where we used the fact that $f^\infty(y, \xi) \leq c_3|\xi|$ and $1 \leq k^{n-1}r^n$. By (f3) and (g3), inequality (8.9) implies that

$$c_2|Dv_r^k|(Q_{x,r}^{v,k}) \leq (c_3|a| + 1)k^{n-1}r^n. \tag{8.10}$$

Changing v_r^k in an \mathcal{L}^n -negligible set, we may assume that $v_r^k(y)$ coincides with the approximate limit of v_r^k at y for every $y \in Q_{x,r}^{v,k} \setminus S_{v_r^k}$ (see [5, Definition 3.63]). By Fubini’s theorem we have

$$\begin{aligned} \int_{k-h}^k \int_{\partial_v^\parallel Q_{x,r}^{v,\lambda}} |v_r^k - \ell_{a \otimes v}| d\mathcal{H}^{n-1} d\lambda &= \frac{2}{r} \int_{Q_{x,r}^{v,k} \setminus Q_{x,r}^{v,k-h}} |v_r^k - \ell_{a \otimes v}| dy \\ &\leq \frac{2}{r} \int_{Q_{x,r}^{v,k}} |v_r^k - \ell_{a \otimes v}| dy. \end{aligned}$$

Since $v_r^k - \ell_{a \otimes v} = 0$ near $\partial_v^\perp Q_{x,r}^{v,k}$, by Poincaré’s inequality on strips we have

$$\frac{1}{r} \int_{Q_{x,r}^{v,k}} |v_r^k - \ell_{a \otimes v}| dy \leq |Dv_r^k - a \otimes v|(Q_{x,r}^{v,k}) \leq |Dv_r^k|(Q_{x,r}^{v,k}) + |a|k^{n-1}r^n. \tag{8.11}$$

Since \mathcal{H}^{n-1} is σ -finite on $S_{v_r^k}$, we have $\mathcal{H}^{n-1}(S_{v_r^k} \cap \partial_v^\parallel Q_{x,r}^{v,\lambda}) = 0$ for \mathcal{L}^1 -a.e. $\lambda \in (k-h, k)$. From (8.10)-(8.11) we deduce that there exists $\lambda_r^k \in (k-h, k)$ such that $\mathcal{H}^{n-1}(S_{v_r^k} \cap \partial_v^\parallel Q_{x,r}^{v,\lambda_r^k}) = 0$ and

$$\int_{\partial_v^\parallel Q_{x,r}^{v,\lambda_r^k}} |v_r^k - \ell_{a \otimes v}| d\mathcal{H}^{n-1} \leq \frac{2}{h} |Dv_r^k|(Q_{x,r}^{v,k}) + \frac{2|a|}{h} k^{n-1}r^n \leq \frac{C}{h} k^{n-1}r^n. \tag{8.12}$$

where $C := 2(c_3|a| + 1)/c_2 + 2|a|$.

To prove (8.8) we need to modify v_r^k so that it attains the affine boundary datum $\ell_{a \otimes v}$ near the whole boundary $\partial Q_{x,r}^{v,k}$, and hence is a competitor for the minimisation problem in the definition of f_{hom}^∞ . The modified function is defined by

$$\hat{v}_r^k := \begin{cases} v_r^k & \text{in } Q_{x,r}^{v,\lambda_r^k}, \\ \ell_{a \otimes v} & \text{in } \mathbb{R}^n \setminus Q_{x,r}^{v,\lambda_r^k}. \end{cases}$$

Then $\hat{v}_r^k \in SBV(Q_{x,r}^{v,k}, \mathbb{R}^m)$ and $\hat{v}_r^k = \ell_{a \otimes v}$ near $\partial Q_{x,r}^{v,k}$. Moreover, since $\mathcal{H}^{n-1}(S_{v_r^k} \cap \partial_v^\parallel Q_{x,r}^{v,\lambda_r^k}) = 0$, by (f4) and (g4) we have

$$\begin{aligned} E^{f^\infty, g_0}(\hat{v}_r^k, Q_{x,r}^{v,k}) &\leq E^{f^\infty, g_0}(v_r^k, Q_{x,r}^{v,k}) + c_3|a|\mathcal{L}^n(Q_{x,r}^{v,k} \setminus Q_{x,r}^{v,\lambda_r^k}) \\ &\quad + c_3 \int_{\partial_v^\parallel Q_{x,r}^{v,\lambda_r^k}} |v_r^k - \ell_{a \otimes v}| d\mathcal{H}^{n-1}. \end{aligned} \tag{8.13}$$

Since $k^{n-1} - (\lambda_r^k)^{n-1} \leq k^{n-1} - (k-h)^{n-1} \leq (n-1)k^{n-2}h$, from (8.9), (8.12), and (8.13) we obtain

$$\begin{aligned} m^{f^\infty, g_0}(\ell_{a \otimes v}, Q_{x,r}^{v,k}) &\leq E^{f^\infty, g_0}(\hat{v}_r^k, Q_{x,r}^{v,k}) \\ &\leq \tilde{m}^{f^\infty, g_0}(\ell_{a \otimes v}, Q_{x,r}^{v,k}) + (n-1)c_3|a|k^{n-2}hr^n + \frac{c_3C}{h}k^{n-1}r^n + 1. \end{aligned}$$

We then divide both sides of the previous inequality by $k^{n-1}r^n$, to obtain

$$\frac{m^{f^\infty, g_0}(\ell_{a \otimes v}, Q_{x,r}^{v,k})}{k^{n-1}r^n} \leq \frac{\tilde{m}^{f^\infty, g_0}(\ell_{a \otimes v}, Q_{x,r}^{v,k})}{k^{n-1}r^n} + (n-1)c_3|a|\frac{h}{k} + \frac{c_3C}{h} + \frac{1}{k^{n-1}r^n}.$$

Taking the limit first as $r \rightarrow +\infty$, then as $k \rightarrow +\infty$, and finally as $h \rightarrow +\infty$, from (8.1) we obtain (8.8), and hence (8.6). □

In the next proposition we characterise the derivative of $\widehat{E}(u, \cdot)$ with respect to $|C(u)|$, for any BV-function u .

Proposition 8.3 (Homogenised Cantor integrand) *Let $f, g, E_\varepsilon, (\varepsilon_k)$, and \widehat{E} be as in Theorem 5.1. Assume that (a) of Theorem 4.1 holds, let f_{hom} be as in (4.2), and let f_{hom}^∞ denote its recession function (whose existence is guaranteed by the fact that $f_{\text{hom}} \in \mathcal{F}$). Then for every $A \in \mathcal{A}$ and every $u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, with $u|_A \in BV(A, \mathbb{R}^m)$, we have that*

$$\frac{d\widehat{E}(u, \cdot)}{d|C(u)|}(x) = f_{\text{hom}}^\infty\left(\frac{dC(u)}{d|C(u)|}(x)\right)$$

for $|C(u)|$ -a.e. $x \in A$.

Proof Let us fix $A \in \mathcal{A}$ and $u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, with $u|_A \in BV(A, \mathbb{R}^m)$. We divide the proof into two steps.

Step 1: We claim that

$$\frac{d\widehat{E}(u, \cdot)}{d|C(u)|}(x) \leq f_{\text{hom}}^\infty\left(\frac{dC(u)}{d|C(u)|}(x)\right) \text{ for } |C(u)|\text{-a.e. } x \in A. \tag{8.14}$$

By Alberti’s rank-one theorem [2] we know that for $|C(u)|$ -a.e. $x \in A$ we have

$$\frac{dC(u)}{d|C(u)|}(x) = a(x) \otimes v(x) \tag{8.15}$$

for a suitable pair $(a(x), v(x)) \in \mathbb{R}^m \times \mathbb{S}^{n-1}$. Moreover, by (a)-(e) of Theorem 5.1 and by [10, Lemma 3.9] we have that for $|C(u)|$ -a.e. $x \in A$ there exists a doubly indexed positive sequence $(t_{\rho,k})$, with $\rho > 0$ and $k \in \mathbb{N}$, such that

$$\text{for every } k \in \mathbb{N} \quad t_{\rho,k} \rightarrow +\infty \text{ and } \rho t_{\rho,k} \rightarrow 0+ \text{ as } \rho \rightarrow 0+, \tag{8.16}$$

$$\frac{d\widehat{E}(u, \cdot)}{d|C(u)|}(x) = \lim_{k \rightarrow +\infty} \limsup_{\rho \rightarrow 0+} \frac{m_{\widehat{E}}(\ell_{t_{\rho,k}a(x) \otimes v(x)}, Q_{\rho}^{v(x),k}(x))}{k^{n-1} \rho^n t_{\rho,k}}. \tag{8.17}$$

Let $x \in A$ be fixed such that (8.15)-(8.17) hold true and set $a := a(x)$ and $v := v(x)$. For every $\rho > 0$ and every $k \in \mathbb{N}$ we have

$$f_{\text{hom}}^\infty(a \otimes v) = \lim_{r \rightarrow +\infty} \frac{m^{f^\infty, g_0}(\ell_{a \otimes v}, Q_r^{v,k}(\frac{rx}{\rho}))}{k^{n-1} r^n}, \tag{8.18}$$

since the above identity directly follows from (8.1) by replacing x with $\frac{x}{\rho}$.

Let us fix $\eta \in (0, \frac{1}{2})$. By (3.4) for every $k \in \mathbb{N}$, $\rho \in (0, 1)$ and $r > 0$ there exists a function $v_r^{\rho,k} \in SBV(Q_r^{v,k}(\frac{rx}{\rho}), \mathbb{R}^m)$ with $v_r^{\rho,k} = \ell_{a \otimes v}$ near $\partial Q_r^{v,k}(\frac{rx}{\rho})$ and such that

$$\begin{aligned} E^{f^\infty, g_0}(v_r^{\rho,k}, Q_r^{v,k}(\frac{rx}{\rho})) &\leq m^{f^\infty, g_0}(\ell_{a \otimes v}, Q_r^{v,k}(\frac{rx}{\rho})) + \eta k^{n-1} r^n \\ &\leq (c_3|a| + 1)k^{n-1} r^n, \end{aligned} \tag{8.19}$$

where we used the fact that $f^\infty(y, \xi) \leq c_3|\xi|$. We extend $v_r^{\rho,k}$ to \mathbb{R}^n by setting $v_r^{\rho,k}(y) = \ell_{a \otimes v}(y)$ for every $y \in \mathbb{R}^n \setminus Q_r^{v,k}(\frac{rx}{\rho})$.

For every $y \in \mathbb{R}^n$ we define $w_r^{\rho,k}(y) := \frac{1}{r}v_r^{\rho,k}(\frac{rx}{\rho} + ry) - \ell_{a \otimes v}(\frac{x}{\rho})$. Clearly $w_r^{\rho,k} \in SBV_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ and $w_r^{\rho,k} = \ell_{a \otimes v}$ near $\partial Q_r^{v,k}$ and in $\mathbb{R}^n \setminus Q_r^{v,k}$, where $Q_r^{v,k} := Q_1^{v,k}(0)$. Moreover, by a change of variables, using the 1-homogeneity of g_0 in the second variable we have

$$E^{f_{r,\rho}, g_0^{r,\rho}}(w_r^{\rho,k}, Q^{v,k}) = \frac{1}{r^n} E^{f^\infty, g_0}(v_r^{\rho,k}, Q_r^{v,k}(\frac{rx}{\rho})), \tag{8.20}$$

where $E^{f_{r,\rho}^\infty, g_0^{r,\rho}}$ is the functional with integrands $f_{r,\rho}^\infty(y, \xi) := f^\infty(\frac{rx}{\rho} + ry, \xi)$ and $g_0^{r,\rho}(y, \zeta, \nu) := g_0(\frac{rx}{\rho} + ry, \zeta, \nu)$; *i.e.*,

$$E^{f_{r,\rho}^\infty, g_0^{r,\rho}}(w, Q^{v,k}) = \int_{Q^{v,k}} f^\infty(\frac{rx}{\rho} + ry, \nabla w(y)) dy + \int_{S_w \cap Q^{v,k}} g_0(\frac{rx}{\rho} + ry, [w](y), \nu_w(y)) d\mathcal{H}^{n-1}(y)$$

for every $w \in SBV(Q^{v,k}, \mathbb{R}^m)$. Note that $f_{r,\rho}^\infty \in \mathcal{F}$ and $g_0^{r,\rho} \in \mathcal{G}$. By the lower bounds (f3) and (g3), from (8.19) and (8.20), using also Poincaré’s inequality, we deduce that $\|w_r^{\rho,k}\|_{L^1(Q^{v,k}, \mathbb{R}^m)} + |Dw_r^{\rho,k}|(Q^{v,k}) \leq Ck^{n-1}$ for every $\rho \in (0, 1)$ and $r > 0$, with $C := 2(c_3|a| + 1)/c_2 + 2|a|$.

By Lemma 4.3 there exist a constant $M_{\eta,k} > 0$, depending also on $|a|$ and C , such that for every $\rho \in (0, 1)$ and $r > 0$ there exists $\tilde{w}_r^{\rho,k} \in SBV(Q^{v,k}, \mathbb{R}^m) \cap L^\infty(Q^{v,k}, \mathbb{R}^m)$ with the following properties: $\tilde{w}_r^{\rho,k} = \ell_{a \otimes \nu}$ near $\partial Q^{v,k}$, $\|\tilde{w}_r^{\rho,k}\|_{L^\infty(Q^{v,k}, \mathbb{R}^m)} \leq M_{\eta,k}$, and

$$E^{f_{r,\rho}^\infty, g_0^{r,\rho}}(\tilde{w}_r^{\rho,k}, Q^{v,k}) \leq E^{f_{r,\rho}^\infty, g_0^{r,\rho}}(w_r^{\rho,k}, Q^{v,k}) + \eta k^{n-1} \leq \frac{m^{f^\infty, g_0}(\ell_{a \otimes \nu}, Q_r^{v,k}(\frac{rx}{\rho}))}{r^n} + 2\eta k^{n-1}, \tag{8.21}$$

where the last inequality follows from (8.18) and (8.19). Let $\tilde{v}_r^{\rho,k} \in SBV_{loc}(\mathbb{R}^n, \mathbb{R}^m)$ be defined by $\tilde{v}_r^{\rho,k}(y) := r\tilde{w}_r^{\rho,k}(\frac{y}{r} - \frac{x}{\rho}) + \ell_{a \otimes \nu}(\frac{rx}{\rho})$. Then $\tilde{v}_r^{\rho,k} = \ell_{a \otimes \nu}$ near $\partial Q_r^{v,k}(\frac{rx}{\rho})$ and, by a change of variables,

$$\|[\tilde{v}_r^{\rho,k}]\|_{L^\infty(S_{\tilde{v}_r^{\rho,k}} \cap Q_r(\frac{rx}{\rho}), \mathbb{R}^m)} \leq 2M_{\eta,k} r, \tag{8.22}$$

$$E^{f^\infty, g_0}(\tilde{v}_r^{\rho,k}, Q_r^{v,k}(\frac{rx}{\rho})) = r^n E^{f_{r,\rho}^\infty, g_0^{r,\rho}}(\tilde{w}_r^{\rho,k}, Q^{v,k}) \leq m^{f^\infty, g_0}(\ell_{a \otimes \nu}, Q_r^{v,k}(\frac{rx}{\rho})) + 2\eta k^{n-1} r^n, \tag{8.23}$$

where the last inequality follows from (8.21). Moreover, by combining (8.19) and (8.23) with the lower bounds (f3) and (g3) we immediately deduce the existence of a constant $C > 0$, depending on $|a|$, such that

$$\frac{1}{k^{n-1} r^n} \int_{Q_r^{v,k}(\frac{rx}{\rho})} |\nabla \tilde{v}_r^{\rho,k}| dy \leq C, \tag{8.24}$$

$$\frac{1}{k^{n-1} r^n} \int_{S_{\tilde{v}_r^{\rho,k}} \cap Q_r^{v,k}(\frac{rx}{\rho})} \|[\tilde{v}_r^{\rho,k}]\| d\mathcal{H}^{n-1} \leq C. \tag{8.25}$$

Let $\rho \in (0, 1)$ and $k \in \mathbb{N}$, with $t_{\rho,k} > 1$. By Lemma 7.1, applied with $t = t_{\rho,k}$, and (8.24), recalling that $\alpha \in (0, 1)$, we obtain that

$$\begin{aligned} & \frac{1}{k^{n-1}r^n} \int_{Q_r^{v,k}(\frac{rx}{\rho})} \left| f^\infty(y, \nabla \tilde{v}_r^{\rho,k}) - \frac{1}{t_{\rho,k}} f(y, t_{\rho,k} \nabla \tilde{v}_r^{\rho,k}) \right| dy \\ & \leq \frac{1}{t_{\rho,k}} c_5(1 + c_4^{1-\alpha}) + \frac{1}{t_{\rho,k}^\alpha} c_5 c_3^{1-\alpha} C^{1-\alpha} \leq \frac{K_1}{t_{\rho,k}^\alpha}, \end{aligned} \tag{8.26}$$

where $K_1 := c_5(1 + c_4^{1-\alpha}) + c_5 c_3^{1-\alpha} C^{1-\alpha}$.

For every $r > 0$ and $\rho \in (0, 1)$ we can apply Lemma 6.1 with $t := \frac{\rho t_{\rho,k}}{r}$. By (8.22) and (8.25) we obtain

$$\begin{aligned} & \frac{1}{k^{n-1}r^n} \int_{S_{\tilde{v}_r^{\rho,k}} \cap Q_r^{v,k}(\frac{rx}{\rho})} \left| g_0(y, [\tilde{v}_r^{\rho,k}], v_{\tilde{v}_r^{\rho,k}}) - \frac{r}{\rho t_{\rho,k}} g(y, \frac{\rho t_{\rho,k}}{r} [\tilde{v}_r^{\rho,k}], v_{\tilde{v}_r^{\rho,k}}) \right| d\mathcal{H}^{n-1} \\ & \leq \lambda(2M_{\eta,k} \rho t_{\rho,k})C \end{aligned} \tag{8.27}$$

for every $r > 0$ and $\rho \in (0, 1)$.

Estimates (8.26) and (8.27), together with (8.18) and (8.23), give

$$\begin{aligned} & \limsup_{r \rightarrow +\infty} \frac{1}{k^{n-1}r^n t_{\rho,k}} \left(\int_{Q_r^{v,k}(\frac{rx}{\rho})} f(y, t_{\rho,k} \nabla \tilde{v}_r^{\rho,k}) dy \right. \\ & \quad \left. + \frac{r}{\rho} \int_{S_{\tilde{v}_r^{\rho,k}} \cap Q_r^{v,k}(\frac{rx}{\rho})} g(y, \frac{\rho t_{\rho,k}}{r} [\tilde{v}_r^{\rho,k}], v_{\tilde{v}_r^{\rho,k}}) d\mathcal{H}^{n-1} \right) \\ & \leq f_{\text{hom}}^\infty(a \otimes v) + 2\eta + K_1 t_{\rho,k}^{-\alpha} + \lambda(2M_{\eta,k} \rho t_{\rho,k})C. \end{aligned} \tag{8.28}$$

Given $k \in \mathbb{N}$ and $\rho \in (0, 1)$, for every $\varepsilon > 0$ and $y \in \mathbb{R}^n$ we define $u_\varepsilon^{\rho,k}(y) := \varepsilon t_{\rho,k} \tilde{v}_r^{\rho,k}(\frac{y}{\varepsilon}) = \frac{\rho}{r} t_{\rho,k} \tilde{v}_r^{\rho,k}(\frac{ry}{\rho})$, with $r := \rho/\varepsilon$. Then $u_\varepsilon^{\rho,k} \in SBV_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, $u_\varepsilon^{\rho,k} = t_{\rho,k} \ell_{a \otimes v}$ near $\partial Q_\rho^{v,k}(x)$ and in $\mathbb{R}^n \setminus Q_\rho^{v,k}(x)$. By a change of variables, from (8.28) we have

$$\limsup_{\varepsilon \rightarrow 0+} \frac{E_\varepsilon(u_\varepsilon^{\rho,k}, Q_\rho^{v,k}(x))}{k^{n-1} \rho^n t_{\rho,k}} \leq f_{\text{hom}}^\infty(a \otimes v) + 2\eta + K_1 t_{\rho,k}^{-\alpha} + \lambda(2M_{\eta,k} \rho t_{\rho,k})C. \tag{8.29}$$

Since $u_\varepsilon^{\rho,k}$ coincides with $t_{\rho,k} \ell_{a \otimes v}$ in $\mathbb{R}^n \setminus Q_\rho^{v,k}(x)$, using Poincaré’s inequality and the lower bounds (f3) and (g3) we deduce from (8.29) that for every $\rho > 0$ there exists $\varepsilon(\rho) > 0$ such that the functions $u_\varepsilon^{\rho,k}$ are bounded in $BV_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ uniformly with respect to $\varepsilon \in (0, \varepsilon(\rho))$. Then there exists a subsequence, not relabelled, of the sequence (ε_j) considered in Theorem 5.1, such that $(u_{\varepsilon_j}^{\rho,k})$ converges in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ to some $u^{\rho,k} \in BV_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ as $j \rightarrow +\infty$ with $u^{\rho,k} = t_{\rho,k} \ell_{a \otimes v}$ in $Q_{(1+\eta)\rho}^{v,k}(x) \setminus Q_\rho^{v,k}(x)$. As a consequence of the Γ -convergence of $E_{\varepsilon_j}(\cdot, Q_{(1+\eta)\rho}^{v,k}(x))$ to $\widehat{E}(\cdot, Q_{(1+\eta)\rho}^{v,k}(x))$, from (8.29) we obtain

$$\begin{aligned} \frac{m \widehat{E}(\ell_{t_{\rho,k} a \otimes v}, Q_{(1+\eta)\rho}^{v,k}(x))}{k^{n-1} \rho^n t_{\rho,k}} &\leq \frac{\widehat{E}(u^\rho, Q_{(1+\eta)\rho}^{v,k}(x))}{k^{n-1} \rho^n t_{\rho,k}} \leq \limsup_{j \rightarrow +\infty} \frac{E_{\varepsilon_j}(u_{\varepsilon_j}^\rho, Q_{(1+\eta)\rho}^{v,k}(x))}{k^{n-1} \rho^n t_{\rho,k}} \\ &\leq f_{\text{hom}}^\infty(a \otimes v) + 2\eta + K_1 t_{\rho,k}^{-\alpha} + \lambda(2M_{\eta,k} \rho t_{\rho,k})C. \end{aligned}$$

Now, passing to the limit as $\rho \rightarrow 0+$, from (8.16) and (3.8) we get

$$\limsup_{\rho \rightarrow 0+} \frac{m \widehat{E}(\ell_{t_{\rho,k} a \otimes v}, Q_{(1+\eta)\rho}^{v,k}(x))}{k^{n-1} \rho^n t_{\rho,k}} \leq f_{\text{hom}}^\infty(a \otimes v) + 2\eta.$$

Finally, passing to the limit as $k \rightarrow +\infty$, by (8.17) we obtain

$$(1 + \eta)^n \frac{d\widehat{E}(u, \cdot)}{d|C(u)|}(x) \leq f_{\text{hom}}^\infty(a \otimes v) + 2\eta.$$

Since $a := a(x)$ and $v = v(x)$, by (8.15) this inequality gives (8.14) by the arbitrariness of $\eta > 0$.

Step 2: We claim that

$$\frac{d\widehat{E}(u, \cdot)}{d|C(u)|}(x) \geq f_{\text{hom}}^\infty\left(\frac{dC(u)}{d|C(u)|}(x)\right) \text{ for } |C(u)|\text{-a.e. } x \in A. \tag{8.30}$$

We extend u to \mathbb{R}^n by setting $u = 0$ on $\mathbb{R}^n \setminus A$. By Γ -convergence there exists $(u_\varepsilon) \subset L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, with $u_\varepsilon|_A \in SBV(A, \mathbb{R}^m)$, such that

$$u_\varepsilon \rightarrow u \text{ in } L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m) \text{ and } \lim_{\varepsilon \rightarrow 0+} E_\varepsilon(u_\varepsilon, A) = \widehat{E}(u, A), \tag{8.31}$$

along the sequence (ε_j) considered in Theorem 5.1.

For $|C(u)|$ -a.e. $x \in A$ there exist $a(x) \in \mathbb{R}^m$ and $v(x) \in \mathbb{S}^{n-1}$ such that for every $k \in \mathbb{N}$ we have

$$\lim_{\rho \rightarrow 0+} \frac{Du(Q_\rho^{v,k}(x))}{|Du|(Q_\rho^{v,k}(x))} = \frac{dC(u)}{d|C(u)}(x) = a(x) \otimes v(x), \tag{8.32}$$

$$\lim_{\rho \rightarrow 0+} \frac{|Du|(Q_\rho^{v,k}(x))}{\rho^n} = +\infty, \tag{8.33}$$

$$\lim_{\rho \rightarrow 0+} \frac{|Du|(Q_\rho^{v,k}(x))}{\rho^{n-1}} = 0, \tag{8.34}$$

$$\lim_{\rho \rightarrow 0+} \frac{\widehat{E}(u, Q_\rho^{v,k}(x))}{|Du|(Q_\rho^{v,k}(x))} = \frac{d\widehat{E}(u, \cdot)}{d|Du|}(x) < +\infty, \tag{8.35}$$

where (8.32) follows by [2, Corollary 3.9], (8.33) and (8.34) are consequences of [4, Proposition 2.2], while (8.35) holds true thanks to a generalised version of the Besicovitch Differentiation Theorem (see [31] and [26, Sections 1.2.1-1.2.2]).

Let us fix $x \in A$ such that (8.32)-(8.35) hold true, and set $a := a(x)$ and $v := v(x)$. For $k \in \mathbb{N}$ and $\rho \in (0, 1)$ we set

$$t_{\rho,k} := \frac{|Du|(Q_\rho^{v,k}(x))}{k^{n-1}\rho^n}.$$

Then

$$\frac{d\widehat{E}(u, \cdot)}{d|C(u)|}(x) = \frac{d\widehat{E}(u, \cdot)}{d|Du|}(x) = \lim_{\rho \rightarrow 0+} \frac{\widehat{E}(u, Q_\rho^{v,k}(x))}{k^{n-1}\rho^n t_{\rho,k}}. \tag{8.36}$$

Note that, by (8.33) and (8.34), the following properties hold:

$$\text{for every } k \in \mathbb{N} \quad t_{\rho,k} \rightarrow +\infty \quad \text{and} \quad \rho t_{\rho,k} \rightarrow 0+ \quad \text{as } \rho \rightarrow 0+. \tag{8.37}$$

Recalling (5.1), we have that $\widehat{E}(u, Q_\rho^{v,k}(x)) \leq c_3|Du|(Q_\rho^{v,k}(x)) + c_4k^{n-1}\rho^n$, hence by (8.37) there exists $\rho_k \in (0, 1)$ such that

$$Q_\rho^{v,k}(x) \subset\subset A \quad \text{and} \quad \frac{\widehat{E}(u, Q_\rho^{v,k}(x))}{k^{n-1}\rho^n t_{\rho,k}} \leq c_3 + 1 \quad \text{for every } \rho \in (0, \rho_k). \tag{8.38}$$

Since $\widehat{E}(u, \cdot)$ is a Radon measure, there exists a set $B_k \subset (0, \rho_k)$, with $(0, \rho_k) \setminus B_k$ at most countable, such that $\widehat{E}(u, \partial Q_\rho^{v,k}(x)) = 0$ for every $\rho \in B_k$. Proceeding as in the proof of (6.18) and (7.14), by (8.31) we can show that for every $\rho \in B_k$

$$\lim_{\varepsilon \rightarrow 0+} E_\varepsilon(u_\varepsilon, Q_\rho^{v,k}(x)) = \widehat{E}(u, Q_\rho^{v,k}(x)). \tag{8.39}$$

Hence, for every $\rho \in B_k$ there exists $\varepsilon(\rho, k) > 0$ such that for every $\varepsilon \in (0, \varepsilon(\rho, k))$

$$\frac{E_\varepsilon(u_\varepsilon, Q_\rho^{v,k}(x))}{k^{n-1}\rho^n t_{\rho,k}} \leq \frac{\widehat{E}(u, Q_\rho^{v,k}(x))}{k^{n-1}\rho^n t_{\rho,k}} + \rho \leq c_3 + 2, \tag{8.40}$$

where in the last inequality we used (8.38).

Now, for every $\rho \in B_k$ and $\varepsilon \in (0, \varepsilon(\rho, k))$ we consider the blow-up functions defined for $y \in Q^{v,k} := Q_1^{v,k}(0)$ by

$$w_\varepsilon^{\rho,k}(y) := \frac{1}{k^{n-1}\rho t_{\rho,k}} \left(u_\varepsilon(x + \rho y) - \frac{1}{k^{n-1}\rho^n} \int_{Q_\rho^{v,k}(x)} u_\varepsilon(z) dz \right)$$

$$w^{\rho,k}(y) := \frac{1}{k^{n-1}\rho t_{\rho,k}} \left(u(x + \rho y) - \frac{1}{k^{n-1}\rho^n} \int_{Q_\rho^{v,k}(x)} u(z) dz \right).$$

Then $w_\varepsilon^{\rho,k} \in SBV(Q^{v,k}, \mathbb{R}^m)$ and $w^{\rho,k} \in BV(Q^{v,k}, \mathbb{R}^m)$. Since $u_\varepsilon \rightarrow u$ in $L^1(Q_\rho^{v,k}(x), \mathbb{R}^m)$ by (8.31), for every $\rho \in B_k$ we obtain

$$w_\varepsilon^{\rho,k} \rightarrow w^{\rho,k} \quad \text{in } L^1(Q^{v,k}, \mathbb{R}^m) \quad \text{as } \varepsilon \rightarrow 0+. \tag{8.41}$$

Moreover, recalling (8.32), we have that the function $w^{\rho,k}$ satisfies

$$\int_{Q^{v,k}} w^{\rho,k}(y) dy = 0 \quad \text{and} \quad Dw^{\rho,k}(Q^{v,k}) = \frac{Du(Q_\rho^{v,k}(x))}{|Du|(Q_\rho^{v,k}(x))} \rightarrow a \otimes v \quad \text{as } \rho \rightarrow 0+.$$

By [4, Theorem 2.3] and [29, Lemma 5.1] up to a subsequence (not relabelled),

$$w^{\rho,k} \rightarrow w^k \quad \text{in } L^1(Q^{v,k}, \mathbb{R}^m) \quad \text{as } \rho \rightarrow 0+, \tag{8.42}$$

where $w^k \in BV(Q^{v,k}, \mathbb{R}^m)$ can be represented as

$$w^k(y) = \psi^k(y \cdot v)a, \tag{8.43}$$

$$\psi^k(\frac{1}{2}) - \psi^k(-\frac{1}{2}) = \frac{1}{k^{n-1}} \quad \text{and} \quad \int_{-1/2}^{1/2} \psi^k(t) dt = 0, \tag{8.44}$$

with ψ_k nondecreasing. By monotonicity these equalities imply that $-\frac{1}{k^{n-1}} \leq \psi^k(-\frac{1}{2}) \leq 0 \leq \psi^k(\frac{1}{2}) \leq \frac{1}{k^{n-1}}$, and so

$$|\psi^k(t)| \leq \frac{1}{k^{n-1}} \quad \text{for every } t \in [-\frac{1}{2}, \frac{1}{2}]. \tag{8.45}$$

By a change of variables we obtain the equality

$$\frac{E_\varepsilon(u_\varepsilon, Q_\rho^{v,k}(x))}{k^{n-1} \rho^n t_{\rho,k}} = E_\varepsilon^{\rho,k}(w_\varepsilon^{\rho,k}, Q^{v,k}), \tag{8.46}$$

where $E_\varepsilon^{\rho,k}$ is the functional corresponding to the integrands $f_\varepsilon^{\rho,k}(y, \xi) := \frac{1}{k^{n-1} t_{\rho,k}} f(\frac{x+\rho y}{\varepsilon}, k^{n-1} t_{\rho,k} \xi)$ and $g_\varepsilon^{\rho,k}(y, \zeta, v) := \frac{1}{k^{n-1} \rho t_{\rho,k}} g(\frac{x+\rho y}{\varepsilon}, k^{n-1} \rho t_{\rho,k} \zeta, v)$; i.e.,

$$\begin{aligned} E_\varepsilon^{\rho,k}(w, Q^{v,k}) &:= \int_{Q^{v,k}} \frac{1}{k^{n-1} t_{\rho,k}} f(\frac{x+\rho y}{\varepsilon}, k^{n-1} t_{\rho,k} \nabla w(y)) dy \\ &+ \int_{S_w \cap Q^{v,k}} \frac{1}{k^{n-1} \rho t_{\rho,k}} g(\frac{x+\rho y}{\varepsilon}, k^{n-1} \rho t_{\rho,k} [w](y), v_w(y)) d\mathcal{H}^{n-1}(y) \end{aligned} \tag{8.47}$$

for every $w \in SBV(Q^{v,k}, \mathbb{R}^m)$. Note that $f_\varepsilon^{\rho,k}$ satisfies (f3) and (f4), the latter with c_4 replaced by $c_4/(k^{n-1} t_{\rho,k})$, while $g_\varepsilon^{\rho,k}$ satisfies (g3) and (g4).

Let ℓ^k be the affine function on \mathbb{R}^n which satisfies $\ell^k(y) = \psi^k(\pm \frac{1}{2})a$ for $y \cdot v = \pm \frac{1}{2}$; i.e.,

$$\ell^k(y) := \frac{1}{k^{n-1}} \ell_{a \otimes v}(y) + (\psi^k(\frac{1}{2}) - \frac{1}{2k^{n-1}})a = (\frac{1}{k^{n-1}} y \cdot v + \psi^k(\frac{1}{2}) - \frac{1}{2k^{n-1}})a \tag{8.48}$$

for every $y \in \mathbb{R}^n$. We now want to modify $w_\varepsilon^{\rho,k}$ in a way such that it attains the boundary datum ℓ^k near $\partial_v^\perp Q^{v,k}$ (see (i) in Section 2 and after (3.4)). This will be done by applying the Fundamental Estimate [11, Proposition 3.1] to the functionals $E_\varepsilon^{\rho,k}$. First note that, since by (8.37) $t_{\rho,k} \rightarrow +\infty$ as $\rho \rightarrow 0+$, we can reduce the value of the constant $\rho_k > 0$ introduced in (8.38) so that $t_{\rho,k} \geq 1$ for every $\rho \in (0, \rho_k)$.

Therefore, for $\eta \in (0, \frac{1}{2})$ fixed, by slightly modifying the proof of [11, Proposition 3.1] we obtain the following property: for every $k \in \mathbb{N}$, $\rho \in B_k$, and $\varepsilon \in (0, \varepsilon(\rho, k))$ there exists a cut-off function $\varphi_\varepsilon^{\rho,k} \in C_c^\infty(Q^{v,k})$, with $0 \leq \varphi_\varepsilon^{\rho,k} \leq 1$ in $Q^{v,k}$, $\text{supp}(\varphi_\varepsilon^{\rho,k}) \subset R_{1-\eta}^{v,k} := R_v([-\frac{k}{2}, \frac{k}{2}]^{n-1} \times [-\frac{1-\eta}{2}, \frac{1-\eta}{2}])$, and $\varphi_\varepsilon^{\rho,k} = 1$ in $R_{1-2\eta}^{v,k} := R_v([-\frac{k}{2}, \frac{k}{2}]^{n-1} \times [-\frac{1-2\eta}{2}, \frac{1-2\eta}{2}])$, such that, setting $\hat{w}_\varepsilon^{\rho,k} := \varphi_\varepsilon^{\rho,k} w_\varepsilon^{\rho,k} + (1 - \varphi_\varepsilon^{\rho,k}) \ell^k$ and $S_{2\eta}^{v,k} := Q^{v,k} \setminus \overline{R_{1-2\eta}^{v,k}}$, we have

$$E_\varepsilon^{\rho,k}(\hat{w}_\varepsilon^{\rho,k}, Q^{v,k}) \leq (1 + \eta)(E_\varepsilon^{\rho,k}(w_\varepsilon^{\rho,k}, Q^{v,k}) + E_\varepsilon^{\rho,k}(\ell^k, S_{2\eta}^{v,k})) + \frac{L}{\eta} \|w_\varepsilon^{\rho,k} - \ell^k\|_{L^1(S_{2\eta}^{v,k}, \mathbb{R}^m)}, \tag{8.49}$$

where $L > 0$ is independent of η, k, ρ , and ε . By definition we clearly have $\hat{w}_\varepsilon^{\rho,k} = \ell^k$ in $Q^{v,k} \setminus R_{1-\eta}^{v,k}$, as desired. Moreover, from the bound $f_\varepsilon^{\rho,k}(y, \xi) \leq c_3|\xi| + c_4/(k^{n-1}t_{\rho,k})$ we obtain the bound

$$E_\varepsilon^{\rho,k}(\ell^k, S_{2\eta}^{v,k}) \leq \int_{S_{2\eta}^{v,k}} (c_3|\nabla \ell^k| + \frac{c_4}{k^{n-1}t_{\rho,k}}) dy = (\frac{c_3|a|}{k^{n-1}} + \frac{c_4}{k^{n-1}t_{\rho,k}}) \mathcal{L}^n(S_{2\eta}^{v,k}) = (c_3|a| + \frac{c_4}{t_{\rho,k}}) 2\eta. \tag{8.50}$$

Hence, from (8.40), (8.46), and (8.49), using also the inequalities $\eta < \frac{1}{2}$ and $t_{\rho,k} \geq 1$, we obtain

$$E_\varepsilon^{\rho,k}(\hat{w}_\varepsilon^{\rho,k}, Q^{v,k}) \leq \frac{3}{2}(c_3 + 2 + c_3|a| + c_4) + \frac{L}{\eta} \|w_\varepsilon^{\rho,k} - \ell^k\|_{L^1(S_{2\eta}^{v,k}, \mathbb{R}^m)}. \tag{8.51}$$

We can estimate the last term in the following way:

$$\|w_\varepsilon^{\rho,k} - \ell^k\|_{L^1(S_{2\eta}^{v,k}, \mathbb{R}^m)} \leq \|w_\varepsilon^{\rho,k} - w^{\rho,k}\|_{L^1(Q^{v,k}, \mathbb{R}^m)} + \|w^{\rho,k} - w^k\|_{L^1(Q^{v,k}, \mathbb{R}^m)} + \|w^k - \ell^k\|_{L^1(S_{2\eta}^{v,k}, \mathbb{R}^m)}. \tag{8.52}$$

By (8.43) and (8.48), thanks to a change of variables we have

$$\begin{aligned} \|w^k - \ell^k\|_{L^1(S_{2\eta}^{v,k}, \mathbb{R}^m)} &= |a| \int_{S_{2\eta}^{v,k}} |\psi^k(y \cdot v) - \frac{1}{k^{n-1}} y \cdot v - \psi^k(\frac{1}{2}) + \frac{1}{2k^{n-1}}| dy \\ &= |a| k^{n-1} \int_{I_\eta \cup J_\eta} |\psi^k(t) - \frac{1}{k^{n-1}} t - \psi^k(\frac{1}{2}) + \frac{1}{2k^{n-1}}| dt, \end{aligned}$$

where $I_\eta = (-\frac{1}{2}, -\frac{1}{2} + \eta)$ and $J_\eta = (\frac{1}{2} - \eta, \frac{1}{2})$. Hence, by (8.44) and (8.45), by using the continuity of ψ^k at the endpoints, there exists a continuous function $\tau_k : [0, \frac{1}{2}] \rightarrow [0, 3]$, with $\tau_k(0) = 0$, such that

$$k^{n-1} |\psi^k(t) - \frac{1}{k^{n-1}}t - \psi^k(\frac{1}{2}) + \frac{1}{2k^{n-1}}| \leq \tau_k(\eta) \quad \text{for every } t \in I_\eta \cup J_\eta.$$

This gives

$$\|w^k - \ell^k\|_{L^1(S_{2\eta}^{v,k}, \mathbb{R}^m)} \leq 2|a|\eta\tau_k(\eta). \tag{8.53}$$

In particular, from (8.41), (8.42), (8.52), and (8.53), possibly reducing the values of the constants $\rho_k > 0$ and $\varepsilon(\rho, k) > 0$, we obtain

$$\|w_\varepsilon^{\rho,k} - \ell^k\|_{L^1(S_{2\eta}^{v,k}, \mathbb{R}^m)} \leq 1, \tag{8.54}$$

for every $\rho \in B_k$ and $\varepsilon \in (0, \varepsilon(\rho, k))$.

Note that (f3), (g3), (8.51), and (8.54) imply that the total variation of \hat{w}_ε^ρ in $Q^{v,k}$ is bounded uniformly with respect to $\rho \in B_k$ and $\varepsilon \in (0, \varepsilon(\rho, k))$. Since $\hat{w}_\varepsilon^\rho = \ell^k$ near $\partial_v^\perp Q^{v,k}$, using Poincaré’s inequality we obtain a uniform bound also for the L^1 norm of \hat{w}_ε^ρ in $Q^{v,k}$. Hence by Lemma 4.3 and Remark 4.4 there exists a constant $M_{\eta,k} > 0$ with the following property: for every $\rho \in B_k$ and $\varepsilon \in (0, \varepsilon(\rho, k))$ there exists $\tilde{w}_\varepsilon^{\rho,k} \in SBV(Q^{v,k}, \mathbb{R}^m) \cap L^\infty(Q^{v,k}, \mathbb{R}^m)$, with $\tilde{w}_\varepsilon^{\rho,k} = \ell^k$ near $\partial_v^\perp Q^{v,k}$, such that

$$\|\tilde{w}_\varepsilon^{\rho,k}\|_{L^\infty(Q^{v,k}, \mathbb{R}^m)} \leq M_{\eta,k} \quad \text{and} \quad E_\varepsilon^{\rho,k}(\tilde{w}_\varepsilon^{\rho,k}, Q^{v,k}) \leq E_\varepsilon^{\rho,k}(\hat{w}_\varepsilon^{\rho,k}, Q^{v,k}) + \eta. \tag{8.55}$$

We now set $r := \frac{\rho}{\varepsilon}$ and $v_\varepsilon^{\rho,k}(y) := r\tilde{w}_\varepsilon^{\rho,k}(\frac{y}{r} - \frac{x}{\rho}) + \frac{r}{\rho} \frac{1}{k^{n-1}} \ell_{a \otimes v}(x) - r(\psi^k(\frac{1}{2}) - \frac{1}{2k^{n-1}})a$; then $v_\varepsilon^{\rho,k} \in SBV(Q_r^{v,k}(\frac{rx}{\rho}), \mathbb{R}^m)$, $v_\varepsilon^{\rho,k} = \frac{1}{k^{n-1}} \ell_{a \otimes v}$ near $\partial_v^\perp Q_r^{v,k}(\frac{rx}{\rho})$, and, by a change of variables

$$\|[v_\varepsilon^{\rho,k}]\|_{L^\infty(S_{v_\varepsilon^{\rho,k}} \cap Q_r^{v,k}(\frac{rx}{\rho}), \mathbb{R}^m)} \leq 2M_{\eta,k} r, \tag{8.56}$$

$$\begin{aligned} & \int_{Q_r^{v,k}(\frac{rx}{\rho})} f(y, k^{n-1}t_{\rho,k} \nabla v_\varepsilon^{\rho,k}) dy \\ & + \frac{1}{\varepsilon} \int_{S_{v_\varepsilon^{\rho,k}} \cap Q_r^{v,k}(\frac{rx}{\rho})} g(y, k^{n-1}\varepsilon t_{\rho,k}[v_\varepsilon^{\rho,k}], v_{v_\varepsilon^{\rho,k}}) d\mathcal{H}^{n-1} \\ & = k^{n-1} r^n t_{\rho,k} E_\varepsilon^{\rho,k}(\tilde{w}_\varepsilon^{\rho,k}, Q^{v,k}). \end{aligned} \tag{8.57}$$

Moreover, recalling (8.47) and combining (8.51), (8.54), (8.55), and (8.57) with the lower bounds (f3) and (g3), we deduce the existence of a constant $C > 0$ such that

$$\frac{1}{r^n} \int_{Q_r^{v,k}(\frac{rx}{\rho})} |\nabla v_\varepsilon^{\rho,k}|(y) dy \leq C \tag{8.58}$$

$$\frac{1}{r^n} \int_{S_{v_\varepsilon^{\rho,k}} \cap Q_r^{v,k}(\frac{rx}{\rho})} |[v_\varepsilon^{\rho,k}]| d\mathcal{H}^{n-1} \leq C. \tag{8.59}$$

for every $\rho \in B_k$ and $\varepsilon \in (0, \varepsilon(\rho, k))$.

By Lemma 7.1, applied with $t = k^{n-1}t_{\rho,k}$, using (8.58) and the inequality $t_{\rho,k} \geq 1$ we obtain

$$\begin{aligned} & \frac{1}{r^n} \int_{Q_r^{v,k}(\frac{rx}{\rho})} \left| f^\infty(y, \nabla v_\varepsilon^{\rho,k}) - \frac{1}{k^{n-1}t_{\rho,k}} f(y, k^{n-1}t_{\rho,k} \nabla v_\varepsilon^{\rho,k}) \right| dy \\ & \leq \frac{1}{t_{\rho,k}} c_5(1 + c_4^{1-\alpha}) + \frac{1}{t_{\rho,k}^\alpha} c_5 c_3^{1-\alpha} C^{1-\alpha} \leq \frac{K_1}{t_{\rho,k}^\alpha}, \end{aligned} \tag{8.60}$$

for every $\rho \in B_k$ and $\varepsilon \in (0, \varepsilon(\rho, k))$, where, as in (8.26), $K_1 := c_5(1 + c_4^{1-\alpha}) + c_5 c_3^{1-\alpha} C^{1-\alpha}$.

Now note that, since by (8.37) $\rho t_{\rho,k} \rightarrow 0+$ as $\rho \rightarrow 0+$, we can reduce the value of the constant $\rho_k > 0$ introduced in (8.38) so that $2M_{\eta,k} k^{n-1} \rho t_{\rho,k} \leq 1$ for every $\rho \in (0, \rho_k)$. By Lemma 6.1, applied with $t := k^{n-1} \varepsilon t_{\rho,k}$, using (8.56) and (8.59), we deduce that

$$\begin{aligned} & \frac{1}{r^n} \int_{S_{v_\varepsilon^{\rho,k}} \cap Q_r^{v,k}(\frac{rx}{\rho})} \left| g_0(y, [v_\varepsilon^{\rho,k}], v_{v_\varepsilon^{\rho,k}}) - \frac{1}{k^{n-1} \varepsilon t_{\rho,k}} g(y, k^{n-1} \varepsilon t_{\rho,k} [v_\varepsilon^{\rho,k}], v_{v_\varepsilon^{\rho,k}}) \right| d\mathcal{H}^{n-1} \\ & \leq \lambda(2M_{\eta,k} k^{n-1} \rho t_{\rho,k}) \frac{1}{r^n} \int_{S_{v_\varepsilon^{\rho,k}} \cap Q_r^{v,k}(\frac{rx}{\rho})} |[v_\varepsilon^{\rho,k}]| d\mathcal{H}^{n-1} \leq \lambda(2M_{\eta,k} k^{n-1} \rho t_{\rho,k}) C, \end{aligned} \tag{8.61}$$

for every $\rho \in B_k$ and $\varepsilon \in (0, \varepsilon(\rho, k))$.

From (8.46), (8.49), (8.50), (8.54), (8.55), (8.57), (8.60), and (8.61) we obtain

$$\begin{aligned} \frac{E^{f^\infty, g_0}(v_\varepsilon^{\rho,k}, Q_r^{v,k}(\frac{rx}{\rho}))}{r^n} & \leq (1 + \eta) \frac{E_\varepsilon(u_\varepsilon, Q_\rho^{v,k}(x))}{k^{n-1} \rho^n t_{\rho,k}} + \left(\frac{3}{2} (c_3 |a| + \frac{2c_4}{t_{\rho,k}}) + 1 \right) \eta \\ & \quad + \frac{L}{\eta} \|w_\varepsilon^{\rho,k} - \ell^k\|_{L^1(S_{2\eta}^{v,k}, \mathbb{R}^m)} + \frac{K_1}{t_{\rho,k}^\alpha} + \lambda(2M_{\eta,k} k^{n-1} \rho t_{\rho,k}) C. \end{aligned} \tag{8.62}$$

By the positive 1-homogeneity of E^{f^∞, g_0} , thanks to (8.7), and recalling that $v_\varepsilon^{\rho,k} = \frac{1}{k^{n-1}} \ell_{a \otimes v}$ near $\partial_v^\perp Q_r^{v,k}(\frac{rx}{\rho})$, we have

$$\frac{\tilde{m}^{f^\infty, g_0}(\ell_{a \otimes v}, Q_r^{v,k}(\frac{rx}{\rho}))}{k^{n-1} r^n} \leq \frac{E^{f^\infty, g_0}(v_\varepsilon^{\rho,k}, Q_r^{v,k}(\frac{rx}{\rho}))}{r^n}.$$

Combining this inequality with (8.62), we obtain

$$\begin{aligned} \frac{\tilde{m}^{f^\infty, g_0}(\ell_{a \otimes v}, Q_r^{v,k}(\frac{rx}{\rho}))}{k^{n-1}r^n} &\leq (1 + \eta) \frac{E_\varepsilon(u_\varepsilon, Q_\rho^{v,k}(x))}{k^{n-1}\rho^n t_{\rho,k}} + \left(\frac{3}{2}(c_3|a| + \frac{2c_4}{t_{\rho,k}}) + 1\right)\eta \\ &\quad + \frac{L}{\eta} \|w_\varepsilon^{\rho,k} - \ell^k\|_{L^1(S_{2\eta}^{v,k}, \mathbb{R}^m)} + \frac{K_1}{t_{\rho,k}^\alpha} + \lambda(2M_{\eta,k} k^{n-1} \rho t_{\rho,k})C. \end{aligned}$$

Passing to the limit as $\varepsilon \rightarrow 0+$, recalling that $r = \frac{\rho}{\varepsilon}$, and thanks to (8.39) and (8.41), we have

$$\begin{aligned} \limsup_{r \rightarrow +\infty} \frac{\tilde{m}^{f^\infty, g_0}(\ell_{a \otimes v}, Q_r^{v,k}(\frac{rx}{\rho}))}{k^{n-1}r^n} &\leq (1 + \eta) \frac{\widehat{E}(u, Q_\rho^{v,k}(x))}{k^{n-1}\rho^n t_{\rho,k}} + \left(\frac{3}{2}(c_3|a| + \frac{2c_4}{t_{\rho,k}}) + 1\right)\eta \\ &\quad + \frac{L}{\eta} \|w^{\rho,k} - \ell^k\|_{L^1(S_{2\eta}^{v,k}, \mathbb{R}^m)} + \frac{K_1}{t_{\rho,k}^\alpha} + \lambda(2M_{\eta,k} k^{n-1} \rho t_{\rho,k})C. \end{aligned}$$

Now, passing to the limit as $\rho \rightarrow 0+$, by (3.8), (8.36), and (8.37), and (8.42),

$$\begin{aligned} \limsup_{r \rightarrow +\infty} \frac{\tilde{m}^{f^\infty, g_0}(\ell_{a \otimes v}, Q_r^{v,k}(\frac{rx}{\rho}))}{k^{n-1}r^n} &\leq (1 + \eta) \frac{d\widehat{E}(u, \cdot)}{d|C(u)|}(x) + \left(\frac{3}{2}c_3|a| + 1\right)\eta + \frac{L}{\eta} \|w^k - \ell^k\|_{L^1(S_{2\eta}^{v,k}, \mathbb{R}^m)}. \end{aligned}$$

Passing to the limit as $\eta \rightarrow 0+$, by (8.53) we deduce that

$$\limsup_{r \rightarrow +\infty} \frac{\tilde{m}^{f^\infty, g_0}(\ell_{a \otimes v}, Q_r^{v,k}(\frac{rx}{\rho}))}{k^{n-1}r^n} \leq \frac{d\widehat{E}(u, \cdot)}{d|C(u)|}(x).$$

Finally, taking the limit as $k \rightarrow +\infty$, by (8.6) we deduce that

$$f_{\text{hom}}^\infty(a \otimes v) = \lim_{k \rightarrow +\infty} \limsup_{r \rightarrow +\infty} \frac{\tilde{m}^{f^\infty, g_0}(\ell_{a \otimes v}, Q_r^{v,k}(\frac{rx}{\rho}))}{k^{n-1}r^n} \leq \frac{d\widehat{E}(u, \cdot)}{d|C(u)|}(x).$$

By (8.32), this concludes the proof of (8.30), since $a = a(x)$ and $v = v(x)$. □

We are now in a position to prove the deterministic homogenisation theorem.

Proof of Theorem 4.1 By Lemma 4.2 the function f_{hom} defined by (4.2) belongs to \mathcal{F} and by Lemma 4.5 the function g_{hom} defined by (4.3) belongs to \mathcal{G} . By Theorem 5.1 for every sequence of positive numbers converging to zero, there exist a subsequence (ε_j) and a functional $\widehat{E}: L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m) \times \mathcal{A} \rightarrow [0, +\infty]$ such that for every $A \in \mathcal{A}$ the functionals $E_{\varepsilon_j}(\cdot, A)$ Γ -converge to $\widehat{E}(\cdot, A)$ in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, as $j \rightarrow +\infty$.

Let us fix $A \in \mathcal{A}$ and $u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$. If $u|_A \in BV(A, \mathbb{R}^m)$, then by Theorem 5.1(c) and (d) the function $\widehat{E}(u, \cdot)$ is a nonnegative bounded Radon measure on $\mathcal{B}(A)$, which satisfies the inequality $\widehat{E}(u, \cdot) \leq c_3|Du| + c_4\mathcal{L}^n$. By the decomposition of the gradient of a BV function (see (f) in Section 2), the measure $\widehat{E}(u, \cdot)$ is absolutely continuous with respect to the measure $\mathcal{L}^n + \mathcal{H}^{n-1} \llcorner S_u + |C(u)|$. Since $\mathcal{L}^n, \mathcal{H}^{n-1} \llcorner S_u$, and $|C(u)|$ are carried by disjoint Borel sets, by the properties of the Radon-Nikodym derivatives mentioned in (j) of Section 2 we have

$$\widehat{E}(u, A) = \int_A \frac{d\widehat{E}(u, \cdot)}{d\mathcal{L}^n} dx + \int_{S_u \cap A} \frac{d\widehat{E}(u, \cdot)}{d\mathcal{H}^{n-1} \llcorner S_u} d\mathcal{H}^{n-1} + \int_A \frac{d\widehat{E}(u, \cdot)}{d|C(u)|} d|C(u)|.$$

Using Propositions 6.2, 7.2, and 8.3 we obtain

$$\begin{aligned} \widehat{E}(u, A) &= \int_A f_{\text{hom}}(\nabla u) dx + \int_{S_u \cap A} g_{\text{hom}}([u], \nu_u) d\mathcal{H}^{n-1} \\ &\quad + \int_A f_{\text{hom}}^\infty\left(\frac{dC(u)}{d|C(u)|}\right) d|C(u)|. \end{aligned}$$

If $u|_A \notin BV(A, \mathbb{R}^m)$, we have $\widehat{E}(u, A) = +\infty$ by Theorem 5.1(c). Therefore,

$$\widehat{E}(u, A) = E_{\text{hom}}(u, A) \quad \text{for every } u \in L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m) \text{ and every } A \in \mathcal{A},$$

where E_{hom} is the functional defined in (4.4). Since the limit does not depend on the subsequence, by the Urysohn property of Γ -convergence in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$ (see [18, Proposition 8.3]) the functionals $E_\varepsilon(\cdot, A)$ Γ -converge to $E_{\text{hom}}(\cdot, A)$ in $L^1_{\text{loc}}(\mathbb{R}^n, \mathbb{R}^m)$, as $\varepsilon \rightarrow 0+$. □

9 Stochastic homogenisation

In this section we prove Theorems 3.17 and 3.18 concerning stationary random integrands, according to Definition 3.12. We adopt the shorthand notation introduced in (3.16).

We start by proving the existence of the limits which define the the homogenised random volume integrand f_{hom} .

Proposition 9.1 (Homogenised random volume integrand) *Let f be a stationary random volume integrand and let g be a stationary random surface integrand with respect to a group $(\tau_z)_{z \in \mathbb{Z}^n}$ of P -preserving transformations on (Ω, \mathcal{T}, P) . Then there exists $\Omega' \in \mathcal{T}$, with $P(\Omega') = 1$, such that for every $\omega \in \Omega', x \in \mathbb{R}^n, \xi \in \mathbb{R}^{m \times n}, \nu \in \mathbb{S}^{n-1}, k \in \mathbb{N}$, and $\rho > 0$ the limit*

$$\lim_{r \rightarrow +\infty} \frac{m_{\omega}^{f, g_0}(\ell_{\xi}, Q_{\rho r}^{\nu, k}(rx))}{k^{n-1} \rho^n r^n} \tag{9.1}$$

exists and is independent of $x, v, k,$ and ρ . More precisely, there exists a random volume integrand $f_{\text{hom}}: \Omega \times \mathbb{R}^{m \times n} \rightarrow [0, +\infty)$ such that for every $\omega \in \Omega', x \in \mathbb{R}^n, \xi \in \mathbb{R}^{m \times n}, v \in \mathbb{S}^{n-1}, k \in \mathbb{N},$ and $\rho > 0$

$$f_{\text{hom}}(\omega, \xi) = \lim_{r \rightarrow +\infty} \frac{m_{\omega}^{f, g_0}(\ell_{\xi}, Q_{\rho r}^{v, k}(rx))}{k^{n-1} \rho^n r^n} = \lim_{r \rightarrow +\infty} \frac{m_{\omega}^{f, g_0}(\ell_{\xi}, Q_r(0))}{r^n}. \tag{9.2}$$

If, in addition, $(\tau_z)_{z \in \mathbb{Z}^n}$ is ergodic, then f_{hom} is independent of ω and

$$f_{\text{hom}}(\xi) = \lim_{r \rightarrow +\infty} \frac{1}{r^n} \int_{\Omega} m_{\omega}^{f, g_0}(\ell_{\xi}, Q_r(0)) dP(\omega).$$

Proof We divide the proof into four main steps.

Step 1: Existence of the limit in (9.1) for $\xi \in \mathbb{Q}^{m \times n}$ and $v \in \mathbb{S}^{n-1} \cap \mathbb{Q}^n$ fixed.

Let $(\Omega, \widehat{\mathcal{T}}, \widehat{P})$ denote the completion of the probability space (Ω, \mathcal{T}, P) . Let $\xi \in \mathbb{Q}^{m \times n}$ and $v \in \mathbb{S}^{n-1} \cap \mathbb{Q}^n$ be fixed. For every $\omega \in \Omega$ and $A \in \mathcal{I}_n$ (see (3.15)) we set

$$\mu_{\xi, v}(\omega, A) := \frac{1}{M_v^n} m_{\omega}^{f, g_0}(\ell_{\xi}, M_v R_v A), \tag{9.3}$$

where R_v is the orthogonal $n \times n$ matrix defined in (h) in Section 2, and M_v is a positive integer such that $M_v R_v \in \mathbb{Z}^{n \times n}$.

We now claim that the map $\mu_{\xi, v}: \Omega \times \mathcal{I}_n \rightarrow \mathbb{R}$ defines an n -dimensional subadditive process on $(\Omega, \widehat{\mathcal{T}}, \widehat{P})$, according to Definition 3.13.

We start observing that the $\widehat{\mathcal{T}}$ -measurability of $\omega \mapsto \mu_{\xi, v}(\omega, A)$ follows from the $\widehat{\mathcal{T}}$ -measurability of $\omega \mapsto m_{\omega}^{f, g_0}(\ell_{\xi}, A)$ for every $A \in \mathcal{A}$, which is ensured by Proposition A.12, taking into account Remark 3.9. We are now going to prove that $\mu_{\xi, v}$ is covariant; that is, we show that there exists a group $(\tau_z^v)_{z \in \mathbb{Z}^n}$ of \widehat{P} -preserving transformations on $(\Omega, \widehat{\mathcal{T}}, \widehat{P})$ such that

$$\mu_{\xi, v}(\omega, A + z) = \mu_{\xi, v}(\tau_z^v(\omega), A), \quad \text{for every } \omega \in \Omega, z \in \mathbb{Z}^n, \text{ and } A \in \mathcal{I}_n.$$

We have

$$M_v R_v(A + z) = M_v R_v A + M_v R_v z = M_v R_v A + z^v,$$

where $z^v := M_v R_v z \in \mathbb{Z}^n$. Then by (9.3) we get

$$\mu_{\xi, v}(\omega, A + z) = \frac{1}{M_v^n} m_{\omega}^{f, g_0}(\ell_{\xi}, M_v R_v A + z^v).$$

Given $u \in SBV(\text{int}(M_v R_v A + z^v), \mathbb{R}^m)$ with $u = \ell_{\xi}$ near $\partial(M_v R_v A + z^v)$, let $v \in SBV(\text{int}(M_v R_v A), \mathbb{R}^m)$ be defined as $v(x) := u(x + z^v) - \xi z^v$ for every $x \in \mathbb{R}^n$. By a change of variables, using the stationarity of f and g_0 we obtain

$$\int_{M_v R_v A + z^v} f(\omega, x, \nabla u) dx + \int_{S_u \cap (M_v R_v A + z^v)} g_0(\omega, x, [u], \nu_u) d\mathcal{H}^{n-1}$$

$$\begin{aligned}
 &= \int_{M_\nu R_\nu A} f(\omega, x + z_\nu, \nabla v) \, dx + \int_{S_\nu \cap (M_\nu R_\nu A)} g_0(\omega, x + z_\nu, [v], \nu_\nu) \, d\mathcal{H}^{n-1} \\
 &= \int_{M_\nu R_\nu A} f(\tau_{z^\nu}(\omega), x, \nabla v) \, dx + \int_{S_\nu \cap (M_\nu R_\nu A)} g_0(\tau_{z^\nu}(\omega), x, [v], \nu_\nu) \, d\mathcal{H}^{n-1}.
 \end{aligned}$$

Since we have $v = \ell_\xi$ near $\partial(M_\nu R_\nu A)$, we deduce that

$$m_\omega^{f, g_0}(\ell_\xi, M_\nu R_\nu A + z^\nu) = m_{\tau_{z^\nu}(\omega)}^{f, g_0}(\ell_\xi, M_\nu R_\nu A),$$

and hence the covariance of $\mu_{\xi, \nu}$ with respect to the group of \widehat{P} -preserving transformations $(\tau_z^\nu)_{z \in \mathbb{Z}^n} := (\tau_{z^\nu})_{z \in \mathbb{Z}^n}$.

We now show that $\mu_{\xi, \nu}$ is subadditive. To this end let $A \in \mathcal{I}_n$ and let $(A_i)_{i=1, \dots, N} \subset \mathcal{I}_n$ be a finite family of pairwise disjoint sets such that $A = \bigcup_{i=1}^N A_i$. For fixed $\eta > 0$ and $i = 1, \dots, N$, let $u_i \in SBV(\text{int}(M_\nu R_\nu A_i), \mathbb{R}^m)$, with $u_i = \ell_\xi$ near $\partial(M_\nu R_\nu A_i)$, be such that

$$\begin{aligned}
 &\int_{M_\nu R_\nu A_i} f(\omega, x, \nabla u_i) \, dx + \int_{S_{u_i} \cap (M_\nu R_\nu A_i)} g_0(\omega, x, [u_i], \nu_{u_i}) \, d\mathcal{H}^{n-1} \\
 &\leq m_\omega^{f, g_0}(\ell_\xi, M_\nu R_\nu A_i) + \eta
 \end{aligned}$$

and on $M_\nu R_\nu A$ define $u(x) := u_i(x)$ if $x \in M_\nu R_\nu A_i$ for $i = 1, \dots, N$. By construction we have that u is a competitor for $m_\omega^{f, g_0}(\ell_\xi, M_\nu R_\nu A)$, since $u \in SBV(M_\nu R_\nu A, \mathbb{R}^m)$ and $u = \ell_\xi$ near $\partial(M_\nu R_\nu A)$. Moreover $S_u \cap \partial(M_\nu R_\nu A_i) = \emptyset$ for every $i = 1, \dots, N$. Therefore it holds

$$\begin{aligned}
 m_\omega^{f, g_0}(\ell_\xi, M_\nu R_\nu A) &\leq \int_{M_\nu R_\nu A} f(\omega, x, \nabla u) \, dx + \int_{S_u \cap (M_\nu R_\nu A)} g_0(\omega, x, [u], \nu_u) \, d\mathcal{H}^{n-1} \\
 &= \sum_{i=1}^N \left(\int_{M_\nu R_\nu A_i} f(\omega, x, \nabla u_i) \, dx \right. \\
 &\quad \left. + \int_{S_{u_i} \cap (M_\nu R_\nu A_i)} g_0(\omega, x, [u_i], \nu_{u_i}) \, d\mathcal{H}^{n-1} \right) \\
 &\leq \sum_{i=1}^N m_\omega^{f, g_0}(\ell_\xi, M_\nu R_\nu A_i) + N\eta,
 \end{aligned}$$

thus the subadditivity of $\mu_{\xi, \nu}$ follows from (9.3), by the arbitrariness of $\eta > 0$. Note that the same proof shows that

$$m_\omega^{f, g_0} \left(\ell_\xi, \bigcup_{i=1}^N A_i \right) \leq \sum_{i=1}^N m_\omega^{f, g_0}(\ell_\xi, A_i) \tag{9.4}$$

for every finite disjoint family $(A_i)_{i=1, \dots, N} \subset \mathcal{I}_n$,

even when $\bigcup_{i=1}^N A_i \notin \mathcal{I}_n$.

Eventually, in view of (f4) we have

$$\begin{aligned} \mu_{\xi,v}(\omega, A) &= \frac{1}{M_v^n} m_{\omega}^{f,g_0}(\ell_{\xi}, M_v R_v A) \\ &\leq \frac{1}{M_v^n} \int_{M_v R_v A} f(\omega, x, \xi) dx \leq (c_3|\xi| + c_4)\mathcal{L}^n(A), \end{aligned} \tag{9.5}$$

for every $\omega \in \Omega$.

We note now that for every $x \in \mathbb{R}^n, k \in \mathbb{N}$, and $\rho > 0$, we have that $(Q_{\rho r}^{e_n,k}(rx))_{r>0}$ is a regular family in \mathcal{I}_n (cf. Definition 3.14). Therefore for every fixed $\xi \in \mathbb{Q}^n$ and $v \in \mathbb{S}^{n-1} \cap \mathbb{Q}^n$ we can apply Theorem 3.15 to the subadditive process $\mu_{\xi,v}$ on $(\Omega, \widehat{\mathcal{T}}, \widehat{P})$ to deduce the existence of a $\widehat{\mathcal{T}}$ -measurable function $\varphi_{\xi,v} : \Omega \rightarrow [0, +\infty)$ and a set $\widehat{\Omega}_{\xi,v} \subset \Omega$, with $\widehat{\Omega}_{\xi,v} \in \widehat{\mathcal{T}}$ and $P(\widehat{\Omega}_{\xi,v}) = 1$ such that

$$\lim_{r \rightarrow +\infty} \frac{\mu_{\xi,v}(\omega, Q_{\rho r}^{e_n,k}(rx))}{k^{n-1} \rho^n r^n} = \varphi_{\xi,v}(\omega), \tag{9.6}$$

for every $\omega \in \widehat{\Omega}_{\xi,v}, x \in \mathbb{R}^n, k \in \mathbb{N}$, and $\rho > 0$. Then, by the properties of the completion there exist a set $\Omega_{\xi,v} \in \mathcal{T}$, with $P(\Omega_{\xi,v}) = 1$, and a \mathcal{T} -measurable function, which we still denote by $\varphi_{\xi,v}$, such that (9.6) holds for every $\omega \in \Omega_{\xi,v}$. Thus choosing in (9.6) $x = 0, k = 1$, and $\rho = 1$, thanks to (9.3) we get

$$\varphi_{\xi,v}(\omega) = \lim_{r \rightarrow +\infty} \frac{\mu_{\xi,v}(\omega, Q_r(0))}{r^n} = \lim_{r \rightarrow +\infty} \frac{m_{\omega}^{f,g_0}(\ell_{\xi}, Q_r^v(0))}{r^n}.$$

Furthermore, if $(\tau_z)_{z \in \mathbb{Z}^n}$ is ergodic, then Theorem 3.15 ensures that $\varphi_{\xi,v}$ is independent of ω .

Step 2: Existence of the limit in (9.1) for every $\xi \in \mathbb{R}^{m \times n}$ and $v \in \mathbb{S}^{n-1}$.

Let $\widetilde{\Omega}$ denote the intersection of the sets $\Omega_{\xi,v}$ for $\xi \in \mathbb{Q}^n$ and $v \in \mathbb{S}^{n-1} \cap \mathbb{Q}^n$; clearly $\widetilde{\Omega} \in \mathcal{T}$ and $P(\widetilde{\Omega}) = 1$. For every $k \in \mathbb{N}$ and $\rho > 0$ we now introduce the auxiliary functions $\underline{f}_{\rho,k}, \overline{f}_{\rho,k} : \widetilde{\Omega} \times \mathbb{R}^n \times \mathbb{R}^{m \times n} \times \mathbb{S}^{n-1} \rightarrow [0, +\infty)$ defined as

$$\begin{aligned} \underline{f}_{\rho,k}(\omega, x, \xi, v) &:= \liminf_{r \rightarrow +\infty} \frac{m_{\omega}^{f,g_0}(\ell_{\xi}, Q_{\rho r}^{v,k}(rx))}{k^{n-1} \rho^n r^n} \\ \overline{f}_{\rho,k}(\omega, x, \xi, v) &:= \limsup_{r \rightarrow +\infty} \frac{m_{\omega}^{f,g_0}(\ell_{\xi}, Q_{\rho r}^{v,k}(rx))}{k^{n-1} \rho^n r^n}. \end{aligned}$$

We notice that

$$\underline{f}_{\rho,k}(\omega, x, \xi, v) = \overline{f}_{\rho,k}(\omega, x, \xi, v) = \varphi_{\xi,v}(\omega) \tag{9.7}$$

for every $\omega \in \widetilde{\Omega}, \xi \in \mathbb{Q}^{m \times n}, v \in \mathbb{S}^{n-1} \cap \mathbb{Q}^n, k \in \mathbb{N}$, and $\rho > 0$. The proof of property (4.8) in Lemma 4.2 can be adapted to the rectangles $Q_{\rho r}^{v,k}(rx)$, obtaining that for every

$\omega \in \tilde{\Omega}, x \in \mathbb{R}^n, v \in \mathbb{S}^{n-1}, k \in \mathbb{N}$, and $\rho > 0$, the functions $\xi \mapsto \underline{f}_{\rho,k}(\omega, x, \xi, v)$ and $\xi \mapsto \overline{f}_{\rho,k}(\omega, x, \xi, v)$ are continuous on $\mathbb{R}^{m \times n}$, and their modulus of continuity does not depend on ω, x, v, k , and ρ . By (9.7) this implies that for every $\xi \in \mathbb{R}^{m \times n}$ and $v \in \mathbb{S}^{n-1} \cap \mathbb{Q}^n$ there exists a \mathcal{T} -measurable function, which we still denote by $\varphi_{\xi,v}$, such that

$$\underline{f}_{\rho,k}(\omega, x, \xi, v) = \overline{f}_{\rho,k}(\omega, x, \xi, v) = \varphi_{\xi,v}(\omega) \tag{9.8}$$

for every $\omega \in \tilde{\Omega}, x \in \mathbb{R}^n, k \in \mathbb{N}$, and $\rho > 0$.

We now show that, for every $\omega \in \tilde{\Omega}, x \in \mathbb{R}^n, \xi \in \mathbb{R}^{m \times n}, k \in \mathbb{N}$, and $\rho > 0$, the functions $v \mapsto \underline{f}_{\rho,k}(\omega, x, \xi, v)$ and $v \mapsto \overline{f}_{\rho,k}(\omega, x, \xi, v)$, restricted to $\widehat{\mathbb{S}}_+^{n-1}$ and $\widehat{\mathbb{S}}_-^{n-1}$, are continuous. We will only prove this property for $\underline{f}_{\rho,k}$ and $\widehat{\mathbb{S}}_+^{n-1}$, the other proofs being analogous. To this end, let $\omega \in \tilde{\Omega}, x \in \mathbb{R}^n, \xi \in \mathbb{R}^{m \times n}, k \in \mathbb{N}$, and $\rho > 0$ be fixed. Let $v \in \widehat{\mathbb{S}}_+^{n-1}$ and let $(v_j) \subset \widehat{\mathbb{S}}_+^{n-1}$ be such that $v_j \rightarrow v$ as $j \rightarrow +\infty$. Since $v \mapsto R_v$ is continuous in $\widehat{\mathbb{S}}_+^{n-1}$, for every $\delta \in (0, 1/2)$ there exists an integer \hat{j} , depending on ρ, k , and δ , such that

$$Q_{(1-\delta)\rho r}^{v_j,k}(rx) \subset\subset Q_{\rho r}^{v,k}(rx) \subset\subset Q_{(1+\delta)\rho r}^{v_j,k}(rx)$$

for every $j \geq \hat{j}$ and $r > 0$. Given $r > 0, j \geq \hat{j}$, and $\eta > 0$, let $u \in SBV(Q_{\rho r}^{v,k}(rx), \mathbb{R}^m)$ be such that $u = \ell_\xi$ near $\partial Q_{\rho r}^{v,k}(rx)$ and

$$\begin{aligned} & \int_{Q_{\rho r}^{v,k}(rx)} f(\omega, y, \nabla u) dy + \int_{S_u \cap Q_{\rho r}^{v,k}(rx)} g_0(\omega, y, [u], v_u) d\mathcal{H}^{n-1} \\ & \leq m_\omega^{f,g_0}(\ell_\xi, Q_{\rho r}^{v,k}(rx)) + \eta k^{n-1} \rho^n r^n. \end{aligned}$$

We define now $v \in SBV(Q_{(1+\delta)\rho r}^{v_j,k}(rx), \mathbb{R}^m)$ as

$$v(y) = \begin{cases} u(y) & \text{if } y \in Q_{\rho r}^{v,k}(rx), \\ \ell_\xi(y) & \text{if } y \in Q_{(1+\delta)\rho r}^{v_j,k}(rx) \setminus Q_{\rho r}^{v,k}(rx). \end{cases}$$

Note that $v = \ell_\xi$ near $\partial Q_{(1+\delta)\rho r}^{v_j,k}(rx)$ and that $\partial Q_{\rho r}^{v,k}(rx) \cap S_v = \emptyset$. Therefore,

$$\begin{aligned} & m_\omega^{f,g_0}(\ell_\xi, Q_{(1+\delta)\rho r}^{v_j,k}(rx)) \\ & \leq \int_{Q_{(1+\delta)\rho r}^{v_j,k}(rx)} f(\omega, y, \nabla v) dy + \int_{S_v \cap Q_{(1+\delta)\rho r}^{v_j,k}(rx)} g_0(\omega, y, [v], v_v) d\mathcal{H}^{n-1} \\ & \leq \int_{Q_{\rho r}^{v,k}(rx)} f(\omega, y, \nabla u) dx \\ & \quad + \int_{S_u \cap Q_{\rho r}^{v,k}(rx)} g_0(\omega, y, [u], v_u) d\mathcal{H}^{n-1} + (c_3|\xi| + c_4)((1 + \delta)^n - 1)k^{n-1} \rho^n r^n \end{aligned}$$

$$\leq m_{\omega}^{f, g_0}(\ell_{\xi}, Q_{\rho r}^{v, k}(rx)) + \eta k^{n-1} \rho^n r^n + (c_3|\xi| + c_4)((1 + \delta)^n - 1)k^{n-1} \rho^n r^n.$$

Dividing by $k^{n-1} \rho^n r^n$ and passing to the liminf as $r \rightarrow +\infty$ we obtain

$$(1 + \delta)^n \underline{f}_{\rho, k}(\omega, x, \xi, v_j) \leq \underline{f}_{\rho, k}(\omega, x, \xi, v) + \eta + (c_3|\xi| + c_4)((1 + \delta)^n - 1).$$

Passing to the limsup first as $j \rightarrow +\infty$, then as $\delta \rightarrow 0+$ and $\eta \rightarrow 0+$ we get

$$\limsup_{j \rightarrow +\infty} \underline{f}_{\rho, k}(\omega, x, \xi, v_j) \leq \underline{f}_{\rho, k}(\omega, x, \xi, v).$$

A similar argument, using the cubes $Q_{(1-\delta)\rho r}^{v_j, k}(rx)$, gives

$$\underline{f}_{\rho, k}(\omega, x, \xi, v) \leq \liminf_{j \rightarrow +\infty} \underline{f}(\omega, x, \xi, v_j),$$

and so the continuity of $v \mapsto \underline{f}_{\rho, k}(\omega, x, \xi, v)$ follows.

It is known that $\mathbb{Q}^n \cap \mathbb{S}^{n-1}$ is dense in \mathbb{S}^{n-1} (see, e.g., [16, Remark A.2]). Arguing as in the proof of [16, Theorem 5.1] it is easy to show that $\mathbb{Q}^n \cap \widehat{\mathbb{S}}_{\pm}^{n-1}$ is dense in $\widehat{\mathbb{S}}_{\pm}^{n-1}$. Therefore, from the continuity property proved above and from (9.8) we deduce that for every $\xi \in \mathbb{R}^{m \times n}$ and $v \in \mathbb{S}^{n-1}$ there exists a \mathcal{T} -measurable function, which we still denote by $\varphi_{\xi, v}$, such that

$$\underline{f}_{\rho, k}(\omega, x, \xi, v) = \overline{f}_{\rho, k}(\omega, x, \xi, v) = \varphi_{\xi, v}(\omega) \tag{9.9}$$

for every $\omega \in \widetilde{\Omega}$, $x \in \mathbb{R}^n$, $\xi \in \mathbb{R}^{m \times n}$, $v \in \mathbb{S}^{n-1}$, $k \in \mathbb{N}$, and $\rho > 0$. This implies that

$$\varphi_{\xi, v}(\omega) = \lim_{r \rightarrow +\infty} \frac{m_{\omega}^{f, g_0}(\ell_{\xi}, Q_{\rho r}^{v, k}(rx))}{k^{n-1} \rho^n r^n} \tag{9.10}$$

for every $\omega \in \widetilde{\Omega}$, $x \in \mathbb{R}^n$, $\xi \in \mathbb{R}^{m \times n}$, $v \in \mathbb{S}^{n-1}$, $k \in \mathbb{N}$, and $\rho > 0$, concluding the proof of Step 2.

Step 3: The limit in (9.1) is independent of v .

We now show that $\varphi_{\xi, v}(\omega)$ does not depend on v ; i.e., we show that

$$\varphi_{\xi, v}(\omega) = \varphi_{\xi, e_n}(\omega) \quad \text{for every } \omega \in \widetilde{\Omega}, \xi \in \mathbb{R}^{m \times n}, v \in \mathbb{S}^{n-1}. \tag{9.11}$$

For every $r > 0$ let $Q_r^v := Q_r^v(0)$ and let $\eta > 0$ be fixed. Let $(Q_{\rho_i}(x_i))$ be a family of pairwise disjoint cubes, with $\rho_i \in (0, 1)$, $i = 1, \dots, N_{\eta}$, with faces parallel to the coordinate axes, such that

$$\bigcup_{i=1}^{N_{\eta}} Q_{\rho_i}(x_i) \subset Q_r^v \quad \text{and} \quad \mathcal{L}^n \left(Q_r^v \setminus \bigcup_{i=1}^{N_{\eta}} Q_{\rho_i}(x_i) \right) = 1 - \sum_{i=1}^{N_{\eta}} \rho_i^n < \eta.$$

By using arguments similar to those used to prove the subadditivity in Step 1, we can prove that

$$m_{\omega}^{f, g_0}(\ell_{\xi}, Q_r^v) \leq m_{\omega}^{f, g_0}\left(\ell_{\xi}, \bigcup_{i=1}^{N_{\eta}} Q_{\rho_i r}(rx_i)\right) + \eta(c_3|\xi| + c_4)r^n.$$

Then, by (9.4), for every $r > 0$ we have

$$\begin{aligned} \frac{m_{\omega}^{f, g_0}(\ell_{\xi}, Q_r^v)}{r^n} &\leq \sum_{i=1}^{N_{\eta}} \frac{m_{\omega}^{f, g_0}(\ell_{\xi}, Q_{\rho_i r}(rx_i))}{r^n} + \eta(c_3|\xi| + c_4) \\ &= \sum_{i=1}^{N_{\eta}} \frac{m_{\omega}^{f, g_0}(\ell_{\xi}, Q_{\rho_i r}(rx_i))}{\rho_i^n r^n} \rho_i^n + \eta(c_3|\xi| + c_4). \end{aligned}$$

Hence, passing to the limit as $r \rightarrow +\infty$ and using (9.10) we obtain

$$\varphi_{\xi, v}(\omega) \leq \varphi_{\xi, e_n}(\omega) \sum_{i=1}^{N_{\eta}} \rho_i^n + \eta(c_3|\xi| + c_4) \leq \varphi_{\xi, e_n}(\omega) + \eta(c_3|\xi| + c_4),$$

thus taking the limit as $\eta \rightarrow 0+$ we get $\varphi_{\xi, v}(\omega) \leq \varphi_{\xi, e_n}(\omega)$. By repeating a similar argument, now using coverings of $Q_1(0)$ by cubes of the form $Q_{\rho_i}^v(x_i)$, we obtain the opposite inequality, and eventually the claim.

Step 4: Definition and properties of f_{hom} .

For every $\omega \in \Omega$ and $\xi \in \mathbb{R}^{m \times n}$ we define

$$f_{\text{hom}}(\omega, \xi) := \begin{cases} \varphi_{\xi, e_n}(\omega) & \text{if } \omega \in \tilde{\Omega}, \\ c_2|\xi| & \text{if } \omega \in \Omega \setminus \tilde{\Omega}. \end{cases}$$

Then (9.2) follows from (9.10) and (9.11). From the measurability of φ_{ξ, e_n} , proved in Step 2, we obtain that $f_{\text{hom}}(\cdot, \xi)$ is \mathcal{T} -measurable in Ω for every $\xi \in \mathbb{R}^{m \times n}$. Moreover, since the function $\xi \mapsto \underline{f}_{\rho, k}(\omega, x, \xi, v)$ is continuous on $\mathbb{R}^{m \times n}$, from (9.9) we deduce that $f_{\text{hom}}(\omega, \cdot)$ is continuous in $\mathbb{R}^{m \times n}$ for every $\omega \in \Omega$, and this implies the $\mathcal{T} \otimes \mathcal{B}^{m \times n}$ -measurability of f_{hom} on $\Omega \times \mathbb{R}^{m \times n}$. Finally, Lemma 4.2 allows us to conclude that $f_{\text{hom}}(\omega, \cdot) \in \mathcal{F}$ for every $\omega \in \Omega$. Therefore, f_{hom} is a random volume integrand according to Definition 3.7. □

The following result is a direct consequence of Propositions 8.1 and 9.1. In the ergodic case, (9.13) can be obtained by integrating (9.12) and observing that, thanks to (9.5), we can apply the Dominated Convergence Theorem.

Proposition 9.2 (Homogenised random Cantor integrand) *Under the assumptions of Proposition 9.1, for every $\omega \in \Omega'$ and $\xi \in \mathbb{R}^{m \times n}$ let*

$$f_{\text{hom}}^{\infty}(\omega, \xi) := \lim_{t \rightarrow +\infty} \frac{f_{\text{hom}}(\omega, t\xi)}{t}$$

(since $f_{\text{hom}}(\omega, \cdot) \in \mathcal{F}$, the existence of the limit is guaranteed by (f5)). Then f_{hom}^∞ is a random volume integrand and for every $\omega \in \Omega'$, $x \in \mathbb{R}^n$, $\xi \in \mathbb{R}^{m \times n}$, $v \in \mathbb{S}^{n-1}$, and $k \in \mathbb{N}$ we have

$$f_{\text{hom}}^\infty(\omega, \xi) = \lim_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g_0}(\ell_\xi, Q_r^{v,k}(rx))}{k^{n-1}r^n} = \lim_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g_0}(\ell_\xi, Q_r)}{r^n}, \tag{9.12}$$

where $Q_r := Q_r(0)$. If, in addition, $(\tau_z)_{z \in \mathbb{Z}^n}$ is ergodic, then f_{hom}^∞ is independent of ω and

$$f_{\text{hom}}^\infty(\xi) = \lim_{r \rightarrow +\infty} \frac{1}{r^n} \int_\Omega m_\omega^{f^\infty, g_0}(\ell_\xi, Q_r) dP(\omega). \tag{9.13}$$

The following proposition establishes the existence of the random surface integrand g_{hom} .

Proposition 9.3 (Homogenised random surface integrand) *Let f be a stationary random volume integrand and let g be a stationary random surface integrand with respect to a group $(\tau_z)_{z \in \mathbb{Z}^n}$ of P -preserving transformations on (Ω, \mathcal{T}, P) . Then there exists $\Omega' \in \mathcal{T}$, with $P(\Omega') = 1$, such that for every $\omega \in \Omega'$, $x \in \mathbb{R}^n$, $\zeta \in \mathbb{R}^m$, $v \in \mathbb{S}^{n-1}$, the limit*

$$\lim_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{rx, \zeta, v}, Q_r^v(rx))}{r^{n-1}} \tag{9.14}$$

exists and is independent of x . More precisely, there exists a random volume integrand $g_{\text{hom}} : \Omega \times \mathbb{R}^m \times \mathbb{S}^{n-1} \rightarrow [0, +\infty)$ such that for every $\omega \in \Omega'$, $x \in \mathbb{R}^n$, $\zeta \in \mathbb{R}^m$, and $v \in \mathbb{S}^{n-1}$

$$g_{\text{hom}}(\omega, \zeta, v) = \lim_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{rx, \zeta, v}, Q_r^v(rx))}{r^{n-1}} = \lim_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{0, \zeta, v}, Q_r^v)}{r^{n-1}},$$

where $Q_r^v := Q_r^v(0)$. If, in addition, $(\tau_z)_{z \in \mathbb{Z}^n}$ is ergodic, then g_{hom} is independent of ω and

$$g_{\text{hom}}(\zeta, v) = \lim_{r \rightarrow +\infty} \frac{1}{r^{n-1}} \int_\Omega m_\omega^{f^\infty, g}(u_{0, \zeta, v}, Q_r^v) dP(\omega).$$

The proof of Proposition 9.3 follows immediately from Propositions 9.4 and 9.5 below. In the first one we prove the existence of the limit in (9.14) for $x = 0$, while in the second one we consider the general case $x \neq 0$ and prove that the limit is independent of x .

Proposition 9.4 *Let f be a stationary random volume integrand and let g be a stationary random surface integrand with respect to a group $(\tau_z)_{z \in \mathbb{Z}^n}$ of P -preserving*

transformations on (Ω, \mathcal{T}, P) . Then there exist $\tilde{\Omega} \in \mathcal{T}$, with $P(\tilde{\Omega}) = 1$, and a random surface integrand $g_{\text{hom}} : \Omega \times \mathbb{R}^m \times \mathbb{S}^{n-1} \rightarrow \mathbb{R}$ such that

$$g_{\text{hom}}(\omega, \zeta, \nu) = \lim_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{0, \zeta, \nu}, Q_r^\nu)}{r^{n-1}}, \tag{9.15}$$

for every $\omega \in \tilde{\Omega}$, $\zeta \in \mathbb{R}^m$, and $\nu \in \mathbb{S}^{n-1}$, where $Q_r^\nu := Q_r^\nu(0)$. If, in addition, $(\tau_z)_{z \in \mathbb{Z}^n}$ is ergodic, then g_{hom} is independent of ω and

$$g_{\text{hom}}(\zeta, \nu) = \lim_{r \rightarrow +\infty} \frac{1}{r^{n-1}} \int_\Omega m_\omega^{f^\infty, g}(u_{0, \zeta, \nu}, Q_r^\nu) dP(\omega). \tag{9.16}$$

Proof We adapt the proof of [17, Theorem 5.1]. The main difference is that now the functional to be minimised depends also on f^∞ , while in [17, Theorem 5.1] it depends only on g . Since this requires some changes, for completeness we prefer to give the whole proof in detail. We divide it into four steps.

Step 1: Existence of the limit in (9.15) for fixed $\zeta \in \mathbb{Q}^m$ and $\nu \in \mathbb{S}^{n-1} \cap \mathbb{Q}^n$.

Let $\nu \in \mathbb{S}^{n-1} \cap \mathbb{Q}^{n-1}$ and $\zeta \in \mathbb{Q}^m$ be fixed, let $R_\nu \in O(n) \cap \mathbb{Q}^{n \times n}$ be the orthogonal $n \times n$ matrix as in (h) in Section 2, and let M_ν be a positive integer such that $M_\nu R_\nu \in \mathbb{Z}^{n \times n}$. Note that, in particular, for every $z' \in \mathbb{Z}^{n-1}$ we have that $M_\nu R_\nu(z', 0) \in \Pi_0^\nu \cap \mathbb{Z}^n$, namely $M_\nu R_\nu$ maps integer vectors perpendicular to e_n into integer vectors perpendicular to ν .

Given $A' = [a_1, b_1) \times \dots \times [a_{n-1}, b_{n-1}) \in \mathcal{I}_{n-1}$ (see (3.15)), we define the (rotated) n -dimensional interval $T_\nu(A')$ as

$$T_\nu(A') := M_\nu R_\nu(A' \times [-c, c)), \quad \text{with } c := \frac{1}{2} \max_{1 \leq j \leq n-1} (b_j - a_j). \tag{9.17}$$

For every $\omega \in \Omega$ and $A' \in \mathcal{I}_{n-1}$ we set

$$\mu_{\zeta, \nu}(\omega, A') := \frac{1}{M_\nu^{n-1}} m_\omega^{f^\infty, g}(u_{0, \zeta, \nu}, T_\nu(A')). \tag{9.18}$$

Now let $(\Omega, \widehat{\mathcal{T}}, \widehat{P})$ denote the completion of the probability space (Ω, \mathcal{T}, P) . We claim that the function $\mu_{\zeta, \nu} : \Omega \times \mathcal{I}_{n-1} \rightarrow \mathbb{R}$ as in (9.18) defines an $(n-1)$ -dimensional subadditive process on $(\Omega, \widehat{\mathcal{T}}, \widehat{P})$. Indeed, thanks to Remark 3.9 and Proposition A.12, for every $A \in \mathcal{A}$ the function $\omega \mapsto m_\omega^{f^\infty, g}(u_{0, \zeta, \nu}, A)$ is $\widehat{\mathcal{T}}$ -measurable. From this, it follows that the function $\omega \mapsto \mu_{\zeta, \nu}(\omega, A')$ is $\widehat{\mathcal{T}}$ -measurable too.

We are now going to prove that $\mu_{\zeta, \nu}$ is covariant; that is, we show that there exists a group $(\tau_{z'}^\nu)_{z' \in \mathbb{Z}^{n-1}}$ of \widehat{P} -preserving transformations on $(\Omega, \widehat{\mathcal{T}}, \widehat{P})$ such that

$$\mu_{\zeta, \nu}(\omega, A' + z') = \mu_{\zeta, \nu}(\tau_{z'}^\nu(\omega), A'), \quad \text{for every } \omega \in \Omega, z' \in \mathbb{Z}^{n-1}, \text{ and } A' \in \mathcal{I}_{n-1}.$$

To this end fix $z' \in \mathbb{Z}^{n-1}$ and $A' \in \mathcal{I}_{n-1}$. Note that, by (9.17),

$$\begin{aligned} T_v(A' + z') &= M_v R_v((A' + z') \times [-c, c]) = M_v R_v(A' \times [-c, c]) + M_v R_v(z', 0) \\ &= T_v(A') + z'_v, \end{aligned}$$

where $z'_v := M_v R_v(z', 0) \in \Pi_0^v \cap \mathbb{Z}^n$. Then, by (9.18)

$$\begin{aligned} \mu_{\zeta,v}(\omega, A' + z') &= \frac{1}{M_v^{n-1}} m_\omega^{f^\infty, g}(u_{0,\zeta,v}, T_v(A' + z')) \\ &= \frac{1}{M_v^{n-1}} m_\omega^{f^\infty, g}(u_{0,\zeta,v}, T_v(A') + z'_v). \end{aligned} \tag{9.19}$$

Given $u \in SBV(\text{int}(T_v(A') + z'_v), \mathbb{R}^m)$ with $u = u_{0,\zeta,v}$ near $\partial(T_v(A') + z'_v)$, let $v \in SBV(\text{int}(T_v(A')), \mathbb{R}^m)$ be defined as $v(x) := u(x + z'_v)$ for every $x \in \mathbb{R}^n$. By a change of variables, using the stationarity of f^∞ and g we obtain

$$\begin{aligned} &\int_{T_v(A') + z'_v} f^\infty(\omega, x, \nabla u) \, dx + \int_{S_u \cap (T_v(A') + z'_v)} g(\omega, x, [u], \nu_u) \, d\mathcal{H}^{n-1} \\ &= \int_{T_v(A')} f^\infty(\omega, x + z'_v, \nabla v) \, dx + \int_{S_v \cap T_v(A')} g(\omega, x + z'_v, [v], \nu_v) \, d\mathcal{H}^{n-1} \\ &= \int_{T_v(A')} f^\infty(\tau_{z'_v}(\omega), x, \nabla v) \, dx + \int_{S_v \cap T_v(A')} g(\tau_{z'_v}(\omega), x, [v], \nu_v) \, d\mathcal{H}^{n-1}. \end{aligned}$$

Since z'_v is perpendicular to v , we have $u_{0,\zeta,v}(x) = u_{0,\zeta,v}(x + z'_v)$ for every $x \in \mathbb{R}^n$. Therefore, from (9.18) and (9.19) we obtain that

$$\mu_{\zeta,v}(\omega, A' + z') = \mu_{\zeta,v}(\tau_{z'_v}^v(\omega), A'),$$

where we set

$$(\tau_{z'}^v)_{z' \in \mathbb{Z}^{n-1}} := (\tau_{z'_v}^v)_{z'_v \in \mathbb{Z}^{n-1}}.$$

We now show that $\mu_{\zeta,v}$ is subadditive in \mathcal{I}_{n-1} . To this end let $A' \in \mathcal{I}_{n-1}$ and let $(A'_i)_{1 \leq i \leq N} \subset \mathcal{I}_{n-1}$ be a finite family of pairwise disjoint sets such that $A' = \bigcup_i A'_i$. For fixed $\eta > 0$ and $i = 1, \dots, N$, let $u_i \in SBV(\text{int}(T_v(A'_i)), \mathbb{R}^m)$ be such that $u_i = u_{0,\zeta,v}$ in a neighbourhood of $\partial T_v(A'_i)$ and

$$\begin{aligned} &\int_{T_v(A'_i)} f^\infty(\omega, x, \nabla u_i) \, dx + \int_{S_{u_i} \cap T_v(A'_i)} g(\omega, x, [u_i], \nu_{u_i}) \, d\mathcal{H}^{n-1} \\ &\leq m_\omega^{f^\infty, g}(u_{0,\zeta,v}, T_v(A'_i)) + \eta. \end{aligned} \tag{9.20}$$

Note that $T_v(A')$ can differ from $\bigcup_i T_v(A'_i)$ (see for instance [17, Figure 2]), but, by construction, we always have $\bigcup_i T_v(A'_i) \subset T_v(A')$.

Now we define

$$u(x) := \begin{cases} u_i(x) & \text{if } x \in T_v(A'_i), i = 1, \dots, N, \\ u_{0,\zeta,v}(x) & \text{if } x \in T_v(A') \setminus \bigcup_i T_v(A'_i); \end{cases}$$

then $u \in SBV(T_v(A'), \mathbb{R}^m)$ and $u = u_{0,\zeta,v}$ near $\partial T_v(A')$. Note that we also have x

$$S_u \cap T_v(A') = \bigcup_{i=1}^N (S_{u_i} \cap T_v(A'_i)).$$

Indeed, $S_u \cap T_v(A') \cap \partial T_v(A'_i) = \emptyset$ for every $i = 1, \dots, N$. Moreover, $u = u_{0,\zeta,v}$ in $T_v(A') \setminus \bigcup_i T_v(A'_i)$, hence $\nabla u = 0$ a.e. in this set. Therefore, recalling that $f^\infty(\omega, \cdot, 0) \equiv 0$, we obtain

$$\begin{aligned} & \int_{T_v(A')} f^\infty(\omega, x, \nabla u) dx + \int_{S_u \cap T_v(A')} g(\omega, x, [u], \nu_u) d\mathcal{H}^{n-1} \\ &= \sum_{i=1}^N \left(\int_{T_v(A'_i)} f^\infty(\omega, x, \nabla u_i) dx + \int_{S_{u_i} \cap T_v(A'_i)} g(\omega, x, [u_i], \nu_{u_i}) d\mathcal{H}^{n-1} \right). \end{aligned}$$

As a consequence, by (9.20),

$$m_\omega^{f^\infty, g}(u_{0,\zeta,v}, T_v(A')) \leq \sum_{i=1}^N m_\omega^{f^\infty, g}(u_{0,\zeta,v}, T_v(A'_i)) + N\eta,$$

thus the subadditivity of $\mu_{\zeta,v}$ follows from (9.18), by the arbitrariness of $\eta > 0$.

Finally, in view of (g4) for every $A' \in \mathcal{I}_{n-1}$ and for \widehat{P} -a.e. $\omega \in \Omega$ we have

$$\begin{aligned} \mu_{\zeta,v}(\omega, A') &= \frac{1}{M_v^{n-1}} m_\omega^{f^\infty, g}(u_{0,\zeta,v}, T_v(A')) \\ &\leq \frac{1}{M_v^{n-1}} \int_{S_{u_{0,\zeta,v}} \cap T_v(A')} g(\omega, x, \zeta, \nu) d\mathcal{H}^{n-1} \\ &\leq \frac{c_3 |\zeta|}{M_v^{n-1}} \mathcal{H}^{n-1}(\Pi_0^v \cap T_v(A')) = c_3 |\zeta| \mathcal{L}^{n-1}(A'), \end{aligned} \tag{9.21}$$

where we used again the fact that $f^\infty(\omega, \cdot, 0) \equiv 0$. This concludes the proof of the fact that $\mu_{\zeta,v}$ is an $(n - 1)$ -dimensional subadditive process.

We can now apply Theorem 3.15 to the subadditive process $\mu_{\zeta,v}$, defined on $(\Omega, \widehat{T}, \widehat{P})$ by (9.18), to deduce the existence of a \widehat{T} -measurable function $\psi_{\zeta,v} : \Omega \rightarrow [0, +\infty)$ and a set $\widehat{\Omega}_{\zeta,v} \subset \Omega$, with $\widehat{\Omega}_{\zeta,v} \in \widehat{T}$ and $P(\widehat{\Omega}_{\zeta,v}) = 1$ such that

$$\lim_{r \rightarrow +\infty} \frac{\mu_{\zeta,v}(\omega, rQ')}{r^{n-1}} = \psi_{\zeta,v}(\omega) \tag{9.22}$$

for every $\omega \in \widehat{\Omega}_{\zeta, \nu}$, where $Q' := [-\frac{1}{2}, \frac{1}{2}]^{n-1}$. Then, by the properties of the completion, there exist a set $\Omega_{\zeta, \nu} \in \mathcal{T}$, with $P(\Omega_{\zeta, \nu}) = 1$, and a \mathcal{T} -measurable function, which we still denote by $\psi_{\zeta, \nu}$, such that (9.22) holds for every $\omega \in \Omega_{\zeta, \nu}$. Using the definition of $\mu_{\zeta, \nu}$ we then have

$$\psi_{\zeta, \nu}(\omega) = \lim_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{0, \zeta, \nu}, Q_r^\nu)}{r^{n-1}} \tag{9.23}$$

for every $\omega \in \Omega_{\zeta, \nu}$.

Step 2: Existence of the limit in (9.15) for every $\zeta \in \mathbb{R}^m$ and $\nu \in \mathbb{S}^{n-1}$.

Let $\widetilde{\Omega}$ denote the intersection of the sets $\Omega_{\zeta, \nu}$ for $\zeta \in \mathbb{Q}^m$ and $\nu \in \mathbb{S}^{n-1} \cap \mathbb{Q}^n$; clearly $\widetilde{\Omega} \in \mathcal{T}$ and $P(\widetilde{\Omega}) = 1$. Let $\underline{g}, \overline{g}: \widetilde{\Omega} \times \mathbb{R}^m \times \mathbb{S}^{n-1} \rightarrow [0, +\infty]$ be the functions defined as

$$\underline{g}(\omega, \zeta, \nu) := \liminf_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{0, \zeta, \nu}, Q_r^\nu)}{r^{n-1}}, \tag{9.24}$$

$$\overline{g}(\omega, \zeta, \nu) := \limsup_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{0, \zeta, \nu}, Q_r^\nu)}{r^{n-1}}. \tag{9.25}$$

By (9.23) we have

$$\underline{g}(\omega, \zeta, \nu) = \overline{g}(\omega, \zeta, \nu) = \psi_{\zeta, \nu}(\omega) \quad \text{for every } \omega \in \widetilde{\Omega}, \zeta \in \mathbb{Q}^m, \text{ and } \nu \in \mathbb{S}^{n-1} \cap \mathbb{Q}^n. \tag{9.26}$$

By Lemma 4.5 (property (4.13)), for every $\omega \in \widetilde{\Omega}$ and $\nu \in \mathbb{S}^{n-1}$ the functions $\zeta \mapsto \underline{g}(\omega, \zeta, \nu)$ and $\zeta \mapsto \overline{g}(\omega, \zeta, \nu)$ are continuous on \mathbb{R}^m and their modulus of continuity does not depend on ω and ν . More precisely, recalling (g4), for every $\omega \in \widetilde{\Omega}$ and $\nu \in \mathbb{S}^{n-1}$ we have

$$\begin{aligned} |\underline{g}(\omega, \zeta_1, \nu) - \underline{g}(\omega, \zeta_2, \nu)| &\leq c_3 \sigma_2(|\zeta_1 - \zeta_2|)(|\zeta_1| + |\zeta_2|), \\ |\overline{g}(\omega, \zeta_1, \nu) - \overline{g}(\omega, \zeta_2, \nu)| &\leq c_3 \sigma_2(|\zeta_1 - \zeta_2|)(|\zeta_1| + |\zeta_2|), \end{aligned} \quad \text{for every } \zeta_1, \zeta_2 \in \mathbb{R}^m. \tag{9.27}$$

From these inequalities and from (9.26) we deduce that for every $\zeta \in \mathbb{R}^m$ and $\nu \in \mathbb{S}^{n-1} \cap \mathbb{Q}^n$ there exists a \mathcal{T} -measurable function, which we still denote by $\psi_{\zeta, \nu}$, such that

$$\underline{g}(\omega, \zeta, \nu) = \overline{g}(\omega, \zeta, \nu) = \psi_{\zeta, \nu}(\omega) \quad \text{for every } \omega \in \widetilde{\Omega}. \tag{9.28}$$

We now claim that for every $\omega \in \widetilde{\Omega}$ and every $\zeta \in \mathbb{R}^m$ the restrictions of the functions $\nu \mapsto \underline{g}(\omega, \zeta, \nu)$ and $\nu \mapsto \overline{g}(\omega, \zeta, \nu)$ to the sets $\widehat{\mathbb{S}}_+^{n-1}$ and $\widehat{\mathbb{S}}_-^{n-1}$ are continuous. We only prove this property for \underline{g} and $\widehat{\mathbb{S}}_+^{n-1}$, the other proofs being analogous. To this end, let us fix $\zeta \in \mathbb{R}^m$, $\nu \in \widehat{\mathbb{S}}_+^{n-1}$, and a sequence $(\nu_j) \subset \widehat{\mathbb{S}}_+^{n-1}$ such that $\nu_j \rightarrow \nu$ as

$j \rightarrow +\infty$. Since the restriction of the function $v \mapsto R_v$ to $\widehat{\mathbb{S}}_+^{n-1}$ is continuous, for every $\delta \in (0, \frac{1}{2})$ there exists an integer j_δ such that

$$Q_{(1-\delta)r}^{v_j} \subset\subset Q_r^v \subset\subset Q_{(1+\delta)r}^{v_j}, \tag{9.29}$$

for every $j \geq j_\delta$ and every $r > 0$. Fix $j \geq j_\delta$, $r > 0$, and $\eta > 0$. Let $u \in SBV(Q_r^v, \mathbb{R}^m)$ be such that $u = u_{0,\zeta,v}$ near ∂Q_r^v and

$$\int_{Q_r^v} f^\infty(\omega, x, \nabla u) dx + \int_{S_u \cap Q_r^v} g(\omega, x, [u], v_u) d\mathcal{H}^{n-1} \leq m_\omega^{f^\infty, g}(u_{0,\zeta,v}, Q_r^v) + \eta r^{n-1}.$$

We define $v \in SBV(Q_{(1+\delta)r}^{v_j}, \mathbb{R}^m)$ as

$$v(x) := \begin{cases} u(x) & \text{if } x \in Q_r^v, \\ u_{0,\zeta,v_j}(x) & \text{if } x \in Q_{(1+\delta)r}^{v_j} \setminus Q_r^v. \end{cases}$$

Then, $v = u_{0,\zeta,v}$ near $\partial Q_{(1+\delta)r}^{v_j}$ and $S_v \subset S_u \cup \Sigma$, where

$$\Sigma := \{x \in \partial Q_r^v : (x \cdot v)(x \cdot v_j) < 0\} \cup (\Pi_0^{v_j} \cap (Q_{(1+\delta)r}^{v_j} \setminus Q_r^v)).$$

Moreover $||[v]|| \leq |\zeta| \mathcal{H}^{n-1}$ -a.e. on Σ . By (9.29) there exists $\zeta_j(\delta) > 0$, independent of r , with $\zeta_j(\delta) \rightarrow (1 + \delta)^{n-1} - 1$ as $j \rightarrow +\infty$, such that $\mathcal{H}^{n-1}(\Sigma) \leq \zeta_j(\delta)r^{n-1}$. Thanks to (g4) we then have

$$\begin{aligned} m_\omega^{f^\infty, g}(u_{0,\zeta,v}, Q_{(1+\delta)r}^{v_j}) &\leq \int_{Q_{(1+\delta)r}^{v_j}} f^\infty(\omega, x, \nabla v) dx \\ &\quad + \int_{S_v \cap Q_{(1+\delta)r}^{v_j}} g(\omega, x, [v], v_v) d\mathcal{H}^{n-1} \\ &\leq \int_{Q_r^v} f^\infty(\omega, x, \nabla u) dx + \int_{S_u \cap Q_r^v} g(\omega, x, [u], v_u) d\mathcal{H}^{n-1} \\ &\quad + c_3 |\zeta| \zeta_j(\delta) r^{n-1} \\ &\leq m_\omega^{f^\infty, g}(u_{0,\zeta,v}, Q_r^v) + \eta r^{n-1} + c_3 |\zeta| \zeta_j(\delta) r^{n-1}, \end{aligned}$$

where we used the fact that $f^\infty(\omega, \cdot, 0) \equiv 0$. Recalling definition (9.24), dividing by r^{n-1} , and passing to the liminf as $r \rightarrow +\infty$, we obtain

$$\underline{g}(\omega, \zeta, v_j)(1 + \delta)^{n-1} \leq \underline{g}(\omega, \zeta, v) + \eta + c_3 |\zeta| \zeta_j(\delta). \tag{9.30}$$

Letting $j \rightarrow +\infty$, then $\delta \rightarrow 0+$, and then $\eta \rightarrow 0+$, we deduce that

$$\limsup_{j \rightarrow +\infty} \underline{g}(\omega, \zeta, v_j) \leq \underline{g}(\omega, \zeta, v).$$

An analogous argument, now using the cubes $Q_{(1-\delta)r}^{\nu_j}$, shows that

$$\underline{g}(\omega, \zeta, \nu) \leq \liminf_{j \rightarrow +\infty} \underline{g}(\omega, \zeta, \nu_j),$$

hence the claim follows. Note that, together with (9.27), this implies that for every $\omega \in \tilde{\Omega}$ the restriction of the function $(\zeta, \nu) \mapsto \underline{g}(\omega, \zeta, \nu)$ to $\mathbb{R}^m \times \widehat{\mathbb{S}}_{\pm}^{n-1}$ is continuous, and the same holds true for $(\zeta, \nu) \mapsto \overline{g}(\omega, \zeta, \nu)$.

As we already observed in the proof of Step 2 of Proposition 9.1, the set $\widehat{\mathbb{S}}_{\pm}^{n-1} \cap \mathbb{Q}^n$ is dense in $\widehat{\mathbb{S}}_{\pm}^{n-1}$. Therefore, from the continuity property proved above and from (9.28) we deduce that for every $\zeta \in \mathbb{R}^m$ and $\nu \in \mathbb{S}^{n-1}$ there exists a \mathcal{T} -measurable function, which we still denote by $\psi_{\zeta, \nu}$, such that

$$\underline{g}(\omega, \zeta, \nu) = \overline{g}(\omega, \zeta, \nu) = \psi_{\zeta, \nu}(\omega) \quad \text{for every } \omega \in \tilde{\Omega}. \tag{9.31}$$

By (9.24) and (9.25) this implies that

$$\psi_{\zeta, \nu}(\omega) = \lim_{r \rightarrow +\infty} \frac{m_{\omega}^{f, \infty, g}(u_{0, \zeta, \nu}, Q_r^{\nu})}{r^{n-1}} \tag{9.32}$$

for every $\omega \in \tilde{\Omega}$, $\zeta \in \mathbb{R}^m$, and $\nu \in \mathbb{S}^{n-1}$, concluding the proof of Step 2.

Step 3: Definition and properties of g_{hom} .

For every $\omega \in \Omega$ and $\zeta \in \mathbb{R}^m$, and $\nu \in \mathbb{S}^{n-1}$ we define

$$g_{\text{hom}}(\omega, \zeta, \nu) := \begin{cases} \psi_{\zeta, \nu}(\omega) & \text{if } \omega \in \tilde{\Omega}, \\ c_2 |\zeta| & \text{if } \omega \in \Omega \setminus \tilde{\Omega}. \end{cases}$$

Then (9.15) follows from (9.32). From the measurability of $\psi_{\zeta, \nu}$, proved in Step 2, we obtain that $g_{\text{hom}}(\cdot, \zeta, \nu)$ is \mathcal{T} -measurable in Ω for every $\zeta \in \mathbb{R}^m$ and $\nu \in \mathbb{S}^{n-1}$. Moreover, since for every $\omega \in \tilde{\Omega}$ the restriction of the function $(\zeta, \nu) \mapsto \underline{g}(\omega, \zeta, \nu)$ to $\mathbb{R}^m \times \widehat{\mathbb{S}}_{\pm}^{n-1}$ is continuous, from (9.31) we deduce that for every $\omega \in \Omega$ the restriction of $g_{\text{hom}}(\omega, \cdot, \cdot)$ to $\mathbb{R}^m \times \widehat{\mathbb{S}}_{\pm}^{n-1}$ is continuous and this implies the $\mathcal{T} \otimes \mathcal{B}^{m \times n} \otimes \mathcal{B}_{\mathbb{S}^n}^n$ -measurability of g_{hom} on $\Omega \times \mathbb{R}^m \times \mathbb{S}^{n-1}$. Finally, Lemma 4.5 allows us to conclude that $g_{\text{hom}}(\omega, \cdot, \cdot) \in \mathcal{G}$ for every $\omega \in \Omega$.

Step 4: In the ergodic case g_{hom} is deterministic.

Set $\widehat{\Omega} := \bigcap_{z \in \mathbb{Z}^n} \tau_z(\tilde{\Omega})$; we clearly have that $\widehat{\Omega} \in \mathcal{T}$, $\widehat{\Omega} \subset \tilde{\Omega}$, and $\tau_z(\widehat{\Omega}) = \widehat{\Omega}$ for every $z \in \mathbb{Z}^n$; moreover, since τ_z is a P -preserving transformation and $P(\tilde{\Omega}) = 1$, we have $P(\widehat{\Omega}) = 1$. We claim that

$$g_{\text{hom}}(\tau_z(\omega), \zeta, \nu) = g_{\text{hom}}(\omega, \zeta, \nu), \tag{9.33}$$

for every $z \in \mathbb{Z}^n$, $\omega \in \widehat{\Omega}$, $\zeta \in \mathbb{R}^m$, and $\nu \in \mathbb{S}^{n-1}$.

We start noting that to prove (9.33) it is enough to show that

$$g_{\text{hom}}(\tau_z(\omega), \zeta, \nu) \leq g_{\text{hom}}(\omega, \zeta, \nu) \tag{9.34}$$

for every $z \in \mathbb{Z}^n$, $\omega \in \widehat{\Omega}$, $\zeta \in \mathbb{R}^m$, and $v \in \mathbb{S}^{n-1}$. Indeed, the opposite inequality is obtained by applying (9.34) with ω replaced by $\tau_z(\omega)$ and z replaced by $-z$.

Let $z \in \mathbb{Z}^n$, $\omega \in \widehat{\Omega}$, $\zeta \in \mathbb{R}^m$, and $v \in \mathbb{S}^{n-1}$ be fixed. For every $r > 3|z|$, let $u_r \in SBV(Q_r^v, \mathbb{R}^m)$ be such that $u_r = u_{0,\zeta,v}$ near ∂Q_r^v , and

$$\begin{aligned} & \int_{Q_r^v} f^\infty(\omega, x, \nabla u_r) dx + \int_{S_{u_r} \cap Q_r^v} g(\omega, x, [u_r], v_{u_r}) d\mathcal{H}^{n-1} \\ & \leq m_\omega^{f^\infty, g}(u_{0,\zeta,v}, Q_r^v) + 1. \end{aligned} \tag{9.35}$$

By the stationarity of f^∞ and g , a change of variables gives

$$m_{\tau_z(\omega)}^{f^\infty, g}(u_{0,\zeta,v}, Q_r^v) = m_\omega^{f^\infty, g}(u_{z,\zeta,v}, Q_r^v(z)). \tag{9.36}$$

We now modify u_r to obtain a competitor for a minimisation problem related to the right-hand side of (9.36). Noting that $Q_r^v \subset\subset Q_{r+3|z|}^v(z)$ we define

$$v_r(x) := \begin{cases} u_r(x) & \text{if } x \in Q_r^v, \\ u_{z,\zeta,v}(x) & \text{if } x \in Q_{r+3|z|}^v(z) \setminus Q_r^v. \end{cases}$$

Clearly $v_r \in SBV(Q_{r+3|z|}^v(z), \mathbb{R}^m)$ and $v_r = u_{z,\zeta,v}$ near $\partial Q_{r+3|z|}^v(z)$. Moreover we notice that $S_{v_r} = S_{u_r} \cup \Sigma_1 \cup \Sigma_2$, where

$$\Sigma_1 := \{x \in \partial Q_r^v : (x \cdot v)((x - z) \cdot v) < 0\} \quad \text{and} \quad \Sigma_2 := \Pi_z^v \cap (Q_{r+3|z|}^v(z) \setminus Q_r^v).$$

Moreover $||[v_r]|| = |\zeta| \mathcal{H}^{n-1}$ -a.e. on $\Sigma_1 \cup \Sigma_2$. Since $3|z| < r$, we have $\mathcal{H}^{n-1}(\Sigma_1) = 2(n-1)|z \cdot v| r^{n-2}$ and $\mathcal{H}^{n-1}(\Sigma_2) = (r+3|z|)^{n-1} - r^{n-1} \leq 3(n-1)|z|(r+3|z|)^{n-2} < 2^n(n-1)|z| r^{n-2}$. Therefore, using the fact that $f^\infty(\omega, \cdot, 0) \equiv 0$, thanks to (g4) we have

$$\begin{aligned} & \int_{Q_{r+3|z|}^v(z)} f^\infty(\omega, x, \nabla v_r) dx + \int_{S_{v_r} \cap Q_{r+3|z|}^v(z)} g(\omega, x, [v_r], v_{v_r}) d\mathcal{H}^{n-1} \\ & \leq \int_{Q_r^v} f^\infty(\omega, x, \nabla u_r) dx + \int_{S_{u_r} \cap Q_r^v} g(\omega, x, [u_r], v_{u_r}) d\mathcal{H}^{n-1} + M_{\zeta,z} r^{n-2}, \end{aligned}$$

where $M_{\zeta,z} := c_3(n-1)(2+2^n)|z||\zeta|$. This inequality, combined with (9.35) yields

$$m_\omega^{f^\infty, g}(u_{z,\zeta,v}, Q_{r+3|z|}^v(z)) \leq m_\omega^{f^\infty, g}(u_{0,\zeta,v}, Q_r^v) + 1 + M_{\zeta,z} r^{n-2}. \tag{9.37}$$

Recalling that $\tau_z(\omega) \in \widehat{\Omega} \subset \widetilde{\Omega}$, by (9.15) and (9.36) we get

$$g_{\text{hom}}(\tau_z(\omega), \zeta, v) = \lim_{r \rightarrow +\infty} \frac{m_{\tau_z(\omega)}^{f^\infty, g}(u_{0,\zeta,v}, Q_r^v)}{r^{n-1}} = \lim_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{z,\zeta,v}, Q_r^v(z))}{r^{n-1}}$$

$$= \lim_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{z, \zeta, \nu}, Q_{r+3|z|}^\nu(z))}{r^{n-1}},$$

where in the last equality we have used the fact that $r^{n-1}/(r + 3|z|)^{n-1} \rightarrow 1$ as $r \rightarrow +\infty$. Therefore, dividing all terms of (9.37) by r^{n-1} and passing to the limit as $r \rightarrow +\infty$, from (9.15) we obtain the inequality

$$g_{\text{hom}}(\tau_z(\omega), \zeta, \nu) \leq g_{\text{hom}}(\omega, \zeta, \nu),$$

which proves (9.34) and hence the claim.

If $(\tau_z)_{z \in \mathbb{Z}^n}$ is ergodic we can invoke [17, Corollary 6.3] to deduce that g_{hom} does not depend on ω and hence is deterministic. In this case, (9.16) can be obtained by integrating (9.15) over Ω and observing that, thanks to (9.21), we can apply the Dominated Convergence Theorem. \square

We now prove that the limit (9.14) that defines g_{hom} is independent of x . More precisely we prove the following result.

Proposition 9.5 *Let f be a stationary random volume integrand and let g be a stationary random surface integrand with respect to a group $(\tau_z)_{z \in \mathbb{Z}^n}$ of P -preserving transformations on (Ω, \mathcal{T}, P) . Then there exist $\Omega' \in \mathcal{T}$, with $P(\Omega') = 1$, and a random surface integrand $g_{\text{hom}} : \Omega \times \mathbb{R}^m \times \mathbb{S}^{n-1} \rightarrow \mathbb{R}$, independent of x , such that*

$$g_{\text{hom}}(\omega, \zeta, \nu) = \lim_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{rx, \zeta, \nu}, Q_r^\nu(rx))}{r^{n-1}}, \tag{9.38}$$

for every $\omega \in \Omega'$, $x \in \mathbb{R}^n$, $\zeta \in \mathbb{R}^m$, $\nu \in \mathbb{S}^{n-1}$.

Proof The proof closely follows that of [17, Theorem 6.1], therefore here we only discuss the main differences with respect to [17].

Let g_{hom} be the random surface integrand introduced in Proposition 9.4. Arguing as in the proof of [17, Theorem 6.1], we can prove the existence of $\Omega' \in \mathcal{T}$, with $P(\Omega') = 1$, such that (9.38) holds for every $\omega \in \Omega'$, $x \in \mathbb{R}^n$, $\zeta \in \mathbb{R}^m$, and $\nu \in \mathbb{S}^{n-1} \cap \mathbb{Q}^{n-1}$. Hence, to conclude it remains to show that (9.38) holds true for every $\nu \in \mathbb{S}^{n-1}$.

To this end, for fixed $\omega \in \Omega'$, $x \in \mathbb{R}^n$, $\zeta \in \mathbb{R}^m$, and $\nu \in \mathbb{S}^{n-1}$, we introduce the auxiliary functions

$$\underline{g}(\omega, x, \zeta, \nu) := \liminf_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{rx, \zeta, \nu}, Q_r^\nu(rx))}{r^{n-1}}, \tag{9.39}$$

$$\overline{g}(\omega, x, \zeta, \nu) := \limsup_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{rx, \zeta, \nu}, Q_r^\nu(rx))}{r^{n-1}}. \tag{9.40}$$

Let $\nu \in \widehat{\mathbb{S}}_+^{n-1}$ be fixed. As we already observed in the proof of Step 2 of Proposition 9.1, the set $\widehat{\mathbb{S}}_+^{n-1} \cap \mathbb{Q}^n$ is dense in $\widehat{\mathbb{S}}_+^{n-1}$, hence there exists a sequence

$(v_j) \subset \widehat{\mathbb{S}}_+^{n-1} \cap \mathbb{Q}^{n-1}$ such that $v_j \rightarrow v$ as $j \rightarrow \infty$. We claim that for every $\delta \in (0, 1/2)$ there exists $j_\delta \in \mathbb{N}$ such that

$$(1 + \delta)^{n-1} \underline{g}(\omega, \frac{x}{1+\delta}, \zeta, v_j) \leq \underline{g}(\omega, x, \zeta, v) + c_3 |\zeta| \varsigma_j(\delta), \tag{9.41}$$

$$\overline{g}(\omega, x, \zeta, v) \leq (1 - \delta)^{n-1} \overline{g}(\omega, \frac{x}{1-\delta}, \zeta, v_j) + c_3 |\zeta| \varsigma_j(\delta), \tag{9.42}$$

for every $j \geq j_\delta$, where $\varsigma_j(\delta)$ is such that $\varsigma_j(\delta) \rightarrow (1 + \delta)^{n-1} - 1$ as $j \rightarrow +\infty$.

The proof of (9.41) and (9.42) is similar to that of (9.30) in Proposition 9.4. Thanks to the continuity of the restriction of $v \mapsto R_v$ to $\widehat{\mathbb{S}}_+^{n-1}$, for every $\delta \in (0, \frac{1}{2})$ there exists an integer j_δ such that

$$Q_{(1-\delta)r}^{v_j}(rx) \subset\subset Q_r^v(rx) \subset\subset Q_{(1+\delta)r}^{v_j}(rx), \tag{9.43}$$

for every $j \geq j_\delta$ and every $r > 0$. Fix $j \geq j_\delta$, $r > 0$, and $\eta > 0$. Let $u \in SBV(Q_r^v(rx), \mathbb{R}^m)$ be such that $u = u_{rx, \zeta, v}$ near $\partial Q_r^v(rx)$ and

$$\begin{aligned} & \int_{Q_r^v(rx)} f^\infty(\omega, y, \nabla u) dy + \int_{S_u \cap Q_r^v(rx)} g(\omega, y, [u], v_u) d\mathcal{H}^{n-1} \\ & \leq m_\omega^{f^\infty, g}(u_{rx, \zeta, v}, Q_r^v(rx)) + \eta r^{n-1}. \end{aligned}$$

We define $v \in SBV(Q_{(1+\delta)r}^{v_j}(rx), \mathbb{R}^m)$ as

$$v(y) := \begin{cases} u(y) & \text{if } y \in Q_r^v(rx), \\ u_{rx, \zeta, v_j}(y) & \text{if } y \in Q_{(1+\delta)r}^{v_j}(rx) \setminus Q_r^v(rx). \end{cases}$$

Then, $v = u_{rx, \zeta, v_j}$ near $\partial Q_{(1+\delta)r}^{v_j}(rx)$, and $S_v \subset S_u \cup \Sigma$, where

$$\begin{aligned} \Sigma := & \{y \in \partial Q_r^v(rx) : ((y - rx) \cdot v)((y - rx) \cdot v_j) \\ & < 0\} \cup (\Pi_{rx}^{v_j} \cap (Q_{(1+\delta)r}^{v_j}(rx) \setminus Q_r^v(rx))). \end{aligned}$$

Moreover $||[v]|| \leq |\zeta| \mathcal{H}^{n-1}$ -a.e. on Σ . By (9.43) there exists $\varsigma_j(\delta) > 0$, independent of r , with $\varsigma_j(\delta) \rightarrow (1 + \delta)^{n-1} - 1$ as $j \rightarrow +\infty$, such that $\mathcal{H}^{n-1}(\Sigma) \leq \varsigma_j(\delta)r^{n-1}$. Thanks to (g4) we then have

$$\begin{aligned} & m_\omega^{f^\infty, g}(u_{rx, \zeta, v_j}, Q_{(1+\delta)r}^{v_j}(rx)) \\ & \leq \int_{Q_{(1+\delta)r}^{v_j}(rx)} f^\infty(\omega, x, \nabla v) dx + \int_{S_v \cap Q_{(1+\delta)r}^{v_j}(rx)} g(\omega, x, [v], v_v) d\mathcal{H}^{n-1} \\ & \leq \int_{Q_r^v(rx)} f^\infty(\omega, x, \nabla u) dx + \int_{S_u \cap Q_r^v(rx)} g(\omega, x, [u], v_u) d\mathcal{H}^{n-1} + c_3 |\zeta| \varsigma_j(\delta)r^{n-1} \\ & \leq m_\omega^{f^\infty, g}(u_{rx, \zeta, v}, Q_r^v(rx)) + \eta r^{n-1} + c_3 |\zeta| \varsigma_j(\delta)r^{n-1}, \end{aligned}$$

where we used the fact that $f^\infty(\omega, \cdot, 0) \equiv 0$. Recalling definition (9.39), dividing by r^{n-1} , and passing to the liminf as $r \rightarrow +\infty$, we obtain

$$(1 + \delta)^{n-1} \underline{g}(\omega, \frac{x}{1+\delta}, \zeta, \nu_j) \leq \underline{g}(\omega, x, \zeta, \nu) + \eta + c_3 |\zeta| \zeta_j(\delta),$$

which gives (9.41) by the arbitrariness of η . The proof of (9.42) is analogous.

From (9.41) and (9.42) we get

$$\begin{aligned} (1 + \delta)^{n-1} \underline{g}(\omega, \frac{x}{1+\delta}, \zeta, \nu_j) - c_3 |\zeta| \zeta_j(\delta) &\leq \underline{g}(\omega, x, \zeta, \nu) \leq \bar{g}(\omega, x, \zeta, \nu) \\ &\leq (1 - \delta)^{n-1} \bar{g}(\omega, \frac{x}{1-\delta}, \zeta, \nu_j) + c_3 |\zeta| \zeta_j(\delta) \end{aligned}$$

for every $j \geq j_\delta$. Since $\nu_j \in \mathbb{S}^{n-1} \cap \mathbb{Q}^n$, and (9.38) holds true for rational directions, we have

$$\underline{g}(\omega, \frac{x}{1+\delta}, \zeta, \nu_j) = \bar{g}(\omega, \frac{x}{1-\delta}, \zeta, \nu_j) = g_{\text{hom}}(\omega, \zeta, \nu_j).$$

This, together with the previous inequality, yields

$$\begin{aligned} (1 + \delta)^{n-1} g_{\text{hom}}(\omega, \zeta, \nu_j) - c_3 |\zeta| \zeta_j(\delta) &\leq \underline{g}(\omega, x, \zeta, \nu) \leq \bar{g}(\omega, x, \zeta, \nu) \\ &\leq (1 - \delta)^{n-1} g_{\text{hom}}(\omega, \zeta, \nu_j) + c_3 |\zeta| \zeta_j(\delta) \end{aligned}$$

for every $j \geq j_\delta$. Hence, taking the liminf as $j \rightarrow +\infty$ and then the limit as $\delta \rightarrow 0+$, we obtain

$$\liminf_{j \rightarrow +\infty} g_{\text{hom}}(\omega, \zeta, \nu_j) \leq \underline{g}(\omega, x, \zeta, \nu) \leq \bar{g}(\omega, x, \zeta, \nu) \leq \liminf_{j \rightarrow +\infty} g_{\text{hom}}(\omega, \zeta, \nu_j)$$

and hence

$$\underline{g}(\omega, x, \zeta, \nu) = \bar{g}(\omega, x, \zeta, \nu) = \liminf_{j \rightarrow +\infty} g_{\text{hom}}(\omega, \zeta, \nu_j).$$

Note that, in particular, all the terms in the above chain of equalities do not depend on x . Then, in view of the definition of \underline{g} and \bar{g} (see (9.39) and (9.40)) we get that the limit

$$\lim_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{rx, \zeta, \nu}, Q_r^\nu(rx))}{r^{n-1}}$$

exists and is independent of x . Therefore we obtain

$$\lim_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{x, \zeta, \nu}, Q_r^\nu(rx))}{r^{n-1}} = \lim_{r \rightarrow +\infty} \frac{m_\omega^{f^\infty, g}(u_{0, \zeta, \nu}, Q_r^\nu(0))}{r^{n-1}} = g_{\text{hom}}(\omega, \zeta, \nu),$$

for every $\omega \in \Omega'$, $x \in \mathbb{R}^n$, $\zeta \in \mathbb{R}^m$, and $\nu \in \widehat{\mathbb{S}}_+^{n-1}$. Since the same property holds for $\nu \in \widehat{\mathbb{S}}_-^{n-1}$, this concludes the proof. □

We are now in a position to prove the theorem concerning the existence, for P -almost every $\omega \in \Omega$, of the limits which define the homogenised integrands.

Proof of Theorem 3.17 Property (a), (3.17), and (3.20) are proved in Proposition 9.1, while property (b), (3.18), and (3.21) are proved in Proposition 9.3. Equalities (3.19) and (3.22) coincide with (9.12) and (9.13), which are proved in Proposition 9.2. \square

We now prove the main result of the paper.

Proof of Theorem 3.18 It is enough to apply Theorem 3.17 together with the deterministic homogenisation result in Theorem 4.1, applied for fixed $\omega \in \Omega'$. \square

Acknowledgements F. Cagnetti was supported by the EPSRC under the Grant EP/P007287/1 “Symmetry of Minimisers in Calculus of Variations”. The research of G. Dal Maso was partially funded by the European Research Council under Grant No. 290888 “Quasistatic and Dynamic Evolution Problems in Plasticity and Fracture”. G. Dal Maso is a member of the Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). The work of C. I. Zeppieri was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) project 3160408400 and under the Germany Excellence Strategy EXC 2044-390685587, Mathematics Münster: Dynamics–Geometry–Structure.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Appendix. Measurability issues

The purpose of this section is to prove the measurability of the functions defined in (3.16). This will be done in Proposition A.12, which requires some preliminary results.

We start by introducing some notation that will be used throughout the proofs. For every $A \in \mathcal{A}$ let $\mathcal{M}_b(A, \mathbb{R}^{m \times n})$ be the Banach space of all $\mathbb{R}^{m \times n}$ -valued bounded Radon measures on A . This space is identified with the dual of the space $C_0(A, \mathbb{R}^{m \times n})$ of all $\mathbb{R}^{m \times n}$ -valued continuous functions on \bar{A} vanishing on ∂A . For every $R > 0$ we set

$$\mathcal{M}_{R,A}^{m \times n} := \{\mu \in \mathcal{M}_b(A, \mathbb{R}^{m \times n}) : |\mu|(A) \leq R\},$$

where $|\mu|$ denotes the variation of μ with respect to the Euclidean norm in $\mathbb{R}^{m \times n}$. On $\mathcal{M}_{R,A}^{m \times n}$ we consider the topology induced by the weak* topology of $\mathcal{M}_b(A, \mathbb{R}^{m \times n})$, which will be called the weak* topology on $\mathcal{M}_{R,A}^{m \times n}$. Since $\mathcal{M}_b(A, \mathbb{R}^{m \times n})$ is the dual of a separable Banach space, there exists a distance $d_{R,A}^{m \times n}$ on $\mathcal{M}_{R,A}^{m \times n}$ which induces the weak* topology on $\mathcal{M}_{R,A}^{m \times n}$ (see [23, Theorem V.5.1]). Moreover, the metric space $(\mathcal{M}_{R,A}^{m \times n}, d_{R,A}^{m \times n})$ is compact by the Banach-Alaoglu Theorem.

For every $\mu \in \mathcal{M}_b(A, \mathbb{R}^{m \times n})$ the absolutely continuous part of μ with respect to the Lebesgue measure \mathcal{L}^n is denoted with μ^a . Note that, if $\mu \in \mathcal{M}_{R,A}^{m \times n}$, then $\mu^a \in \mathcal{M}_{R,A}^{m \times n}$.

The following lemma concerns the mesurability properties of the density of μ^a with respect to \mathcal{L}^n .

Lemma A.1 *Let $A \in \mathcal{A}$ and $R > 0$. Then there exists a $\mathcal{B}(A) \otimes \mathcal{B}(\mathcal{M}_{R,A}^{m \times n})$ -measurable function $\gamma : A \times \mathcal{M}_{R,A}^{m \times n} \rightarrow \mathbb{R}^{m \times n}$ such that*

$$\begin{aligned} \gamma(\cdot, \mu) &\in L^1(A, \mathbb{R}^{m \times n}) \quad \text{for every } \mu \in \mathcal{M}_{R,A}^{m \times n}, \\ \mu^a(B) &= \int_B \gamma(x, \mu) dx \quad \text{for every } \mu \in \mathcal{M}_{R,A}^{m \times n} \text{ and } B \in \mathcal{B}(A). \end{aligned} \tag{A.1}$$

Proof For every $(x, \mu) \in A \times \mathcal{M}_{R,A}^{m \times n}$ let $\gamma(x, \mu) \in \mathbb{R}^{m \times n}$ be defined as

$$\gamma(x, \mu) := \begin{cases} \lim_{\rho \rightarrow 0^+} \frac{\mu(B_\rho(x) \cap A)}{\omega_n \rho^n} & \text{if the limit exists in } \mathbb{R}^{m \times n}, \\ 0 & \text{otherwise,} \end{cases}$$

where ω_n denotes the volume of the unit ball of \mathbb{R}^n . From the theory of differentiation of measures (see, e.g., [26, Theorem 1.155]), for every $\mu \in \mathcal{M}_{R,A}^{m \times n}$ we have that $\gamma(\cdot, \mu) \in L^1(A, \mathbb{R}^{m \times n})$ and

$$\mu^a(B) = \int_B \gamma(x, \mu) dx \quad \text{for every } B \in \mathcal{B}(A),$$

which proves (A.1).

To prove the measurability of the function γ it suffices to show that for every $\rho > 0$ the function

$$(x, \mu) \mapsto \mu(B_\rho(x) \cap A) \tag{A.2}$$

from $A \times \mathcal{M}_{R,A}^{m \times n}$ to $\mathbb{R}^{m \times n}$ is $\mathcal{B}(A) \otimes \mathcal{B}(\mathcal{M}_{R,A}^{m \times n})$ -measurable. To this end, for a fixed $\rho > 0$ we introduce an increasing sequence of nonnegative functions $(\varphi_j) \subset C_c(\mathbb{R}^n)$ pointwise converging to the characteristic function of the open ball $B_\rho(0)$, and we observe that

$$\mu(B_\rho(x) \cap A) = \lim_{j \rightarrow +\infty} \int_A \varphi_j(y - x) d\mu(y),$$

by the Monotone Convergence Theorem.

Let $A_\rho := \{x \in A : \text{dist}(x, \partial A) > \rho\}$. Since for every $j \in \mathbb{N}$ the function

$$(x, \mu) \mapsto \int_A \varphi_j(y - x) d\mu(y)$$

is continuous on $A_\rho \times \mathcal{M}_{R,A}^{m \times n}$ (considering on $\mathcal{M}_{R,A}^{m \times n}$ the weak* topology), the function (A.2) from $A_\rho \times \mathcal{M}_{R,A}^{m \times n}$ to $\mathbb{R}^{m \times n}$ is $\mathcal{B}(A_\rho) \otimes \mathcal{B}(\mathcal{M}_{R,A}^{m \times n})$ -measurable. By the arbitrariness of $\rho > 0$ we obtain that the same function considered on $A \times \mathcal{M}_{R,A}^{m \times n}$ is $\mathcal{B}(A) \otimes \mathcal{B}(\mathcal{M}_{R,A}^{m \times n})$ -measurable. \square

To prove the measurability of the map $\mu \mapsto \mu^a$, from $\mathcal{M}_{R,A}^{m \times n}$ to $\mathcal{M}_{R,A}^{m \times n}$, we need the following lemma.

Lemma A.2 *Let $A \in \mathcal{A}$, let $R > 0$, let (Y, \mathcal{E}) be a measurable space, and let $h : A \times Y \rightarrow \mathbb{R}^{m \times n}$ be a $\mathcal{B}(A) \otimes \mathcal{E}$ -measurable function such that*

$$\int_A |h(x, y)| dx \leq R \quad \text{for every } y \in Y.$$

For every $y \in Y$, we define the $\mathbb{R}^{m \times n}$ -valued measure $\lambda_y \in \mathcal{M}_{R,A}^{m \times n}$ as

$$\lambda_y(B) := \int_B h(x, y) dx \quad \text{for every } B \in \mathcal{B}(A).$$

Then the map $y \mapsto \lambda_y$ is measurable from (Y, \mathcal{E}) to $(\mathcal{M}_{R,A}^{m \times n}, \mathcal{B}(\mathcal{M}_{R,A}^{m \times n}))$.

Proof We start by observing that for every $\varphi \in C_c(A, \mathbb{R}^{m \times n})$ the scalar function

$$y \mapsto \int_A \varphi(x) \cdot d\lambda_y(x) \quad \text{is } \mathcal{E}\text{-measurable,} \tag{A.3}$$

where \cdot denotes the Euclidean scalar product between matrices. Indeed, by definition we have

$$\int_A \varphi(x) \cdot d\lambda_y(x) = \int_A \varphi(x) \cdot h(x, y) dx,$$

and the measurability with respect to y follows from the Fubini Theorem.

Note now that a basis for the open sets of the space $\mathcal{M}_{R,A}^{m \times n}$ (endowed with the weak* topology) is given by the collection of sets

$$\left\{ \lambda \in \mathcal{M}_{R,A}^{m \times n} : \left| \int_A \varphi_i(x) \cdot d\lambda(x) - \int_A \varphi_i(x) \cdot d\hat{\lambda}(x) \right| < \eta \text{ for } i = 1, \dots, l \right\},$$

with $\eta > 0$, $\hat{\lambda} \in \mathcal{M}_{R,A}^{m \times n}$, $l \in \mathbb{N}$, and $\varphi_1, \dots, \varphi_l \in C_c(A, \mathbb{R}^{m \times n})$. By (A.3), the pre-image of these sets under the function $y \mapsto \lambda_y$ belongs to \mathcal{E} . This implies that this function is measurable from (Y, \mathcal{E}) to $(\mathcal{M}_{R,A}^{m \times n}, \mathcal{B}(\mathcal{M}_{R,A}^{m \times n}))$, since the weak* topology in $\mathcal{M}_{R,A}^{m \times n}$ has a countable basis. \square

The following lemma shows the measurable dependence of μ^a on μ .

Lemma A.3 *The map $\mu \mapsto \mu^a$ is measurable from $(\mathcal{M}_{R,A}^{m \times n}, \mathcal{B}(\mathcal{M}_{R,A}^{m \times n}))$ to $(\mathcal{M}_{R,A}^{m \times n}, \mathcal{B}(\mathcal{M}_{R,A}^{m \times n}))$.*

Proof Thanks to (A.1), the conclusion follows from Lemma A.2 with $(Y, \mathcal{E}) = (\mathcal{M}_{R,A}^{m \times n}, \mathcal{B}(\mathcal{M}_{R,A}^{m \times n}))$ and $h = \gamma$. □

Given $A \in \mathcal{A}$, we set

$$BV_{R,A}^m := \{u \in BV(A, \mathbb{R}^m) : \|u\|_{L^1(A, \mathbb{R}^m)} \leq R \text{ and } |Du|(A) \leq R\}. \tag{A.4}$$

On $BV_{R,A}^m$ we consider the topology induced by the distance $d_{R,A}^m$ defined by

$$d_{R,A}^m(u, v) := \|u - v\|_{L^1(A, \mathbb{R}^m)} + d_{R,A}^{m \times n}(Du, Dv),$$

where $d_{R,A}^{m \times n}$ is the distance on $\mathcal{M}_{R,A}^{m \times n}$ that metrizes the weak* topology.

Note that $BV(A, \mathbb{R}^m)$ is the dual of a separable space, and that, when A has Lipschitz boundary, the topology just defined coincides with the topology induced on $BV_{R,A}^m$ by the weak* topology of $BV(A, \mathbb{R}^m)$ (see [5, Remark 3.12]).

The following lemma will be crucial in the proof of Proposition A.12.

Lemma A.4 *Assume that $A \in \mathcal{A}$ has Lipschitz boundary. Then the metric space $(BV_{R,A}^m, d_{R,A}^{m \times n})$ is compact.*

Proof Let (u_k) be a sequence in $BV_{R,A}^m$. By (A.4) this sequence is bounded in $BV(A, \mathbb{R}^m)$. Recalling the compact embedding of $BV(A, \mathbb{R}^m)$ into $L^1(A, \mathbb{R}^m)$ and the compactness of $\mathcal{M}_{R,A}^{m \times n}$, there exist a subsequence, not relabelled, and a function $u \in BV(A, \mathbb{R}^m)$ such that $u_k \rightarrow u$ strongly in $L^1(A, \mathbb{R}^m)$ and $Du_k \rightharpoonup Du$ weakly* in $\mathcal{M}_b(A, \mathbb{R}^{m \times n})$. It is easy to see that $u \in BV_{R,A}^m$ and that $d_{R,A}^{m \times n}(u_k, u) \rightarrow 0$. □

We now prove the measurability with respect to (ω, u) of the integral functional corresponding to a random volume integrand.

Lemma A.5 *Let $A \in \mathcal{A}$ with Lipschitz boundary, let $R > 0$, and let f be a random volume integrand as in Definition 3.7. Then, the function*

$$(\omega, u) \mapsto \int_A f(\omega, x, \nabla u) dx$$

from $\Omega \times BV_{R,A}^m$ to \mathbb{R} is $\mathcal{T} \otimes \mathcal{B}(BV_{R,A}^m)$ -measurable.

Proof Let γ be the function introduced in Lemma A.1. We observe that for every $u \in BV_{R,A}^m$

$$\gamma(x, Du) = \nabla u(x) \quad \text{for } \mathcal{L}^n\text{-a.e. } x \in A.$$

Therefore,

$$\int_A f(\omega, x, \nabla u) dx = \int_A f(\omega, x, \gamma(x, Du)) dx.$$

We claim that the function $(x, u) \mapsto \gamma(x, Du)$ from $A \times BV_{R,A}^m$ to $\mathbb{R}^{m \times n}$ is $\mathcal{B}(A) \otimes \mathcal{B}(BV_{R,A}^m)$ -measurable. Indeed, it is the composition of the functions $(x, u) \mapsto (x, Du)$, which is continuous from $A \times BV_{R,A}^m$ to $A \times \mathcal{M}_{R,A}^{m \times n}$, and the function $(x, \mu) \mapsto \gamma(x, \mu)$ from $A \times \mathcal{M}_{R,A}^{m \times n}$ to $\mathbb{R}^{m \times n}$, which is $\mathcal{B}(A) \otimes \mathcal{B}(BV_{R,A}^m)$ -measurable, by Lemma A.1. Therefore, the function $(\omega, x, u) \mapsto (\omega, x, \gamma(x, Du))$ is measurable from $(\Omega \times A \times BV_{R,A}^m, \mathcal{T} \otimes \mathcal{B}(A) \otimes \mathcal{B}(BV_{R,A}^m))$ to $(\Omega \times \mathbb{R}^n \times \mathbb{R}^{m \times n}, \mathcal{T} \otimes \mathcal{B}^n \otimes \mathcal{B}^{m \times n})$. By the $\mathcal{T} \otimes \mathcal{B}^n \otimes \mathcal{B}^{m \times n}$ -measurability of f we deduce that the function $(\omega, x, u) \mapsto f(\omega, x, \gamma(x, Du))$ from $\Omega \times A \times BV_{R,A}^m$ to \mathbb{R} is $\mathcal{T} \otimes \mathcal{B}(A) \otimes \mathcal{B}(BV_{R,A}^m)$ -measurable. The conclusion then follows from Fubini's Theorem. \square

The following two lemmas are used to prove the measurable dependence on u of the surface integral functional corresponding to a continuous surface integrand.

For every $A \in \mathcal{A}$, $\mu \in \mathcal{M}_b(A, \mathbb{R}^{m \times n})$, $x \in A$, and $\rho > 0$ we set

$$\theta_{A,\rho}(\mu, x) := \frac{\mu(B_\rho(x) \cap A)}{\omega_{n-1}\rho^{n-1}}, \tag{A.5}$$

where ω_{n-1} denotes the volume of the unit ball of \mathbb{R}^{n-1} .

Lemma A.6 *Let $A \in \mathcal{A}$ and $u \in BV(A, \mathbb{R}^m)$. Then*

$$\lim_{\rho \rightarrow 0^+} \theta_{A,\rho}(Du, x) = ([u](x) \otimes \nu_u(x)) \chi_{S_u}(x) \text{ for } \mathcal{H}^{n-1}\text{-a.e. } x \in A, \tag{A.6}$$

where $\chi_{S_u}(x) = 1$ if $x \in S_u$ and $\chi_{S_u}(x) = 0$ if $x \in A \setminus S_u$.

Proof *Step 1.* We claim that

$$\lim_{\rho \rightarrow 0^+} \theta_{A,\rho}(Du^d, x) = 0 \text{ for } \mathcal{H}^{n-1}\text{-a.e. } x \in A. \tag{A.7}$$

We now recall that, for a positive Radon measure μ in A and for $d \in \mathbb{N}$, the d -dimensional upper density of μ at $x \in A$ is defined as

$$\Theta^{*,d}(\mu, x) = \limsup_{\rho \rightarrow 0^+} \frac{\mu(B_\rho(x) \cap A)}{\omega_d \rho^d},$$

where ω_d denotes the volume of the unit ball of \mathbb{R}^d (see, e.g., [5, Definition 2.55]). To prove (A.7) it is then sufficient to show that

$$\Theta^{*,n-1}(|D^a u|, x) = 0 \text{ for } \mathcal{H}^{n-1}\text{-a.e. } x \in A. \tag{A.8}$$

To do so, for any $t > 0$ we define the set

$$E_t := \{x \in A : \Theta^{*,n-1}(|D^a u|, x) > t\};$$

note that

$$E_t \subset \{x \in A : \Theta^{*,n}(|D^a u|, x) = +\infty\}.$$

By the Lebesgue Differentiation Theorem we have $\mathcal{L}^n(E_t) = 0$ and, since $|D^a u| < \infty$, we have $|D^a u|(E_t) = 0$.

Since $|D^a u|$ is a finite Radon measure, for every $k \in \mathbb{N}$ there exists an open set $A_k \subset A$ with $E_t \subset A_k$ such that

$$|D^a u|(A_k) < \frac{1}{k}.$$

Thanks to [24, Section 2.10.19(3) and Section 2.10.6] this implies that

$$t\mathcal{H}^{n-1}(E_t) \leq |D^a u|(A_k) < \frac{1}{k} \quad \text{for every } k \in \mathbb{N}.$$

Taking the limit as $k \rightarrow \infty$, we obtain that

$$\mathcal{H}^{n-1}(E_t) = 0 \quad \text{for every } t > 0.$$

From this, it follows that

$$\mathcal{H}^{n-1}(\{x \in A : \Theta^{*,n-1}(|D^a u|, x) > 0\}) = 0$$

and this proves (A.8), which gives (A.7).

Step 2. We claim that

$$\lim_{\rho \rightarrow 0^+} \theta_{A,\rho}(C(u), x) = 0 \quad \text{for } \mathcal{H}^{n-1}\text{-a.e. } x \in A. \tag{A.9}$$

As before, it is sufficient to show that

$$\Theta^{*,n-1}(|C(u)|, x) = 0 \quad \text{for } \mathcal{H}^{n-1}\text{-a.e. } x \in A. \tag{A.10}$$

To do so, for any $t > 0$ we define the set

$$E_t := \{x \in A : \Theta^{*,n-1}(|C(u)|, x) > t\}.$$

Now, let $K \subset E_t$ be a compact set with $\mathcal{H}^{n-1}(K) < +\infty$ so that, in particular, $|C(u)|(K) = 0$. Then, by [24, Section 2.10.19(3) and Section 2.10.6] we have that

$$t\mathcal{H}^{n-1}(K) \leq |C(u)|(V) \quad \text{for every open set } V \text{ containing } K.$$

Since $C(u)$ is a finite Radon measure, taking the infimum of the above inequality over all open sets V containing K we obtain that

$$t\mathcal{H}^{n-1}(K) \leq |C(u)|(K).$$

Since $|C(u)|(K) = 0$, from the above inequality it follows that $\mathcal{H}^{n-1}(K) = 0$. Using the fact that E_t is a Borel (and hence Suslin) set, by [24, Corollary 2.10.48] we have that

$$\mathcal{H}^{n-1}(E_t) = \sup\{\mathcal{H}^{n-1}(K) : K \text{ compact, } K \subset E_t, \mathcal{H}^{n-1}(K) < +\infty\},$$

and so $\mathcal{H}^{n-1}(E_t) = 0$ for every $t > 0$, which implies (A.10) and, in turn, (A.9).

Step 3. We claim that

$$\lim_{\rho \rightarrow 0^+} \theta_{A,\rho}(D^j u, x) = 0 \text{ for } \mathcal{H}^{n-1}\text{-a.e. } x \in A \setminus S_u. \tag{A.11}$$

Observe that $|D^j u|(A \setminus S_u) = 0$. By [24, Section 2.10.19(4) and Section 2.10.6] we have immediately that

$$\Theta^{*,n-1}(|D^j u|, x) = 0 \text{ for } \mathcal{H}^{n-1}\text{-a.e. } x \in A \setminus S_u,$$

which implies (A.11).

Step 4. By Besicovich Derivation Theorem (see [5, Theorems 2.22, 2.83, and 3.78]) we have that

$$\lim_{\rho \rightarrow 0^+} \theta_{A,\rho}(D^j u, x) = [u](x) \otimes \nu_u(x) \text{ for } \mathcal{H}^{n-1}\text{-a.e. } x \in S_u.$$

Together with the previous steps, this gives (A.6). □

Lemma A.7 *Let $A \in \mathcal{A}$ and let $g : A \times \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$ be a continuous function. Assume that there exists a $a > 0$ such that*

$$|g(x, \xi)| \leq a|\xi| \tag{A.12}$$

for every $(x, \xi) \in A \times \mathbb{R}^{m \times n}$. Then for every $u \in BV(A, \mathbb{R}^m)$

$$\lim_{\eta \rightarrow 0^+} \lim_{\rho \rightarrow 0^+} \int_A \frac{g(x, \theta_{A,\rho}(Du, x))}{|\theta_{A,\rho}(Du, x)| \vee \eta} d|Du|(x) = \int_{A \cap S_u} g(x, [u](x) \otimes \nu_u(x)) d\mathcal{H}^{n-1}(x),$$

where $\theta_{A,\rho}$ is defined in (A.5).

Proof Let $u \in BV(A, \mathbb{R}^m)$ be fixed.

Step 1. Thanks to (A.12) and to the bound

$$\int_{A \cap S_u} |[u](x)| d\mathcal{H}^{n-1}(x) < +\infty, \tag{A.13}$$

the function $x \mapsto g(x, [u](x) \otimes \nu_u(x))$ is \mathcal{H}^{n-1} -integrable on $A \cap S_u$.

Step 2. We claim that for every $\eta > 0$

$$\lim_{\rho \rightarrow 0^+} \int_A \frac{g(x, \theta_{A,\rho}(Du, x))}{|\theta_{A,\rho}(Du, x)| \vee \eta} d|D^a u + C(u)|(x) = 0.$$

By Lemma A.6 we have that

$$\lim_{\rho \rightarrow 0^+} \theta_{A,\rho}(Du, x) = 0 \text{ for } |D^a u + C(u)|\text{-a.e. } x \in A,$$

and by (A.12) we have the inequality

$$\frac{|g(x, \theta_{A,\rho}(Du, x))|}{|\theta_{A,\rho}(Du, x)| \vee \eta} \leq a. \tag{A.14}$$

The claim then follows from the Dominated Convergence Theorem, since g is continuous, $g(x, 0) = 0$, and $|D^a u + C(u)|$ is a bounded measure.

Step 3. Recalling (f) in Section 2, to conclude the proof it is sufficient to show that

$$\begin{aligned} & \lim_{\eta \rightarrow 0^+} \lim_{\rho \rightarrow 0^+} \int_{A \cap S_u} \frac{g(x, \theta_{A,\rho}(Du, x))}{|\theta_{A,\rho}(Du, x)| \vee \eta} |[u](x) \otimes v_u(x)| d\mathcal{H}^{n-1}(x) \\ &= \int_{A \cap S_u} g(x, [u](x) \otimes v_u(x)) d\mathcal{H}^{n-1}(x). \end{aligned}$$

By Lemma A.6 and by the continuity of g we have that for every $\eta > 0$

$$\begin{aligned} & \lim_{\rho \rightarrow 0^+} \int_{A \cap S_u} \frac{g(x, \theta_{A,\rho}(Du, x))}{|\theta_{A,\rho}(Du, x)| \vee \eta} |[u](x) \otimes v_u(x)| d\mathcal{H}^{n-1}(x) \\ &= \int_{A \cap S_u} g(x, [u](x) \otimes v_u(x)) \frac{|[u](x) \otimes v_u(x)|}{|[u](x) \otimes v_u(x)| \vee \eta} d\mathcal{H}^{n-1}(x), \end{aligned}$$

where we used the Dominated Convergence Theorem, thanks to (A.13) and (A.14). Note that for \mathcal{H}^{n-1} -almost every $x \in S_u$

$$\lim_{\eta \rightarrow 0^+} \frac{|[u](x) \otimes v_u(x)|}{|[u](x) \otimes v_u(x)| \vee \eta} = \sup_{\eta > 0} \frac{|[u](x) \otimes v_u(x)|}{|[u](x) \otimes v_u(x)| \vee \eta} = 1,$$

since $[u](x) \neq 0$. Thanks to (A.12) and (A.13) we can apply again the Dominated Convergence Theorem and deduce the claim in the limit $\eta \rightarrow 0^+$. \square

We are now in a position to prove the measurable dependence on u of the integral functional corresponding to a continuous surface integrand.

Lemma 9.6 *Let $A \in \mathcal{A}$ with Lipschitz boundary, let $R > 0$, and let g be as in Lemma A.7. Then the function*

$$u \mapsto \int_{S_u \cap A} g(x, [u](x) \otimes v_u(x)) d\mathcal{H}^{n-1}(x)$$

from $BV_{R,A}^m$ to \mathbb{R} is $\mathcal{B}(BV_{R,A}^m)$ -measurable.

Proof By Lemma A.7 the thesis follows by proving that for every $\rho, \eta > 0$ the function

$$u \mapsto \int_A \frac{g(x, \theta_{A,\rho}(Du, x))}{|\theta_{A,\rho}(Du, x)| \vee \eta} d|Du|(x) \tag{A.15}$$

from $BV_{R,A}^m$ to \mathbb{R} is $\mathcal{B}(BV_{R,A}^m)$ -measurable. Let $\rho, \eta > 0$ be fixed. First note that, by the $\mathcal{B}(A) \otimes \mathcal{B}(\mathcal{M}_{R,A}^{m \times n})$ -measurability of (A.2) and the continuity of g , the function

$$(x, \mu) \mapsto \frac{g(x, \theta_{A,\rho}(\mu, x))}{|\theta_{A,\rho}(\mu, x)| \vee \eta}$$

is $\mathcal{B}(A) \otimes \mathcal{B}(\mathcal{M}_{R,A}^{m \times n})$ -measurable. Moreover it is bounded by (A.14). So by [17, Corollary A.3]

$$\mu \mapsto \int_A \frac{g(x, \theta_{A,\rho}(\mu, x))}{|\theta_{A,\rho}(\mu, x)| \vee \eta} d|\mu|(x)$$

is $\mathcal{B}(\mathcal{M}_{R,A}^{m \times n})$ -measurable. Since $u \mapsto Du$ is continuous from $(BV_{R,A}^m, d_{R,A}^m)$ to $(\mathcal{M}_{R,A}^{m \times n}, d_{R,A}^{m \times n})$, the $\mathcal{B}(BV_{R,A}^m)$ -measurability of (A.15) follows. \square

We now prove the measurability with respect to (ω, u) of the integral functional corresponding to a random surface integrand, with no continuity assumption with respect to x .

Lemma A.8 *Let $A \in \mathcal{A}$ with Lipschitz boundary, let $R > 0$, and let $g: \Omega \times A \times \mathbb{R}^m \times \mathbb{S}^{n-1} \rightarrow \mathbb{R}$ be a $\mathcal{T} \otimes \mathcal{B}(A) \otimes \mathcal{B}^m \otimes \mathcal{B}_S^n$ -measurable function. Assume that there exists $a > 0$ such that*

$$g(\omega, x, \zeta, \nu) = g(\omega, x, -\zeta, -\nu), \tag{A.16}$$

$$|g(\omega, x, \zeta, \nu)| \leq a|\zeta|, \tag{A.17}$$

for every $(\omega, x, \zeta, \nu) \in \Omega \times A \times \mathbb{R}^m \times \mathbb{S}^{n-1}$. Then the function

$$(\omega, u) \mapsto \int_{S_u \cap A} g(\omega, x, [u](x), \nu_u(x)) d\mathcal{H}^{n-1}(x)$$

from $\Omega \times BV_{R,A}^m$ to \mathbb{R} is $\mathcal{T} \otimes \mathcal{B}(BV_{R,A}^m)$ -measurable.

Proof We recall that a matrix $\xi \in \mathbb{R}^{m \times n}$ has rank ≤ 1 if and only if $\xi = \zeta \otimes \nu$ for some $\zeta \in \mathbb{R}^m$ and $\nu \in \mathbb{S}^{n-1}$, and that the pair (ζ, ν) is uniquely determined by ξ , up to a change of sign of both terms. Therefore, thanks to (A.16), we can define a $\mathcal{T} \otimes \mathcal{B}(A) \otimes \mathcal{B}^{m \times n}$ -measurable function $\tilde{g}: \Omega \times A \times \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$ by setting for every $(\omega, x, \xi) \in \Omega \times A \times \mathbb{R}^{m \times n}$

$$\tilde{g}(\omega, x, \xi) := \begin{cases} g(\omega, x, \zeta, \nu) & \text{if } \xi = \zeta \otimes \nu, \text{ with } \zeta \in \mathbb{R}^m \text{ and } \nu \in \mathbb{S}^{n-1}, \\ 0 & \text{if } \text{rank}(\xi) > 1. \end{cases}$$

By (A.17) we have $|\tilde{g}(\omega, x, \xi)| \leq a|\xi|$ for every $(\omega, x, \xi) \in \Omega \times A \times \mathbb{R}^{m \times n}$.

To prove the thesis it is enough to show that

$$(\omega, u) \mapsto \int_{S_u \cap A} \tilde{g}(\omega, x, [u](x) \otimes \nu_u(x)) d\mathcal{H}^{n-1} \text{ is } \mathcal{T} \otimes \mathcal{B}(BV_{R,A}^m)\text{-measurable.} \tag{A.18}$$

Note that the function \tilde{g} can be written as

$$\tilde{g}(\omega, x, \xi) = \hat{g}(\omega, x, \xi)a|\xi|, \tag{A.19}$$

where \hat{g} is $\mathcal{T} \otimes \mathcal{B}(A) \otimes \mathcal{B}(\mathbb{R}^{m \times n})$ -measurable and satisfies $|\hat{g}| \leq 1$.

Let \mathcal{R} be the set of all bounded $\mathcal{T} \otimes \mathcal{B}(A) \otimes \mathcal{B}^{m \times n}$ -measurable functions $\hat{g}: \Omega \times A \times \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$ such that the function \tilde{g} defined as in (A.19) satisfies the claim (A.18).

In order to conclude the proof, we need to show that \mathcal{R} contains all bounded $\mathcal{T} \otimes \mathcal{B}(A) \otimes \mathcal{B}^{m \times n}$ -measurable functions. To prove this property, note that \mathcal{R} is a vector space of bounded real-valued functions that contains the constants and is closed both under uniform convergence and under monotone convergence of uniformly bounded sequences. Let \mathcal{C} be the set of all functions $\hat{g}: \Omega \times A \times \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$ that can be written as

$$\hat{g}(\omega, x, \xi) = \alpha(\omega)\beta(x, \xi),$$

where $\alpha: \Omega \rightarrow \mathbb{R}$ is bounded and \mathcal{T} -measurable, and $\beta: A \times \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$ is bounded and continuous. Note that \mathcal{C} is stable under multiplication and that the σ -algebra generated by \mathcal{C} is $\mathcal{T} \otimes \mathcal{B}(A) \otimes \mathcal{B}^{m \times n}$.

By Lemma 9.6 we have $\mathcal{C} \subset \mathcal{R}$. Hence the functional form of the Monotone Class Theorem (see [22, Chapter I, Theorem 21]), implies that \mathcal{R} contains all bounded $\mathcal{T} \otimes \mathcal{B}(A) \otimes \mathcal{B}^{m \times n}$ -measurable functions, and this concludes the proof. \square

We now prove the measurability of the map $u \mapsto D^j u$.

Lemma A.9 *Let $A \in \mathcal{A}$ with Lipschitz boundary and let $R > 0$. Then the map*

$$u \mapsto D^j u$$

is measurable from $(BV_{R,A}^m, \mathcal{B}(BV_{R,A}^m))$ to $(\mathcal{M}_{R,A}^{m \times n}, \mathcal{B}(\mathcal{M}_{R,A}^{m \times n}))$.

Proof As in the proof of Lemma A.2 it is sufficient to show that

$$u \mapsto \int_A \varphi(x) \cdot d(D^j u)(x) = \int_{A \cap S_u} \varphi(x) \cdot ([u](x) \otimes \nu_u(x)) d\mathcal{H}^{n-1}(x)$$

from $BV_{R,A}^m$ to \mathbb{R} is $\mathcal{B}(BV_{R,A}^m)$ -measurable for every $\varphi \in C_c(A, \mathbb{R}^{m \times n})$. To this end we set $g(x, \zeta, \nu) := \varphi(x) \cdot (\zeta \otimes \nu)$, and note that $|g(x, \zeta, \nu)| \leq a|\zeta|$, where a is the

maximum of $|\varphi|$. Therefore, by Lemma A.8 it follows that

$$u \mapsto \int_{S_u \cap A} g(x, [u](x), \nu_u(x)) d\mathcal{H}^{n-1}(x)$$

from $BV_{R,A}^m$ to \mathbb{R} is $\mathcal{B}(BV_{R,A}^m)$ -measurable, and hence the claim. □

The following corollary deals with the Cantor part.

Corollary A.11 *Let $A \in \mathcal{A}$ with Lipschitz boundary and let $R > 0$. Then the map*

$$u \mapsto C(u)$$

is measurable from $(BV_{R,A}^m, \mathcal{B}(BV_{R,A}^m))$ to $(\mathcal{M}_{R,A}^{m \times n}, \mathcal{B}(\mathcal{M}_{R,A}^{m \times n}))$.

Proof Since $C(u) = Du - D^a u - D^j u$, the result follows from the continuity of the map $u \mapsto Du$ from $(BV_{R,A}^m, d_{R,A}^m)$ to $(\mathcal{M}_{R,A}^{m \times n}, d_{R,A}^{m \times n})$, using Lemmas A.3 and A.9. □

We are now ready to prove the main result of the section.

Proposition A.12 *Let f and g be random volume and surface integrands, respectively, according to Definition 3.7, and let $(\Omega, \widehat{\mathcal{T}}, \widehat{P})$ be the completion of the probability space (Ω, \mathcal{T}, P) . Let $A \in \mathcal{A}$, let $w \in SBV(A, \mathbb{R}^m)$, and for every $\omega \in \Omega$ let $m_\omega^{f,g}(w, A)$ be defined as in (3.4) and (3.16). Then the function $\omega \mapsto m_\omega^{f,g}(w, A)$ is $\widehat{\mathcal{T}}$ -measurable.*

Proof For every $\omega \in \Omega$, $B \in \mathcal{B}(A)$, and $u \in BV(A, \mathbb{R}^m)$, we define

$$E(\omega)(u, B) := \int_B f(\omega, x, \nabla u) dx + \int_{S_u \cap B} g(\omega, x, [u], \nu) d\mathcal{H}^{n-1}.$$

Let us fix a sequence (A_j) of open sets with Lipschitz boundary, with $A_j \subset\subset A_{j+1}$ for every $j \in \mathbb{N}$ and $\cup_j A_j = A$. It follows easily from the definition that

$$\begin{aligned} & m_\omega^{f,g}(w, A) \\ &= \lim_{j \rightarrow +\infty} \inf\{E(\omega)(u, A) : u \in SBV(A, \mathbb{R}^m), u = w \text{ in } A \setminus A_j\} \\ &= \lim_{j \rightarrow +\infty} \left(\inf\{E(\omega)(u, A_{j+1}) : u \in SBV(A_{j+1}, \mathbb{R}^m), u = w \text{ in } A_{j+1} \setminus A_j\} \right. \\ & \quad \left. + E(\omega)(w, A \setminus A_{j+1}) \right) \\ &= \lim_{j \rightarrow +\infty} \inf\{E(\omega)(u, A_{j+1}) : u \in SBV(A_{j+1}, \mathbb{R}^m), u = w \text{ in } A_{j+1} \setminus A_j\}, \end{aligned}$$

where in the last equality we used the fact that $E(\omega)(w, A \setminus A_{j+1}) \rightarrow 0$ as $j \rightarrow +\infty$ since, by (f4) and (g4), we have $E(\omega)(w, A \setminus A_{j+1}) \leq c_3 |Dw|(A \setminus A_{j+1}) + c_4 \mathcal{L}^n(A \setminus A_{j+1})$.

Let us fix $j \in \mathbb{N}$. It is obvious that

$$\begin{aligned} & \inf\{E(\omega)(u, A_{j+1}) : u \in SBV(A_{j+1}, \mathbb{R}^m), u = w \text{ in } A_{j+1} \setminus A_j\} \\ &= \inf\{E(\omega)(u, A_{j+1}) : u \in SBV(A_{j+1}, \mathbb{R}^m), E(\omega)(u, A_{j+1}) \\ &\leq E(\omega)(w, A_{j+1}), u = w \text{ in } A_{j+1} \setminus A_j\}. \end{aligned}$$

By (f4) and (g4) we have $E(\omega)(w, A_{j+1}) \leq c_3|Dw|(A_{j+1}) + c_4\mathcal{L}^n(A_{j+1})$. Therefore, from Remark 3.2 we obtain that there exists $R_1 > 0$, depending on A_{j+1} and w , such that

$$\begin{aligned} & \inf\{E(\omega)(u, A_{j+1}) : u \in SBV(A_{j+1}, \mathbb{R}^m), u = w \text{ in } A_{j+1} \setminus A_j\} \\ &= \inf\{E(\omega)(u, A_{j+1}) : u \in SBV(A_{j+1}, \mathbb{R}^m), \\ &|Du|(A_{j+1}) \leq R_1, u = w \text{ in } A_{j+1} \setminus A_j\}. \end{aligned}$$

Thanks to Poincaré’s inequality, there exists $R \geq R_1$, depending on A_{j+1} , w , and R_1 , such that every function $u \in BV(A_{j+1}, \mathbb{R}^m)$, satisfying $|Du|(A_{j+1}) \leq R_1$ and $u = w$ in $A_{j+1} \setminus A_j$, satisfies also $\|u\|_{L^1(A_{j+1}, \mathbb{R}^m)} \leq R$. This implies that

$$\begin{aligned} & \inf\{E(\omega)(u, A_{j+1}) : u \in SBV(A_{j+1}, \mathbb{R}^m), u = w \text{ in } A_{j+1} \setminus A_j\} \\ &= \inf\{E(\omega)(u, A_{j+1}) : u \in SBV(A_{j+1}, \mathbb{R}^m), \|u\|_{L^1(A_{j+1}, \mathbb{R}^m)} \leq R, \\ &|Du|(A_{j+1}) \leq R, u = w \text{ in } A_{j+1} \setminus A_j\}. \end{aligned}$$

Therefore, to prove the proposition it is enough to show that the function

$$\omega \mapsto \inf\{E(\omega)(u, A_{j+1}) : u \in SBV(A_{j+1}, \mathbb{R}^m) \cap BV_{R, A_{j+1}}^m, u = w \text{ in } A_{j+1} \setminus A_j\} \tag{A.20}$$

is $\widehat{\mathcal{T}}$ -measurable.

We define $H : BV_{R, A_{j+1}}^m \rightarrow [0, +\infty]$ as

$$H(u) := \begin{cases} 0 & \text{if } C(u) = 0 \text{ and } u = w \text{ in } A_{j+1} \setminus A_j, \\ +\infty & \text{otherwise,} \end{cases}$$

where the equality $C(u) = 0$ means that $C(u)(B) = 0$ for every $B \in \mathcal{B}(A_{j+1})$. By (A.20) to conclude the proof it suffices to show that the function

$$\omega \mapsto \inf\{E(\omega)(u, A_{j+1}) + H(u) : u \in BV_{R, A_{j+1}}^m\} \tag{A.21}$$

is $\widehat{\mathcal{T}}$ -measurable.

To this aim, we apply the Projection Theorem. Note that the function $(\omega, u) \mapsto E(\omega)(u, A_{j+1})$ from $\Omega \times BV_{R, A_{j+1}}^m$ to \mathbb{R} is $\mathcal{T} \otimes \mathcal{B}(BV_{R, A_{j+1}}^m)$ -measurable, by Lemmas A.5 and A.8. Moreover, the function $u \mapsto H(u)$ from $BV_{R, A_{j+1}}^m$ to \mathbb{R} is

$\mathcal{B}(BV_{R,A_{j+1}}^m)$ -measurable, since the set $\{u \in BV_{R,A_{j+1}}^m : C(u) = 0\}$ belongs to $\mathcal{B}(BV_{R,A_{j+1}}^m)$ by Corollary A.11, while the set $\{u \in BV_{R,A_{j+1}}^m : u = w \text{ in } A_{j+1} \setminus A_j\}$ is closed in $(BV_{R,A_{j+1}}^m, d_{R,A_{j+1}}^m)$. Hence, for every $t > 0$ we have

$$\{(\omega, u) \in \Omega \times BV_{R,A_{j+1}}^m : E(\omega)(u, A_{j+1}) + H(u) < t\} \in \mathcal{T} \otimes \mathcal{B}(BV_{R,A_{j+1}}^m). \quad (\text{A.22})$$

Since the metric space $(BV_{R,A_{j+1}}^m, d_{R,A_{j+1}}^m)$ is compact thanks to Lemma A.4, by the Projection Theorem (see, e.g., [22, Theorem III.13 and 33(a)]) the projection onto Ω of the set above belongs to $\widehat{\mathcal{T}}$. On the other hand, the projection onto Ω of the set in (A.22) coincides with the set of points $\omega \in \Omega$ such that

$$\inf\{E(\omega)(u, A_{j+1}) + H(u) : u \in BV_{R,A_{j+1}}^m\} < t.$$

Since this set belongs to $\widehat{\mathcal{T}}$ for every $t > 0$, the function in (A.21) is $\widehat{\mathcal{T}}$ -measurable. \square

References

1. Akcoglu, M.A., Krengel, U.: Ergodic theorems for superadditive processes. *J. Reine Angew. Math.* **323**, 53–67 (1981)
2. Alberti, G.: Rank one property for derivatives of functions with bounded variation. *Proc. Roy. Soc. Edinburgh Sect. A* **123**, 239–274 (1993)
3. Alicandro, R., Cicalese, M., Ruf, M.: Domain formation in magnetic polymer composites: an approach via stochastic homogenization. *Arch. Ration. Mech. Anal.* **218**, 945–984 (2015)
4. Ambrosio, L., Dal Maso, G.: On the relaxation in $BV(\Omega; \mathbb{R}^m)$ of quasi-convex integrals. *J. Funct. Anal.* **109**, 76–97 (1992)
5. Ambrosio, L., Fusco, N., Pallara, D.: *Functions of Bounded Variation and Free Discontinuity Problems*. Clarendon Press, Oxford (2000)
6. Barchiesi, M., Dal Maso, G.: Homogenization of fiber reinforced brittle materials: the extremal cases. *SIAM J. Math. Anal.* **41**, 1874–1889 (2009)
7. Barchiesi, M., Focardi, M.: Homogenization of the Neumann problem in perforated domains: an alternative approach. *Calc. Var. Partial Differential Equations* **42**, 257–288 (2011)
8. Barchiesi, M., Lazzaroni, G., Zeppieri, C.I.: A bridging mechanism in the homogenisation of brittle composites with soft inclusions. *SIAM J. Math. Anal.* **48**, 1178–1209 (2016)
9. Barroso, A., Bouchitté, G., Buttazzo, G., Fonseca, I.: Relaxation of bulk and interfacial energies. *Arch. Ration. Mech. Anal.* **135**, 103–173 (1996)
10. Bouchitté, G., Fonseca, I., Mascarenhas, L.: A global method for relaxation. *Arch. Ration. Mech. Anal.* **145**, 51–98 (1998)
11. Braides, A., Defranceschi, A., Vitali, E.: Homogenization of free discontinuity problems. *Arch. Ration. Mech. Anal.* **135**, 297–356 (1996)
12. Braides, A., Maslennikov, M., Sigalotti, L.: Homogenization by blow-up. *Applicable Anal.* **87**, 1341–1356 (2008)
13. Braides, A., Piatnitski, A.: Homogenization of surface and length energies for spin systems. *J. Funct. Anal.* **264**, 1296–1328 (2013)
14. Braides, A., Solci, M.: Multi-scale free-discontinuity problems with soft inclusions. *Boll. Unione Mat. Ital.* (9) **6**, no. 1, 29–51 (2013)
15. Cagnetti, F., Scardia, L.: An extension theorem in SBV and an application to the homogenization of the Mumford-Shah functional in perforated domains. *J. Math. Pures Appl.* **95**, 349–381 (2011)
16. Cagnetti, F., Dal Maso, G., Scardia, L., Zeppieri, C.I.: Γ -convergence of free-discontinuity problems. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **36**, 1035–1079 (2019)

17. Cagnetti, F., Dal Maso, G., Scardia, L., Zeppieri, C.I.: Stochastic homogenisation of free-discontinuity problems. *Arch. Ration. Mech. Anal.* **233**, 935–974 (2019)
18. Dal Maso, G.: An introduction to Γ -convergence. Birkhäuser, Boston (1993)
19. Dal Maso, G., Modica, L.: Nonlinear stochastic homogenization and ergodic theory. *J. Reine Angew. Math.* **368**, 28–42 (1986)
20. Dal Maso, G., Zeppieri, C.I.: Homogenization of fiber reinforced brittle materials: the intermediate case. *Adv. Calc. Var.* **3**, 345–370 (2010)
21. De Giorgi, E., Ambrosio L.: New functionals in the calculus of variations. (Italian) *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.* (8) **82** (1988), no. 2, 199–210 (1989)
22. Dellacherie, C., Meyer, P.-A.: *Probabilities and Potential*. North-Holland Publishing Company, Amsterdam (1978)
23. Dunford, N., Schwartz, J.T.: *Linear Operators*. Interscience, New York (1957)
24. Federer, H.: *Geometric Measure Theory*. Springer-Verlag, Berlin Heidelberg (1996)
25. Focardi, M., Gelli, M.S., Ponsiglione, M.: Fracture mechanics in perforated domains: a variational model for brittle porous media. *Math. Models Methods Appl. Sci.* **19**, 2065–2100 (2009)
26. Fonseca, I., Leoni, G.: *Modern Methods in the Calculus of Variations: L^p Spaces*. Springer, New York (2007)
27. Fonseca, I., Müller, S.: Quasiconvex integrands and lower semicontinuity in L^1 . *SIAM J. Math. Anal.* **23**, 1081–1098 (1992)
28. Giacomini, A., Ponsiglione, M.: A Γ -convergence approach to stability of unilateral minimality properties. *Arch. Ration. Mech. Anal.* **180**, 399–447 (2006)
29. Larsen, C.J.: Quasiconvexification in $W^{1,1}$ and optimal jump microstructure in BV relaxation. *SIAM J. Math. Anal.* **29**, 823–848 (1998)
30. Licht, C., Michaille, G.: Global-Local subadditive ergodic theorems and application to homogenisation in elasticity. *Ann. Math. Blaise Pascal* **9**, 21–62 (2002)
31. Morse, A.P.: Perfect blankets. *Trans. Amer. Math. Soc.* **61**, 418–442 (1947)
32. Pellet, X., Scardia, L., Zeppieri, C.I.: Homogenization of high-contrast Mumford-Shah energies. *SIAM J. Math. Anal.* **51**, 1696–1729 (2019)
33. Pellet X., Scardia L., and Zeppieri C.I.: Stochastic homogenisation of free-discontinuity functionals in random perforated domains, submitted. Arxiv preprint available at [arXiv:2002.01389](https://arxiv.org/abs/2002.01389)
34. Scardia, L.: Damage as Γ -limit of microfractures in anti-plane linearized elasticity. *Math. Models Methods Appl. Sci.* **18**, 1703–1740 (2008)
35. Scardia, L.: Damage as the Γ -limit of microfractures in linearized elasticity under the non-interpenetration constraint. *Adv. Calc. Var.* **3**, 423–458 (2010)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.