



SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI

SISSA Digital Library

A “noisy” electrical stimulation protocol favors muscle regeneration in vitro through release of endogenous ATP

*Original*

A “noisy” electrical stimulation protocol favors muscle regeneration in vitro through release of endogenous ATP / Bosutti, A.; Bernareggi, A.; Massaria, G.; D'Andrea, P.; Taccola, G.; Lorenzon, P.; Sciancalepore, M.. - In: EXPERIMENTAL CELL RESEARCH. - ISSN 0014-4827. - 381:1(2019), pp. 121-128. [10.1016/j.yexcr.2019.05.012]

*Availability:*

This version is available at: 20.500.11767/95474 since: 2019-06-28T08:20:27Z

*Publisher:*

*Published*

DOI:10.1016/j.yexcr.2019.05.012

*Terms of use:*

Testo definito dall'ateneo relativo alle clausole di concessione d'uso

*Publisher copyright*  
Elsevier

This version is available for education and non-commercial purposes.

note finali coverpage

(Article begins on next page)

# **A “noisy” electrical stimulation protocol favors muscle regeneration *in vitro* through release of endogenous ATP**

Alessandra Bosutti<sup>a</sup>, Annalisa Bernareggi<sup>a</sup>, Gabriele Massaria<sup>a,b</sup>, Paola D’Andrea<sup>a</sup>, Giuliano Taccola<sup>c,d</sup>, Paola Lorenzon<sup>a</sup> and Marina Sciancalepore<sup>a</sup>

<sup>a</sup>Department of Life Sciences and Centre for Neuroscience B.R.A.I.N., University of Trieste, Via A. Fleming 22, I-34127 Trieste, Italy

<sup>b</sup>Area Science Park, Padriciano, 99, I-34149 Trieste, Italy

<sup>c</sup>Department of Neuroscience, SISSA, Via Bonomea 265, 34136 Trieste, Italy

<sup>d</sup>SPINAL (Spinal Person Injury Neurorehabilitation Applied Laboratory), Istituto di Medicina Fisica e Riabilitazione (IMFR), via Gervasutta 48, 33100 Udine, Italy.

**Short title:** EMGstim favors muscle regeneration *in vitro*

**Correspondence to:** Marina Sciancalepore, PhD

Department of Life Sciences, University of Trieste, Via Fleming, 22, 34127 Trieste, Italy

Email: [msciancalepore@units.it](mailto:msciancalepore@units.it)

Tel. +390405588613

## ABSTRACT

An *in vitro* system of electrical stimulation was used to explore whether an innovative “noisy” stimulation protocol derived from human electromyographic recordings (EMG*stim*) could promote muscle regeneration. EMG*stim* was delivered to cultured mouse myofibers isolated from *Flexor Digitorum Brevis*, preserving their satellite cells. In response to EMG*stim*, immunostaining for the myogenic regulatory factor myogenin, revealed an increased percentage of elongated myogenin-positive cells surrounding the myofibers. Conditioned medium collected from EMG*stim*-treated cell cultures, promoted satellite cells differentiation in unstimulated myofiber cell cultures, suggesting that extracellular soluble factors could mediate the process. Interestingly, the myogenic effect of EMG*stim* was mimicked by exogenously applied ATP (0.1  $\mu$ M), reduced by the ATP diphosphohydrolase apyrase and prevented by blocking endogenous ATP release with carbenoxolone.

In conclusion, our results show that “noisy” electrical stimulations favor muscle progenitor cell differentiation most likely via the release of endogenous ATP from contracting myofibres. Our data also suggest that “noisy” stimulation protocols could be potentially more efficient than regular stimulations to promote *in vivo* muscle regeneration after traumatic injury or in neuropathological diseases.

**Keywords:** Differentiation, Electrical stimulation, Skeletal muscle, Myofibers, ATP, Satellite cells.

### Abbreviations:

AU	Arbitrary Units
AUC	Area Under the Curve
CBX	Carbenoxolone
CM	Conditioned Medium
DAPI	4',6-diamidino-2-phenylindole
EMG <i>stim</i>	Electromyographic stimulation
ES	Electrical Stimulation
FDB	Flexor Digitorum Brevis
SC	Satellite Cell

## Introduction

Direct electrical stimulation (ES) of muscles has been demonstrated to mimic nerve activity and therefore has been proposed as a tool to counteract muscle atrophy and to enhance muscle strength in patients suffering from chronic debilitating conditions. Benefits depend on the efficacy of the stimulation patterns and the most appropriate stimulation parameters are still under debate [1,2]. We have already demonstrated that a “noisy” ES protocol derived from human electromyographic recordings, named *EMGstim*, is capable of eliciting *in vitro* mouse myotube contractions more efficiently than “regular” conventional stimulation protocols. More specifically, we observed that in cultured myotubes, *EMGstim*, characterized by biphasic voltage pulses of various duration, amplitude and frequency, was able to elicit action potentials and  $\text{Ca}^{2+}$  release from the sarcoplasmic reticulum of cultured myotubes *in vitro*, at a lower stimulus strength than regular 1 Hz, 1 ms pulse stimulations. These results suggest that *EMGstim*, as a “noisy” protocol, could be potentially more effective in eliciting contractile activity in human patients, thereby limiting the common occurrence of pain and fatigue during ES [3]. Although the effect of ES protocols has been quite extensively studied at the level of skeletal muscle fibers [4-6], the effects on the regenerative response of satellite cells (SCs), the resident myogenic precursors, still remains unknown. SCs within adult skeletal muscle represent a heterogeneous population of undifferentiated mononuclear cells, located under the basal lamina of each myofiber [7]. Their number depends on the species, age and muscle fiber type. Even if they constitute around 5% of the muscle nuclei in the adult mice fiber [8], SCs are important for the maintenance and regeneration of skeletal muscle. Activation of SCs by external stimuli like exercise or injury induces cell proliferation and differentiation, recapitulating the embryonic and fetal program of muscle differentiation.

Satellite cell activation, proliferation and differentiation can be studied in isolated and cultured mouse FDB myofibers [9,10]. After muscle cell plating, SCs detach from the myofibers, divide, migrate and fuse into myotubes in 4-6 days. The different phases of SC myogenesis can be identified by evaluating the expression of specific transcription factors [11].

In the present work, *EMGstim* was delivered to isolated mouse FDB myofibers along with their resident SCs. Immunolabeling for Pax7 and myogenin were then carried out to monitor the effect of *EMGstim* on SC number and differentiation, respectively.

Our results show that, in the absence of motor and sensory innervation, the muscle cell activity promoted by *EMGstim* favors myogenic progenitor differentiation. A molecular mechanism based on the release of endogenous ATP from myofibers during the *EMGstim*-induced contractile activity is proposed.

## Materials and methods

### *FDB myofiber culture*

Skeletal mouse muscle fibers with the resident SCs were obtained from the dissociation of *Flexor Digitorum Brevis* (FDB) muscles of 6 to 8-weeks old C57BL/6J male mice [12,13].

For muscle dissection, mice were anesthetized and sacrificed by cervical dislocation as approved by local Animal Care Committee and in agreement with the European legislation (2010/63/EU). FDB myofibers were isolated from both hind feet muscles of a single mouse for each preparation. Briefly, FDB muscles were enzymatically treated for 1 h in ice and 1 h at 37 °C with Type I collagenase 0.3 % (wt/v), in Tyrode's solution supplemented with Fetal Bovine Serum (10%), penicillin (100 Units/ml) and streptomycin (100 µg/ml). Single fibers (750-900 for each mouse) were isolated by mechanical dissociation with Pasteur pipettes with decreasing tip diameters and allowed to settle on matrigel-coated (1 mg/ml) glass coverslips accommodated in 35-mm Petri dishes. Cultures were covered with DMEM high glucose enriched with Horse Serum (5%), L-Glutamine (2 mM), penicillin (100 IU/ml) and streptomycin (100 µg/ml). Dishes were maintained in an incubator at 37°C in saturated humidity and in CO<sub>2</sub> (5%)-enriched air. The medium was replaced every 48 h.

### *Assessment of cell contractions*

Myofiber twitching was visualized using an inverted phase-contrast microscope (Axiovert S100, Carl Zeiss, Jena, Germany) equipped with a 40x objective. Images were collected by a digital camera (Sony Alpha 6300) at a sampling frequency of 100 Hz.

### *Immunofluorescence staining*

FDB myofibers seeded on coverslips were fixed with a solution of 4% (w/v) paraformaldehyde in PBS for 20 min at 4 °C, followed by washing three times with PBS (10 min each). Cell permeabilization and the blocking of non-specific protein binding were conducted by incubation with 5% normal goat serum in PBS/0.1% Triton-X100 for 30 min. The staining was performed by incubation with a primary mouse monoclonal anti-Pax7 (1:8 dilution) or a mouse monoclonal anti-myogenin (1:20 dilution), overnight in normal goat serum at 4 °C. After three washes of 10 min with PBS/0.1% Triton-X100, coverslips were incubated with the secondary antibodies for 1.5 h at 4 °C (1:50, Alexa Fluor 594-conjugated Affine Pure goat anti-mouse IgG for anti-Pax7 staining or 1:100, Alexa Fluor 488-conjugated Affine Pure goat anti-mouse IgG for anti-myogenin staining). Nuclei were counterstained by

4',6-diamidino-2-phenylindole (DAPI,1:50). Finally, cells were washed three times with PBS/0.1% Triton-X100 and then mounted onto slides and visualized under a Leica DMLS fluorescence microscope (Leica Microsystems, Wetzlar, Germany). Images were captured with a Leica DC300F camera, coupled to a Leica LM50 acquisition software. Image analysis, measurements of cell morphology and nuclei counting were performed by ImageJ-NIH software. Image sizing, cropping and overlays were obtained with Adobe Photoshop CC (Adobe Systems Incorporated, San Jose, CA). Only dishes containing a comparable number of myofibers (around 100-150) were used for immunofluorescence detection. In the count, we considered the SCs associated to the fibers and those that migrate from them. The aspect ratio (ratio between main and minor axis) of single cells was measured in bright field microscopy in cells expressing myogenin positive nuclei.

Each set of experiments was carried out on three independent cell culture preparations. At least 30 different optical fields were examined in each coverslip corresponding to a minimum myogenic cell number equal to 300.

#### *Electrical stimulation*

ES was carried out on FDB myofibers 20 h after seeding, when SCs were still adhered to the fibers. Coverslips with isolated myofibers were mounted on the stage of an inverted Axiovert microscope (Carl Zeiss, Germany) and stimulated in a 35 mm Petri dish by a custom-built device. Field ES was delivered by the programmable stimulator STG 4002 (Multi Channel Systems, Reutlingen, Germany) connected to a couple of parallel platinum-iridium electrodes (0.2 mm in diameter), placed 1 cm apart and positioned 1-2 mm over the cells. Unless otherwise specified, the maximum voltage strength used was that capable of eliciting twitching in 60% of muscle fibers. EMGstim or bipolar 1 Hz, 1 ms pulses were always delivered in fresh DMEM. Cell cultures were maintained at 37 °C in 5 % CO<sub>2</sub> until they were processed for immunofluorescence.

For Ca<sup>2+</sup> imaging experiments, each single myofiber was electrically stimulated with focal field stimulations using two concentric bipolar platinum electrodes (0.2 mm in diameter) positioned within 3 mm from each other. The inner pole was inserted in a glass pipette surrounded by the outer pole arranged as a wire coil. The concentric electrodes were positioned 1-2 mm over the considered myofiber.

### *Ca<sup>2+</sup> imaging*

Intracellular calcium concentration  $[Ca^{2+}]_i$  was monitored using the fluorescent  $Ca^{2+}$  indicator fura-2 pentacetoxymethyl ester (Fura-2 AM).  $Ca^{2+}$  imaging experiments were performed on skeletal myofibers plated on matrigel-coated coverslips 20 h after seeding. Cell loading was carried out at room temperature, in dark condition, in DMEM plus 5  $\mu$ M Fura-2 AM. After 30 min, the loading solution was removed and the cells washed and bathed in DMEM for 15 min to allow complete de-esterification of the dye. Cells were visualized with a Zeiss Axiovert S100TV inverted microscope (Carl Zeiss, Jena, Germany) using an oil immersion 40x objective.  $Ca^{2+}$  measurements were carried out at room temperature exciting alternately, the cells at 340 and 380 nm, selected by a monochromator device equipped with integrated light source (Polychrome IV, Till Photonics, Gräfelfing, Germany). Fluorescence signals were collected by a CCD camera (SensiCam; PCO Computer Optics, Kelheim, Germany) at a sampling rate of 6.45 Hz. The monochromator and CCD camera were controlled by a software package (TILLvision, Till Photonics) also used for image processing. The ratio of fluorescence images (340/380) and the corresponding temporal plots representing the mean value of the fluorescence signal in regions of interest, were calculated off-line.

$Ca^{2+}$  imaging experiments were carried out on a single cell culture preparation. A total of 8 myofibers were considered for the analysis; area and peaks of at least 40  $Ca^{2+}$  transients for each cell were averaged.

### *Chemicals*

L-Glutamine, penicillin and streptomycin were purchased from Euroclone (Milano, Italy); Fetal Bovine Serum from Gibco (Burlington, ON, Canada); Matrigel from Corning (Tewksbury, MA, USA). Mouse monoclonal anti-Pax7 MAB1675 was from R&D Systems (Minneapolis, MN, USA); mouse monoclonal anti-myogenin (5FD), sc-52903 from Santa Cruz Biotechnology (Dallas, TX, USA). Alexa Fluor 594-conjugated Affine Pure goat anti-mouse IgG and Alexa Fluor 488-conjugated Affine Pure goat anti-mouse IgG were from Jackson ImmunoResearch Laboratories (West Grove, PA, USA). All the other chemicals, unless otherwise stated, were from Sigma (St. Louis, MA, USA).

### *Statistical analysis*

Data were analyzed with GraphPad Prism 4.00 (GraphPad Software, San Diego, CA, USA). A normality *test* was used to determine whether sample data were drawn from a Gaussian distributed population. For parametric data, statistical significance was determined using *t*-



test or one-way ANOVA analysis followed by the Tukey's multiple comparison test. Non-parametric data were analysed by Kruskal-Wallis test followed by Dunn's Multiple Comparison Test. Unless otherwise stated, results were expressed as mean  $\pm$  standard error (SEM). A  $p$  value of  $<0.05$  was considered statistically significant.

## Results

### *EMGstim elicited $[Ca^{2+}]_i$ transients and contractions in FDB myofibers*

$Ca^{2+}$  transients elicited in response to focal ES were measured in single FDB myofibers loaded with the  $Ca^{2+}$  indicator Fura-2AM. The “noisy” protocol *EMGstim*, variable in amplitude, frequency and pulse duration [3], Fig. 1a), was able to elicit intracellular  $Ca^{2+}$  transients and contractions at around 1 Hz as well as the standard protocol of ES at 1 Hz of stereotyped 1 ms pulses (Fig. 1b). To compare the ES-elicited  $Ca^{2+}$  transients, the two protocols were applied to each imaged myofiber at the same maximum strength (mean peak-to-peak amplitude  $3.45 \pm 0.17$  V,  $n = 8$ ). The area under the curve (AUC), relative to the baseline, and the peak amplitude were measured and averaged. Interestingly, *EMGstim* elicited  $Ca^{2+}$  transients with significant larger area and higher amplitude ( $p < 0.05$ ) than 1 Hz stimulation protocol (Fig. 1c). In the same cells, the mean  $Ca^{2+}$  AUC induced by *EMGstim* and 1 Hz ES were  $2438.00 \pm 288.80$  AU ms and  $1434.00 \pm 146.70$  AU ms, respectively. The mean  $Ca^{2+}$  peak amplitude induced by *EMGstim* was  $10.47 \pm 1.44$  Arbitrary Units (AU) whereas it was  $6.58 \pm 0.72$  AU ( $n = 8$  myofibers) at 1 Hz ES.

### *EMGstim did not affect the number of myogenic precursor cells*

Immunolabelling for the marker of quiescent and activated SCs Pax7, performed 2 h after FDB cell plating, revealed the presence of SCs adherent to the single myofibers (1-2 for each of them, Fig. 2). SCs proliferate then as myoblasts [14,15] and migrate far from the myofibers forming new myotubes during the time in culture. In our experimental conditions, migration occurred 36 h after seeding. To determine the effect of *EMGstim* on the number of myogenic precursors, FDB myofibers were stimulated 20 h after seeding and Pax7 positive cells were then detected 36, 48 and 72 h after plating, in stimulated and unstimulated (control) cultures (Fig. 3a). At 36 and 48 h, the percentage of Pax7 positive nuclei per area in *EMGstim*-stimulated cultures did not differ from the controls (36 h:  $63.75 \pm 3.70\%$  vs  $62.09 \pm 4.17\%$ ; 48 h:  $68.47 \pm 4.17\%$  vs  $61.94 \pm 3.13\%$ ; Fig. 3b). In addition, 72 h after seeding, the percentage of Pax7-expressing nuclei per area tended to decrease in both stimulated ( $45.65 \pm 4.20\%$ ) and controls ( $50.08 \pm 5.86\%$ ; Fig. 3b). The latter observation was in line with the increasing number of cells committed to terminal differentiation reported around 72 h in culture [13], when Pax7 is down-regulated and the expression of the transcription factor myogenin increases.

### *EMGstim increased the proportion of myogenin-positive cells*

72 h after FDB myofibre plating, SCs differentiation was determined by counting the percentage of myogenin-positive cells. Compared with the controls, EMGstim delivered for 1 h at 12 V maximum peak-to-peak voltage, increased the proportion of myogenin-positive nuclei ( $44.70 \pm 0.84\%$  vs  $36.76 \pm 0.80\%$ , Fig. 4a, b). Furthermore, in EMGstim the prevalence of myogenic cells with an elongated morphology increased comparing with the controls, as shown by the increase in the aspect ratio ( $5.12 \pm 0.17 \mu\text{m}$  vs  $3.96 \pm 0.16 \mu\text{m}$ ; Fig. 4 c, d).

Regular 1 Hz and 10 Hz, 1 ms ES protocols were delivered to the cultures at 24 V, the output voltage able to induce 60 % of myofiber contractions. 1 Hz, 1 ms ES protocol induced no significant changes in the proportion of myogenin-positive nuclei with respect to control cultures, even if the ES was delivered at a double strength (24 V,  $42.81 \pm 1.41\%$  vs  $39.10 \pm 1.43\%$ , control).

In a different set of experiments 10 Hz, 1 ms regular pulses were elicited at 24 V inducing significant change in the proportion of myogenin-positive nuclei with respect to control cultures ( $55.63 \pm 1.41\%$  vs  $43.88 \pm 1.22\%$ , control,  $p < 0.001$ ). In the same cultures, compared with the controls, EMGstim delivered for 1 h at 12 V maximum peak-to-peak voltage, increased the proportion of myogenin-positive nuclei ( $60.86 \pm 1.46\%$ ,  $p < 0.001$ ), significantly different ( $p < 0.05$ ) from what observed at 10 Hz. The prevalence of myogenic cells with an elongated morphology at 10 Hz was significantly different from the controls (aspect ratio  $6.93 \pm 0.34 \mu\text{m}$  vs  $5.00 \pm 0.21 \mu\text{m}$ ,  $p < 0.001$ ) but not from EMGstim (aspect ratio  $7.83 \pm 0.30$ ,  $p > 0.05$ ).

### *Soluble factors mediated the effect of EMGstim on myogenic differentiation*

In an additional set of experiments, we tested the possibility that the stimulatory effects of EMGstim were due to the release of soluble factor (s) from skeletal myofibers. To this end, immunostaining for myogenin was performed in FDB myofibers in four different experimental conditions: i) controls; ii) 1 h stimulation with EMGstim, started 20 h after cell seeding; iii) unstimulated FDB myofibres cultured in the medium collected from another culture immediately after EMGstim-treatment (conditioned medium, CM); iv) EMGstim delivered as in (ii) but immediately followed by washout with fresh medium (replaced medium, RM). Immunostaining was performed 72 h after plating.

The exposure to CM partially mimicked the effect of EMGstim on cell differentiation. The percentage of myogenin-positive muscle progenitors in cultures exposed to CM ( $46.57 \pm 1.62 \%$ ), was significantly higher than controls ( $39.06 \pm 1.50 \%$ ) but remained lower than in EMGstim cultures ( $57.25 \pm 1.89 \%$ ; Fig. 5a). In addition, CM, similarly to EMGstim, induced elongation of myogenin-positive cells (aspect ratio:  $6.88 \pm 0.35 \mu\text{m}$ ; Fig. 5b). Collectively, the exposure to CM partially mimicked the effect of EMGstim on muscle cell differentiation. Moreover, in the RM cultures, both percentages of myogenin-positive cells ( $40.29 \pm 1.33\%$  vs  $39.06 \pm 1.50\%$ ) and the cell aspect ratio ( $4.18 \pm 0.23$  vs  $4.91 \pm 0.29$ ) were similar to controls (Fig. 5a, b).

Taking into account all these observations, the results suggested a role for soluble factors released from myofibers in determining the stimulatory effects of EMGstim on SCs.

#### *ATP as a potential soluble mediator for EMGstim-induced effect on myogenic differentiation*

Extracellular ATP is released from contracting skeletal muscle fibers during ES [4,6] and promote SC activation [16,17]. This evidence prompted us to study if ATP released from ES/contracting fibers could be implicated in the favoring effect of EMGstim on myogenic differentiation. To test this hypothesis, we firstly analyzed the effect of exogenously-applied ATP on muscle cell differentiation. In particular, 20 h after seeding, increasing concentrations (0.01, 0.1 and  $1 \mu\text{M}$ ) of ATP were added to the FDB cell cultures, incubated for a further 48 h (up to 72 h after seeding). At this time, the percentage of myogenin-positive cells was determined and compared to controls. Results showed that  $0.1 \mu\text{M}$  ATP increased the percentage of myogenin-positive cells ( $58.55 \pm 1.13\%$ ; Fig. 6).

The ATP diphosphohydrolase apyrase is an enzyme catalyzing the hydrolysis of ATP to its unphosphorylated catabolites. To test the involvement of ATP in mediating the EMGstim effect, cells were incubated in with apyrase ( $10 \text{ U/ml}$ ) 30 min before delivering the ES and during EMGstim. The presence of apyrase prevented both the increase in the number and the elongation of myogenic-positive cells induced by either EMGstim treatment and by CM (Fig. 7a, b). To further confirm the role of ATP, the connexin and pannexin 1 inhibitor carbenoxolone (CBX) was used to inhibit ATP release [18,19]. Preincubation in CBX ( $5 \mu\text{M}$ ) prevented the EMGstim effect both on the percentage of myogenin-positive cells (Fig. 7 c) and cell morphology (Fig. 7d).

## Figures

**Fig. 1**  $[Ca^{2+}]$  transients elicited by EMGstim and regular 1 Hz pulses. Representative intracellular  $[Ca^{2+}]$  transients detected in the same cultured myofiber, during EMGstim (**a**) or 1 Hz pulses (**b**) elicited at 3 V (focal ES). On the top of each Figure, the stimulation protocol is shown. The fluorescence 340/380 ratio values are in Arbitrary Units. Scale bars, 5 s. The the mean areas under the curve (AUCs) and the mean  $Ca^{2+}$  peak amplitudes induced by EMGstim and 1 Hz ES, are represented (**c**, n= 8 myofibers). \*  $p<0.05$ , \*\* $p<0.01$ .

**Fig. 2** Distribution of Pax7 positive cells in freshly isolated FDB myofibers. Representative merged image of Pax7 immunolabeled SCs (red) with DAPI nuclear counterstain (blue), 2 h after FDB myofiber seeding. Scale bar, 50  $\mu m$ .

**Fig. 3** EMGstim effect on Pax7 positive cells. **a** Merged images of Pax7 positive cells 36, 48 and 72 h after seeding in control conditions and after 1 h EMGstim. **b** Proportion (%) of Pax7-positive cells at 36, 48 and 72 h in culture in controls and after EMGstim. Scale bars, 100  $\mu m$ .

**Fig. 4** EMGstim effect on myogenin-positive cells. **a** Representative immunofluorescence-labeled myogenin-positive cells (green), DAPI counterstained nuclei (blue) and corresponding merged micrographs, at 72 h of seeding, in control conditions and in EMGstim treated cultures. Scale bars, 100  $\mu m$ . **b** Percentage of myogenin-positive cells in control and EMGstim treated cultures. **c** Representative immunofluorescence-labeled myogenin-positive cells (green) merged with the same images captured in bright field microscopy in control and EMGstim treated cultures. Scale bars, 20  $\mu m$ . **d** The aspect ratio of myogenin positive-cells in control and in EMGstim treated cultures. \*\*\* $p<0.001$  vs controls.

**Fig. 5** Soluble factors mediated the EMGstim effect. **a** Percentage of myogenin-positive cells 72 h post seeding in control conditions, in cultures that underwent EMGstim (EMGstim), in cultures exposed to conditioned medium collected from EMGstim-treated cells (CM) and in cultures in which the medium was replaced immediately after stimulation (RM). See the text

for further details. **b** Quantification of the myogenin-positive cell morphology in EMGstim, CM and RM cultures. \*\*\*  $p < 0.001$ , \*\* $p < 0.01$  vs controls; §§  $p < 0.01$ , §§§  $p < 0.001$  vs EMGstim.

**Fig. 6** Exogenous ATP facilitated myogenic differentiation. Percentage of myogenin-positive cells in cell cultures treated with 0.01, 0.1 and 1  $\mu$ M ATP. \*\* $p < 0.01$  vs controls.

**Fig. 7** Dephosphorylation of ATP and blockage of ATP release prevented the EMGstim effects. **a** Percentage of myogenin-positive cells (expressed as ratio on controls) in cultures stimulated with EMGstim (EMGstim), stimulated with EMGstim in the presence of apyrase (10 U/mL; AP+EMGstim) or cultured in CM from EMGstim treated cultures to which apyrase was added (AP+CM). **b** Morphology of myogenin-positive cells in the experimental conditions as in a. **c** Percentage of myogenin-positive cells (expressed as ratio on controls) in cultures stimulated with EMGstim (EMGstim) or stimulated with EMGstim in the presence of carbenoxolone (5  $\mu$ M; CBX+EMGstim). **d** Morphology of myogenin-positive cells in the experimental conditions as in c. §  $p < 0.05$ , §§  $p < 0.01$ , §§§  $p < 0.001$  vs EMGstim.

## Discussion

ES is recognized as a strategy for providing rehabilitation for individuals suffering from muscle atrophy, especially when physical therapy is limited. Mimicking nerve activity, it not only induces contractions but also drives gene expression patterns, activation of metabolic pathways [20-24] and SC-mediated muscle regeneration [25-27]. Intensive ES, activating a higher number of motor units, could likely be more efficient in inducing muscle potentiation but is often associated with pain [28,29] and muscle fatigue [30,31]. The optimization of the most appropriate parameters for ES thus remains under debate and investigation [25,2].

In the present work, we have studied the effect of the “noisy” stimulation pattern, *EMGstim*, on SC behavior. Compared with a regular stimulation pattern, the “noisy” protocol was already proved to be more effective in inducing firing,  $[Ca^{2+}]_i$  changes and contractions in cultured mouse myotubes [3]. Here, we have shown that the “noisy” stimulation pattern *EMGstim* elicits higher  $Ca^{2+}$  transients in adult mouse FDB myofibers and favors the differentiation of their associated SCs. Moreover, our results indicate that endogenous ATP released by FDB myofibers during the contractile activity could be a crucial factor in enhancing the differentiation of the myogenic precursors.

The “noisy” stimulation pattern *EMGstim* was obtained from human electromyographic recordings and originally designed for the optimal recruitment of neuronal spinal networks [32-33]. In the present work, we provide evidence that in isolated adult FDB myofibers *EMGstim*, delivered at the same maximum strength of regular 1 Hz stimulations, is able to induce significantly higher increases in  $[Ca^{2+}]_i$ . *EMGstim* is characterized by a temporal summation of stochastic pulses with rest intervals of around 1 second. Most likely, the augmented  $Ca^{2+}$  signals induced by *EMGstim* depends on the features of the number of pulses characterizing *EMGstim* signals continuously variable in amplitude, duration and frequency [5]. The intrinsic variability characterizing the “noisy” ES, might facilitate a cumulative membrane depolarization responsible for the  $Ca^{2+}$  release from the sarcoplasmic reticulum during the excitation-contraction coupling, resulting in  $Ca^{2+}$  transients with bigger peaks and area.

Using the isolated FDB myofibers with their resident SCs, we also investigated the effect of *EMGstim* on muscle progenitors since the early phases of activation, proliferation and differentiation. We firstly observed that *EMGstim* did not alter the number of Pax7 positive cells indicating that it did not affect the survival and/or the proliferation of the myogenic precursors. Secondly, we noticed that the “noisy” waveform *EMGstim*, increased the number

of myogenin-positive cells, a phenomenon not observed in response to 1 Hz ES, even if delivered at a double voltage strength. Moreover, the elongated shape of the myogenin-positive cells, with respect to controls, highlights the favoring effect of *EMGstim* on myogenic differentiation. Since the ATP release in FDB fibers was found to depend on ES frequency, 10 Hz regular ES was tested, at which such release was found to be optimal [6]. A slight but significant higher effect of *EMGstim* on increasing the myogenin-positive cells was found, comparing with 10 Hz ES. The fact that *EMGstim* could be elicited at half maximum voltage strengthens the efficacy of such “noisy” ES protocol in inducing cell differentiation.

We also observed that CM, collected from *EMGstim*-treated FDB myofibers, fairly reproduced the effect of *EMGstim* on SCs. In particular, cell exposure to CM mimicked the effect on the elongation of myogenin-positive cells and increased the percentage of myogenin-positive cells, albeit less than *EMGstim*. From these data emerges the contribution of extracellular soluble factors released by the skeletal muscle fibers during ES stimulation. The lower efficacy of CM on SC differentiation could be attributable to the degradation of key signaling molecules mediating the *EMGstim* effect on muscle cell differentiation.

Exercise [34-35] as well as ES *in vivo* [36-37] have been associated with the secretion of ATP from skeletal muscle fibers, that not only circulates and attracts monocytes [38] but can also exert an autocrine/paracrine effect, easily unveiled in a controlled system such as C2C12 cell lines [17] or isolated myofibers [4]. Our observation that exogenous ATP mimicked the *EMGstim* effect on the number and morphology of myogenin-positive cells migrating from adult FDB myofibers suggested that the nucleotide could represent one of the key signaling molecules.

It is not excluded that other nucleotides such as guanosine 5' triphosphate might enhance muscle cell differentiation [39,40].

Extracellular ATP, released after muscle fiber contraction, has been already proposed to act autocrinally as a mediator of the excitation-transcription mechanism in skeletal muscle cells [4,41,6]. Our observations indicate that ATP could also act paracrinally on SCs. The finding that apyrase inhibited the effect of *EMGstim* and CM, rules out the involvement of ATP catabolites such as adenosine. As observed by others [4,42], also in our experimental conditions, ATP release was inhibited by CBX, suggesting that pannexins could mediate a regulated ATP efflux from skeletal myofibers. ATP is supposed to mediate different effects during myogenesis via the activation of purinergic receptors. In mouse SCs [43] and C2C12 myoblasts [17], ATP has been reported to activate ionotropic P2X receptors inhibiting



proliferation and inducing differentiation. ATP application in primary mouse cultures increased the expression of markers of muscle differentiation (myogenin, the cell cycle regulator p21 and myosin heavy chain) via the MAPK signaling cascade [43]. In C2C12 extracellular ATP controlled the levels of myogenin and the number of myotubes [17]. Moreover, other soluble factors could participate to EMG*stim*-induced SCs differentiation [11]. It is worth mentioning that, in FDB myofibers, extracellular ATP increases the expression of the proinflammatory cytokine IL-6 [4,44], which is known to promote muscle cell differentiation [45].

In conclusion, considering the higher efficacy of the EMG*stim* protocol both in inducing contractile activity in myofibers and differentiation of the myogenic precursors, our data strongly encourage the testing of innovative “noisy” stimulation protocols *in vivo* to better counteract muscle atrophy and favor muscle regeneration during rehabilitation training.

### **Ethical standards**

The experiments comply with the current Italian laws (D.L. 4/3/2014 n.26)

### **ACKNOWLEDGMENTS**

We are grateful to Dr. Andrew Constanti (UCL, School of Pharmacy, London, UK) for critically reading the manuscript and to Gianluigi Nocera for doing preliminary experiments.

This work was supported by *Fondazione Beneficentia Stiftung* (Liechtenstein), by MEMORINET project, Interreg V-A Italia-Slovenija 2014-2020 Programme and by Agenzia Spaziale Italiana (ASI; n. 2018-13-U.0).

G.M. is working within *MADE in TRIESTE*, an Area Science Park initiative financed by Commissariato del Governo nella Regione Friuli Venezia Giulia – “Fondo Trieste”.

### **CONFLICT OF INTEREST**

The authors report no conflict of interest.

## References

1. Bergquist AJ, Clair JM, Lagerquist O, Mang CS, Okuma Y, Collins DF (2011) Neuromuscular electrical stimulation: implications of the electrically evoked sensory volley. *Eur J Appl Physiol* 111: 2409–2426.
2. Nikolić N, Görgens SW, Thoresen GH, Aas V, Eckel J, Eckardt K (2017) Electrical pulse stimulation of cultured skeletal muscle cells as a model for in vitro exercise – possibilities and limitations. *Acta Physiol* 220: 310–331.
3. Sciancalepore M, Coslovich T, Lorenzon P, Ziraldo G, Taccola G, (2015) Extracellular stimulation with human “noisy” electromyographic patterns facilitates myotube activity. *J Muscle Res Cell Motil* 36:349–357.
4. Buvinic S, Almarza G, Bustamante M, Casas M, Lopez J, Riquelme M, Saez JC, Huidobro-Toro JP, Jaimovich E (2009) ATP released by electrical stimuli elicits calcium transients and gene expression in skeletal muscle. *J Biol Chem* 284: 34490–34505.
5. Casas M, Figueroa R, Jorquera G, Escobar M, Molgo J, Jaimovich E (2010) IP(3)-dependent, post-tetanic calcium transients induced by electrostimulation of adult skeletal muscle fibers. *J Gen Physiol* 136:455–467.
6. Jorquera G, Altamirano F, Contreras-Ferrat A, Almarza G, Buvinic S, Jacquemond V, et al. (2013). Cav1.1 controls frequency-dependent events regulating adult skeletal muscle plasticity. *J Cell Sci* 126:1189–1198.
7. Mauro A (1961) Satellite cells of skeletal muscle fibers. *J Biophys Biochem Cytol* 9:493–495.
8. Bischoff R (1994) The satellite cell and muscle regeneration. In *Myology* (ed. A.G. Engel and C.Franzini-Armstrong), McGraw Hill, New York, pp. 97–118.
9. Grohovaz F, Lorenzon P, Ruzzier F, Zorec R (1993) Properties of acetylcholine receptors in adult rat skeletal muscle fibers in culture. *J Membr Biol* 136:31–42.
10. Rosenblatt JD, Lunt AI, Parry DJ and Partridge TA (1995) Culturing Satellite Cells from Living Single Muscle Fiber Explants. *In Vitro Cell Dev Biol Anim* 31:773–779.
11. Hawke TJ, Garry DJ (2001) Myogenic satellite cells: physiology to molecular biology. *J Appl Physiol* 91: 534–551.
12. Bekoff A and Betz WJ (1977) Physiological properties of dissociated muscle fibres obtained from innervated and denervated adult rat muscle. *J Physiol* 271(1): 25–40.

13. Yablonka-Reuveni Z & Rivera AJ (1997) Proliferative dynamics and the role of FGF2 during myogenesis of rat satellite cells on isolated fibers. *Basic Appl Myol* 7: 189-202.
14. Yablonka-Reuveni Z & Rivera AJ (1994) Temporal expression of regulatory and structural muscle proteins during myogenesis of satellite cells on isolated adult rat fibers. *Dev Biol* 164: 588-603.
15. Keire P, Shearer A, Shefer G, Yablonka-Reuveni Z (2013) Isolation and culture of skeletal muscle myofibers as a means to analyze satellite cells. *Methods Mol Biol* 946:431-468.
16. Meyer MP, Groschel-Stewart U, Robson T, Burnstock G (1999) Expression of two ATP-gated ion channels, P2X5 and P2X6, in developing chick skeletal muscle. *Dev Dyn* 216:442-449.
17. Araya R, Riquelme MA, Brandan E, Saez JC (2004) The formation of skeletal muscle myotubes requires functional membrane receptors activated by extracellular ATP. *Brain Res Rev* 47: 174-199.
18. Oishi S, Sasano T, Tateishi Y, Tamura N, Isobe M, Furukawa T. (2012) Stretch of atrial myocytes stimulates recruitment of macrophages via ATP released through gap-junction channels. *J Pharmacol Sci* 120:296-304.
19. Dahl G, Qiu F, Wang J (2013) The bizarre pharmacology of the ATP release channel pannexin1. *Neuropharmacology*. 75:583-93.
20. Murgia M, Serrano AL, Calabria E, Pallafacchina G, LomoT, Schiaffino S (2000) Ras is involved in nerve-activity-dependent regulation of muscle genes. *Nat Cell Biol* 2: 142-147.
21. Araya R, Liberona JL, Cardenas JC, Riveros N, Estrada M, Powell JA, Carrasco MA, Jaimovich E. (2003) Dihydropyridine receptors as voltage sensor for a depolarization-evoked, IP3R-mediated, slow calcium signal in skeletal muscle cells. *J Gen Physiol* 121: 3-16.
22. Silveira LR, Pilegaard H, Kusuhara K, Curi R, Hellsten Y (2006) The contraction induced increase in gene expression of peroxisome proliferator-activated receptor (PPAR)-gamma coactivator 1alpha (PGC-1alpha), mitochondrial uncoupling protein 3 (UCP3) and hexokinase II (HKII) in primary rat skeletal muscle cells is dependent on reactive oxygen species. *Biochim Biophys Acta* 1763:969–976.
23. Park H, Bhalla R, Saigal R, Radisic M, Watson N, Langer R, Vunjak-Novakovic G (2008) Effects of electrical stimulation in C2C12 muscle constructs. *J Tissue Eng Regen Med* 2: 279-287.

24. Burch N, Arnold AS, Item F, Summermatter S, Brochmann Santana Santos G, Christie M et al (2010) Electric pulse stimulation of cultured murine muscle cells reproduces gene expression changes of trained mouse muscle. *PLoS ONE* 5:e10970
25. Zhang BT, Yeung SS, Liu Y, Wang HH, Wan YM, Ling SK, Zhang HY, Li YH, Yeung EW (2010) The effects of low frequency electrical stimulation on satellite cell activity in rat skeletal muscle during hindlimb suspension. *BMC Cell Biol* 18,11:87.
26. Guo BS, Cheung KK, Yeung SS, Zhang BT, Yeung E W (2012) Electrical stimulation influences satellite cell proliferation and apoptosis in unloading-induced muscle atrophy in mice. *PLoS One*, 7:e30348.
27. Di Filippo ES, Mancinelli R, Marrone M, Doria C, Verratti V, Toniolo L, Dantas JL, Fulle S, Pietrangelo T (2017) Neuromuscular electrical stimulation improves skeletal muscle regeneration through satellite cell fusion with myofibers in healthy elderly subjects. *J Appl Physiol* 123(3):501-512.
28. Delitto A, Strube MJ, Shulman AD, Minor SD (1992) A study of discomfort with electrical stimulation. *Phys Ther* 72:410–421.
29. Naaman SC, Stein RB, Thomas C (2000) Minimizing discomfort with surface neuromuscular stimulation. *Neurorehabil Neural Repair* 14:223–228.
30. Bickel CS, Slade JM, Warren GL, Dudley GA (2003) Fatigability and variable-frequency train stimulation of human skeletal muscles. *Phys Ther* 83:366–373
31. Gregory CM, Bickel CS (2005) Recruitment patterns in human skeletal muscle during electrical stimulation. *Phys Ther* 85:358–364.
32. Taccola G (2011) The locomotor central pattern generator of the rat spinal cord in vitro is optimally activated by noisy dorsal root waveforms. *J Neurophysiol* 106:872–884.
33. Dose F, Menosso R, Taccola G (2013) Rat locomotor spinal circuits in vitro are activated by electrical stimulation with noisy waveforms sampled from human gait. *Physiol Rep* 1:1–15.
34. Hellsten Y, Maclean D, Rådegran G, Saltin B, Bangsbo J (1998) Adenosine concentrations in the interstitium of resting and contracting human skeletal muscle. *Circulation* 98: 6-8.
35. Li J, King NC, Sinoway LI (2005) Interstitial ATP and norepinephrine concentrations in active muscle. *Circulation* 111: 2748-51.
36. Li J, King NC, Sinoway LI (2003) ATP concentrations and muscle tension increase linearly with muscle contraction. *J Appl Physiol* 95: 577-83.

37. Cunha RA, Sebastiao, AM (1993) Adenosine and adenine nucleotides are independently released from both the nerve terminals and the muscle fibres upon electrical stimulation of the innervated skeletal muscle of the frog. *Pflügers Arch.* 424, 503e510.
38. Burnstock G and Boeynaems JM (2014) Purinergic signalling and immune cells. *Purinergic Signal* 10:529–564.
39. Pietrangelo T, Guarnieri S, Fulle S, Fanò G, Mariggiò MA (2006) Signal transduction events induced by extracellular guanosine 5'-triphosphate in excitable cells. *Purinergic Signal* 2:633-636.
40. Pietrangelo T, Di Filippo ES, Locatelli M, Piacenza F, Farina M, Pavoni E, DiDonato A, Innosa D, Provinciali M, Fulle S (2018) Extracellular Guanosine 5'-triphosphate induces human muscle satellite cells to release exosomes stuffed with guanosine. *Front Pharmacol* 9: 152 doi: 10.3389.
41. Casas M, Buvinic S, Jaimovich E (2014) ATP signaling in skeletal muscle: from fiber plasticity to regulation of metabolism. *Exerc Sport Sci Rev* 42: 110-6.
42. Riquelme MA, Cea LA, Vega JL, Boric MP, Monyer H, Bennett MVL, Frank M, Willecke K, Sáez JC (2013) The ATP required for potentiation of skeletal muscle contraction is released via pannexin hemichannels. *Neuropharmacology* 75:594e603.
43. Ryten M, Dunn PM, Burnstock G (2002) ATP regulates the differentiation of mammalian skeletal muscle by activation of a P2X5 receptor on satellite cells. *J. Cell Biol* 158:345–355.
44. Bustamante M, Fernández-Verdejo, Jaimovich E, Buvinic S (2014) Electrical stimulation induces IL-6 in skeletal muscle through extracellular ATP by activating  $\text{Ca}^{2+}$  signals and an IL-6 autocrine loop. *Am J Physiol Endocrinol Metab* 306:E869-E882.
45. Joannis S, Parise G. (2016) Cytokine Mediated Control of Muscle Stem Cell Function. *Adv Exp Med Biol.* 900:27-44.