



SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI

SISSA Digital Library

The non-equivariant coherent-constructible correspondence and a conjecture of King

*Original*

The non-equivariant coherent-constructible correspondence and a conjecture of King / Scherotzke, S.; Sibilla, N.. - In: SELECTA MATHEMATICA. - ISSN 1022-1824. - 22:1(2016), pp. 389-416. [10.1007/s00029-015-0193-y]

*Availability:*

This version is available at: 20.500.11767/117717 since: 2021-01-21T19:14:32Z

*Publisher:*

*Published*

DOI:10.1007/s00029-015-0193-y

*Terms of use:*

Testo definito dall'ateneo relativo alle clausole di concessione d'uso

*Publisher copyright*

note finali coverpage

(Article begins on next page)

# THE NONEQUIVARIANT COHERENT-CONSTRUCTIBLE CORRESPONDENCE AND A CONJECTURE OF KING

SARAH SCHEROTZKE AND NICOLÒ SIBILLA

**ABSTRACT.** The coherent-constructible (CC) correspondence is a relationship between coherent sheaves on a toric variety  $X$  and constructible sheaves on a real torus  $\mathbb{T}$ . This was discovered by Bondal, and established in the equivariant setting by Fang, Liu, Treumann and Zaslow. In this paper we explore various aspects of the nonequivariant CC correspondence. Also, we use the nonequivariant CC correspondence to prove the existence of tilting complexes in the derived categories of toric orbifolds satisfying certain combinatorial conditions. This has applications to a conjecture of King.

## 1. INTRODUCTION

The coherent-constructible correspondence (*CC correspondence*) is an equivalence of categories that relates coherent sheaves on a toric variety  $X$ , and constructible sheaves on a real torus  $\mathbb{T}$  which is dual to the compact torus acting on  $X$ . The CC correspondence was first conjectured by Bondal in his influential preprint [Bo], and was established in the equivariant setting by Fang, Liu, Treumann, and Zaslow in [FLTZ1, FLTZ2]. In this paper we establish some general results towards the nonequivariant CC correspondence. Further, we achieve a complete proof of the nonequivariant CC correspondence for an interesting class of toric DM stacks: our main application is the construction of tilting complexes in their derived categories. Below we give a more detailed sketch of our results.

**1.1. The non-equivariant coherent constructible correspondence.** Let  $\Sigma = (N, \Sigma, \beta)$  be a stacky fan (see Section 3 for a definition), and let  $M = \text{Hom}(N, \mathbb{Z})$  and  $M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}$ . Denote  $\mathcal{X}_{\Sigma}$  the corresponding toric orbifold. In [FLTZ1, FLTZ2] the authors define a conical Lagrangian  $\tilde{\Lambda}_{\Sigma} \subset T^*M_{\mathbb{R}}$ , and construct an equivalence between the torus equivariant category of perfect complexes  $\mathcal{P}\text{erf}_T(\mathcal{X}_{\Sigma})$ , and the category of compactly supported constructible sheaves on  $M_{\mathbb{R}}$  with singular support in  $\tilde{\Lambda}_{\Sigma}$ ,  $Sh_c(M_{\mathbb{R}}, \tilde{\Lambda}_{\Sigma})$ . The notion of singular support is due to Kashiwara and Schapira [KS], informally it is an invariant that measures the way sections of constructible sheaves propagate. We recall its definition and some of its basic properties in Section 4.

The authors in [FLTZ1, FLTZ2] conjecture that a similar statement holds when replacing the equivariant category with ordinary perfect complexes,  $\mathcal{P}\text{erf}(\mathcal{X}_{\Sigma})$ . For toric varieties partial results in this direction were obtained by Treumann in the preprint [Tr]. In Section 5 we extend Treumann's work to toric orbifolds. More precisely, let  $\mathbb{T}$  be the quotient of  $M_{\mathbb{R}}$  by the integral lattice  $M$ ,  $\mathbb{T} = M_{\mathbb{R}}/M$ . As  $\tilde{\Lambda}$  is invariant under translation by  $M$  we obtain a conical Lagrangian subset  $\Lambda = \tilde{\Lambda}/M$  of  $T^*\mathbb{T}$ .

**Theorem 1.1.** *There is a fully faithful functor  $\kappa : \text{Perf}(\mathcal{X}_\Sigma) \rightarrow \text{Sh}_c(\mathbb{T}, \Lambda_\Sigma)$  which makes the diagram commute (up to natural equivalence)*

$$\begin{array}{ccc} \text{Perf}_T(\mathcal{X}_\Sigma) & \xrightarrow[\cong]{\tilde{\kappa}} & \text{Sh}_c(M_\mathbb{R}, \tilde{\Lambda}_\Sigma) \\ \text{Forg} \downarrow & & \downarrow p! \\ \text{Perf}(\mathcal{X}_\Sigma) & \xrightarrow{\kappa} & \text{Sh}_c(\mathbb{T}, \Lambda_\Sigma), \end{array}$$

where  $p : M_\mathbb{R} \rightarrow M_\mathbb{R}/M_\mathbb{Z}$  is the quotient map, and  $\text{Forg}$  is the forgetful functor.

If  $\mathcal{X}_\Sigma$  is a toric orbifold, we say that the non-equivariant CC correspondence holds for  $\mathcal{X}_\Sigma$  if  $\kappa$  is an equivalence. Following [FLTZ1, FLTZ2] we make the following Conjecture.

**Conjecture 1.2.** *The non-equivariant CC correspondence holds for all toric orbifolds.*

One of our main results is a proof of Conjecture 1.2 for a class of stacky fans  $\Sigma$  called *cragged* (see Definition 6.3), which satisfy some special combinatorial conditions. Bondal and Ruan announced that the non-equivariant CC correspondence holds for weighted projective space [BR]: as weighted projective spaces have cragged fans, this follows from our work.

**Theorem 1.3.** *If  $\Sigma$  is a cragged stacky fan then the functor  $\kappa : \text{Perf}(\mathcal{X}_\Sigma) \rightarrow \text{Sh}_c(\mathbb{T}, \Lambda_\Sigma)$  is an equivalence.*

The conditions defining cragged fans are satisfied in a number of interesting cases: we have already mentioned weighted projective spaces, also all toric Fano surfaces have cragged fans. The proof of Theorem 1.1 depends on a careful study of the category of constructible sheaves  $\text{Sh}_c(\mathbb{T}, \Lambda_\Sigma)$ . When  $\Sigma$  is cragged  $\text{Sh}_c(\mathbb{T}, \Lambda_\Sigma)$  has some important properties, including the fact that it is closed under tensor product of constructible sheaves. This is one of the key ingredients in the proof of Theorem 1.1.

These properties of  $\text{Sh}_c(\mathbb{T}, \Lambda_\Sigma)$  follow from general results about constructible sheaves. In Section 4 we define *cragged Lagrangians*: if  $\Sigma$  is a cragged fan then  $\Lambda_\Sigma$  is a cragged Lagrangian in this sense. Using techniques from Kashiwara and Schapira's microlocal sheaf theory [KS] we prove that if  $M$  is a real analytic manifold and  $\Lambda \subset T^*M$  is cragged then  $\text{Sh}_c(M, \Lambda)$  is always closed under tensor product, and the restriction of the tautological  $t$ -structure of constructible sheaves restrict to a  $t$ -structure on this category.

**1.2. King's Conjecture and tilting.** Section 7 contains an application of the non-equivariant CC correspondence to the algebraic geometry of toric DM stacks. Let us recall the definition of tilting objects in triangulated categories:

**Definition 1.4.** Let  $k$  be a field. Let  $T$  be a  $k$ -linear triangulated category, and let  $\mathcal{E}$  be an object of  $T$ . We say that  $\mathcal{E}$  is a *tilting object* if the following conditions are satisfied:

- $\text{Hom}_T(\mathcal{E}, \mathcal{E}[i]) = 0$  for all  $i \neq 0$ .
- The algebra  $A = \text{Hom}_T(\mathcal{E}, \mathcal{E})$  is finite-dimensional over  $k$ , and has finite global dimension.
- $\mathcal{E}$  generates  $T$  as a triangulated category.

Motivated by the theory of moduli of quiver representations in [Ki] King formulated an influential conjecture:

**Conjecture 1.5** ([Ki] Conjecture 9.3). *Let  $X$  be a smooth complete toric variety. Then  $D^b(\text{Coh}(X))$  admits a tilting object  $\mathcal{E}$  that decomposes as a direct sum of line bundles.*

A lot of work has been done towards establishing King's Conjecture and its extension to toric orbifolds in several classes of examples, see for instance [CM] and [BH] and references therein.<sup>1</sup> Recently however King's Conjecture has been proved to be false [HP, Ef]. It seems reasonable to revise the statement of King's conjecture by dropping the condition that the tilting object is a direct sum of line bundles. At the same time, it is natural to extend its scope to general smooth and proper DM stacks.

**Conjecture 1.6.** *Let  $\mathcal{X}$  be a smooth and proper toric DM stack. Then  $\text{Perf}(\mathcal{X})$  admits a tilting complex.<sup>2</sup>*

In Section 7 we prove Conjecture 1.6 for toric DM stacks with cragged fans.

**Theorem 1.7.** *If  $\Sigma$  is a cragged stacky fan then  $\text{Perf}(\mathcal{X}_\Sigma)$  admits a tilting complex.*

Our technique to produce tilting complexes in  $\text{Perf}(\mathcal{X}_\Sigma)$  is novel and relies in a crucial way on the CC correspondence. The key point is that as  $\Lambda_\Sigma$  is cragged, the tautological  $t$ -structure of  $Sh_c(\mathbb{T})$  restricts to a  $t$ -structure on  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$ , and we have a good control over its heart. From the perspective of toric geometry the condition that  $\Sigma$  is cragged is quite artificial, however at present it cannot be removed: if  $\Sigma$  is not cragged the canonical  $t$ -structure of  $Sh_c(\mathbb{T})$  (essentially) never restricts to a  $t$ -structure on  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$ . We expect that this technical obstruction might be lifted by selecting different  $t$ -structures on constructible sheaves on  $\mathbb{T}$ , such as the ones arising from MacPherson's theory of cellular perverse sheaves [Po, Vy], and investigating their interplay with the microlocal conditions encoded in  $\Lambda_\Sigma$ . We will return to this in future work.

**1.3. The CC correspondence and Homological Mirror Symmetry.** We conclude with a brief discussion of the relations between our work and mirror symmetry.<sup>3</sup> The Fukaya category of a symplectic manifold is an intricate invariant that measures the quantum intersection theory of Lagrangian submanifolds [FOOO, S]. Work of Nadler and Zaslow [NZ, N] shows that, in the case of cotangent bundles, an alternative and much more manageable description of the Fukaya category is available. Let  $M$  be an analytic manifold and  $\Lambda \subset T^*M$  a conical Lagrangian. Restricting to Lagrangian submanifolds of  $T^*M$  which are asymptotic to  $\Lambda$  cuts out a subcategory of the Fukaya category,  $Fuk(T^*M, \Lambda) \subset Fuk(T^*M)$ .

**Theorem 1.8** ([NZ, N]). *There is a quasi-equivalence  $\mu : Fuk(T^*M, \Lambda) \rightarrow Sh_c(M, \Lambda)$ .*

Using Theorem 1.8 we can reformulate Theorems 1.1 and 1.3 as follows:

**Theorem 1.9.** *Let  $\mathcal{X}_\Sigma$  be a toric orbifold.*

- (1) *There is a fully-faithful embedding  $\tau := \mu^{-1} \circ \kappa : \text{Perf}(\mathcal{X}_\Sigma) \rightarrow Fuk(\mathbb{T}, \Lambda_\Sigma)$ .*

<sup>1</sup>A different but closely related result is due to Kawamata [Kaw] who proves that all toric varieties admit exceptional collections composed of coherent sheaves.

<sup>2</sup>A related statement is proposed as Conjecture 1.6 in [Ef].

<sup>3</sup>For an introduction to mirror symmetry with special emphasis on Kontsevich's HMS we recommend the nice overview [Bal].

(2) *Further, if  $\Sigma$  is cragged  $\tau$  is an equivalence.*

Theorem 1.9 (1) extends to toric orbifolds Theorem 2 of [FLTZ3]. In view of Theorem 1.9, we can regard the CC correspondence as a version of Kontsevich's Homological Mirror Symmetry conjecture (HMS) for toric orbifolds. This perspective is explained in great detail in [FLTZ3]. The authors show that the functor  $\tau$  can be interpreted as implementing T-duality which, according to the Strominger-Yau-Zaslow conjecture [SYZ], is the geometric mechanism responsible for mirror symmetry. We remark that the ordinary formulation of HMS for toric orbifolds differs from the one expounded in [FLTZ3], although it is conjecturally equivalent to it. We explain this next.

Standard HMS in the toric setting [Ko] predicts that there is an equivalence between  $\mathcal{P}erf(\mathcal{X}_\Sigma)$  and the Fukaya-Seidel category of the mirror *Landau-Ginzburg (LG) model*. The LG model is given by a holomorphic map  $W_\Sigma : (\mathbb{C}^*)^n \rightarrow \mathbb{C}$  encoding the combinatorics of the fan  $\Sigma$ ; the Fukaya-Seidel category  $\overrightarrow{Fuk}((\mathbb{C}^*)^n, W_\Sigma)$  is a symplectic invariant of the pair  $((\mathbb{C}^*)^n, W_\Sigma)$ . In the last fifteen years this formulation of HMS for toric orbifolds has been intensely studied, see [AKO1], [AKO2] and references therein. We also note the papers [A1, A2], by Abouzaid, which take an approach to HMS for toric varieties that is closer in spirit to the CC correspondence: a detailed discussion of the relationship between Abouzaid's work and the CC correspondence can be found in Appendix C2 of [FLTZ3].

Although seemingly quite different, the ordinary formulation of HMS for toric orbifolds is conjecturally related to the CC correspondence. In fact, applications to HMS provided the main motivation for Bondal's original work on the CC correspondence. According to [Bo, BR] the symplectic geometry of the LG model  $((\mathbb{C}^*)^n, W_\Sigma)$  can be read off from the sheaf theory of the compact torus  $\mathbb{T} \subset (\mathbb{C}^*)^n$ , which can be identified with  $M_{\mathbb{Z}} \otimes_{\mathbb{Z}} U(1)$ . In particular, when  $\Sigma$  is the stacky fan of a weighted projective space Bondal and Ruan conjecture that there is an equivalence:

$$Sh_c(\mathbb{T}, \Lambda_\Sigma) \cong \overrightarrow{Fuk}((\mathbb{C}^*)^2, W_\Sigma).^4$$

We can extract from our work a proof of Bondal and Ruan's claim in the simplest non-trivial class of examples: weighted projective planes. Indeed, Theorem 1.3 together with the proof of ordinary HMS for weighted projective planes given by Auroux, Katzarkov and Orlov in [AKO1] implies the following:

**Proposition 1.10.** *Let  $\Sigma$  be the stacky fan of a weighted projective plane. Then there is an equivalence:  $Sh_c(\mathbb{T}, \Lambda_\Sigma) \cong \overrightarrow{Fuk}((\mathbb{C}^*)^2, W_\Sigma)$ .*

**Acknowledgements:** We thank David Treumann and Eric Zaslow for their interest in this project. We thank particularly David Treumann for many useful discussions. Claus Ringel kindly answered some of our questions. S.S. thanks the Mathematical Institute for funding and providing a stimulating environment. N.S. thanks the Max Planck Institute for Mathematics for excellent working conditions.

---

<sup>4</sup>Bondal's original statement involved considering first a stratification  $\mathcal{S}_\Sigma$  on  $\mathbb{T}$  capturing the combinatorics of  $\Sigma$ , and then the subcategory of sheaves constructible with respect to  $\mathcal{S}_\Sigma$ ,  $Sh_c(\mathbb{T}, \mathcal{S}_\Sigma)$ . However the categories  $Sh_c(\mathbb{T}, \mathcal{S}_\Sigma)$  and  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$  coincide: this follows from a mild generalization to non-Whitney stratifications of Remark 4.5 below.

## 2. CONVENTIONS AND BACKGROUND

Throughout the paper we work over the field of complex numbers  $\mathbb{C}$ .

**2.1. Categories.** By *dg category* we shall always mean a  $\mathbb{C}$ -linear differential graded category, see [Ke] and [Dr] for definitions and basic properties. If  $C$  is a dg category and  $x, y$  are objects in  $C$ , we denote  $\text{hom}_C(x, y)$  the complex of morphisms between  $x$  and  $y$ . We denote  $D(C)$  the *homotopy category* of the dg category  $C$ :  $D(C)$  has the same objects then  $C$ , while the set of morphisms  $\text{Hom}_{D(C)}(x, y)$  is given by the zero-th cohomology of the complex  $\text{hom}_C(x, y)$ . It will be often useful to work with *triangulated* dg categories, which are defined for instance in Section 7 of [To]. If  $E$  is the set of objects in a triangulated dg category  $C$  we denote  $\langle E \rangle \subset C$  the smallest triangulated dg subcategory of  $C$  containing  $E$ . The homotopy category  $D(C)$  of a triangulated dg category  $C$  is triangulated in the classical sense. We say that a triangulated dg category  $C$  has a t-structure if its homotopy category  $D(C)$  has one. The reader can consult [Lu] Section 1.2.1 for basic facts on t-structures in the context of stable  $(\infty, 1)$ -categories: all the statements there can be adapted to triangulated dg categories in a straightforward way. We sometimes consider diagrams of functors between dg categories: we say that such a diagram is *commutative* if it is commutative up to canonical natural equivalence.

For future reference we recall the notion of *Postnikov systems* in triangulated dg categories, see Chapter 4 of [GM]. Postnikov systems give a convenient way of talking about resolutions in triangulated categories.

**Definition 2.1.** Let  $C$  be a triangulated dg category, and let

$$X^\bullet = X^c \xrightarrow{d^c} X^{c+1} \xrightarrow{d^{c+1}} \dots \xrightarrow{d^0} X^0$$

be a sequence of objects in  $C$  with degree zero maps between them, such that  $d^{i+1}d^i = 0$  in the homotopy category.

- A (*left*) *Postnikov system* attached to  $X^\bullet$  is a diagram in  $C$  of the form

$$\begin{array}{ccccccc} X^c & \xrightarrow{d^c} & X^{c+1} & \xrightarrow{d^{c+1}} & X^{c+2} & \xrightarrow{d^{c+2}} & \dots \xrightarrow{d^{-1}} X^0 \\ & \searrow Id & \uparrow f^c = d^c & \searrow g^{c+1} & \uparrow f^{c+1} & \searrow g^{c+2} & \uparrow f^{c+2} \\ & & Y^c = X^c & \xleftarrow{+1} & Y^{c+1} & \xleftarrow{+1} & Y^{c+2} \xleftarrow{+1} \dots \xleftarrow{+1} Y^0, \\ & & & & \uparrow f^{c+1} & \uparrow f^{c+2} & \uparrow f^{-1} \\ & & & & & & \uparrow g^0 \end{array}$$

which satisfies the following conditions:

- (1)  $f^i$  and  $g^i$  have degree zero, and  $f^i g^i = d^i$  in the homotopy category.
  - (2)  $Y^i \xrightarrow{f^i} X^{i+1} \xrightarrow{g^{i+1}} Y^{i+1}$  is a cofiber sequence.
- We call an object  $Y$  of  $C$  a *left convolution* of  $X^\bullet$  if there is a Postnikov system attached to  $X^\bullet$  with the property that  $Y = Y^0$ .

**Remark 2.2.** Postnikov systems are preserved by dg functors of triangulated dg categories. Note also that if  $Y \in C$  is a left convolution of  $X^\bullet$  then  $Y$  belongs to the subcategory  $\langle X^c, \dots, X^0 \rangle \subset C$ .

**2.2. Constructible sheaves.** The classical reference for microlocal sheaf theory is [KS]. Although the theory of [KS] is cast in the language of ordinary triangulated categories, it is straightforward to adapt it to the setting of dg categories. This issue is discussed at greater length in [NZ, N] to which we refer the reader for further details. If  $M$  is a topological space we denote  $Sh(M)$  the dg category of bounded complexes of sheaves of  $\mathbb{C}$ -vector spaces on  $M$  localized at quasi-isomorphisms:  $Sh(M)$  is a dg enhancement of the usual derived category of sheaves over  $M$ , which is equivalent to the homotopy category  $D(Sh(M))$ . For the theory of localization in the dg setting we refer the reader to [Dr].

Assume that  $M$  is a real analytic manifold. A sheaf  $\mathcal{F}$  of  $\mathbb{C}$ -vector spaces is called *quasi-constructible* if there is a Whitney stratification of  $M$  such that the restriction of  $\mathcal{F}$  to each stratum is locally constant.  $\mathcal{F}$  is called *constructible* if additionally its stalks  $\mathcal{F}_x$  are finite dimensional vector spaces for all  $x \in M$ . We denote  $Sh_c(M)$  (resp.  $Sh_{qc}(M)$ ) the triangulated dg category of complexes with constructible (resp. quasi-constructible) cohomology. We often refer to objects in  $Sh_c(M)$  simply as constructible sheaves. An important invariant of an object  $\mathcal{F} \in Sh_c(M)$  is its singular support, see [KS] chapter V. We give a detailed review of this concept in Section 4. If  $\Lambda \subset T^*M$  is a  $\mathbb{R}_{>0}$ -invariant Lagrangian, the triangulated dg category of constructible sheaves whose singular support is contained in  $\Lambda$  is denoted  $Sh_c(M, \Lambda) \subset Sh_c(M)$ .

The theory of Grothendieck's (derived) six operations  $f_*, f^*, f!, f^!, \otimes, \mathcal{H}om$  can be lifted to the dg category of constructible sheaves  $Sh_c(M)$ , see [N] Section 2.2. We denote  $\Gamma(-)$  the functor of global sections. If  $\mathcal{F} \in Sh_c(M)$  and  $Z \subset M$  is a closed subset we denote  $\Gamma_Z(\mathcal{F})$  the derived functor of the subsheaf of sections supported on  $Z$ . Throughout the paper, it will always be understood that all the functors between triangulated dg categories are dg derived: thus, for instance, we denote the derived push-forward simply  $f_*$ , and not  $Rf_*$ , and similarly for  $\Gamma(-)$  and  $\Gamma_Z(-)$ .

**2.3. Deligne-Mumford stacks.** An introduction to Deligne-Mumford (DM) stacks can be found in the appendix of [Vi]. We refer to [Vi] also for a definition of (quasi)-coherent sheaves and vector bundles on DM stacks. If  $\mathcal{X}$  is a DM stack we denote  $\mathcal{QCoh}(\mathcal{X})$  the triangulated dg category of bounded complexes of quasi-coherent sheaves on  $\mathcal{X}$ :  $\mathcal{QCoh}(\mathcal{X})$  is obtained from the dg category of bounded complexes of quasi-coherent sheaves on  $\mathcal{X}$  by localizing at quasi-isomorphisms. A perfect complex is an object of  $\mathcal{QCoh}(\mathcal{X})$  which is locally quasi-isomorphic to a complex of vector bundles on  $\mathcal{X}$ . The triangulated dg category of perfect complexes is denoted  $\mathcal{Perf}(\mathcal{X}) \subset \mathcal{QCoh}(\mathcal{X})$ . Let  $G$  be an algebraic group acting on  $\mathcal{X}$  in the sense of [Ro]. The  $G$ -equivariant categories of quasi-coherent sheaves and perfect complexes on  $\mathcal{X}$  will be denoted respectively  $\mathcal{QCoh}_G(\mathcal{X})$  and  $\mathcal{Perf}_G(\mathcal{X})$ .

### 3. TORIC ORBIFOLDS AND STACKY FANS

In this section we review the definition of *toric DM stacks* and *stacky fans* due to Borisov, Chen and Smith [BCS], see also [FMN] and [Iw]. We also recall a few facts in toric geometry which will be important later. We follow quite closely [BCS] except we work in somewhat lesser generality. Indeed, in order to simplify the exposition we restrict to *toric orbifolds*, i.e. toric DM stacks with trivial generic stabilizers.

**Definition 3.1.** A (*strict*) *stacky fan*  $\Sigma$  is given by a tuple  $(N, \Sigma, \beta)$ , where:

- $N \cong \mathbb{Z}^n$  is a finitely generated free abelian group.
- $\Sigma$  is a finite rational simplicial fan in  $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$ , such that its set of one dimensional cones  $\Sigma(1) = \{\rho_1, \dots, \rho_r\}$  spans  $N_{\mathbb{R}}$ .
- $\beta : \mathbb{Z}^r = \bigoplus_{i=1}^r \mathbb{Z} \cdot e_i \rightarrow N$  is a homomorphism such that for all  $i \in \{1 \dots r\}$   $b_i := \beta(e_i)$  belongs to  $\rho_i$ .

To any stacky fan we can attach a toric orbifold  $\mathcal{X}_{\Sigma}$  in the following way. Let  $M = \text{Hom}(N, \mathbb{Z})$  and let  $\beta^* : M \rightarrow M' := \text{Hom}(\mathbb{Z}^r, \mathbb{Z})$  be the dual of  $\beta$ . It follows from our assumptions that  $\beta^*$  is injective. Denote  $\text{Gale}(N)$  the cokernel of  $\beta^*$ , and set

$$G_{\Sigma} := \text{Hom}(\text{Gale}(N), \mathbb{C}^*).$$

The morphism  $\beta^*$  induces an embedding  $G_{\Sigma} \hookrightarrow \text{Hom}(M', \mathbb{C}^*) \cong (\mathbb{C}^*)^r$ . For all  $\sigma \in \Sigma$  denote  $Z_{\sigma} \subset \text{Hom}(M', \mathbb{C}) \cong \mathbb{C}^r$  the closed subset defined as follows

$$Z_{\sigma} := \{(z_1, \dots, z_r) \in \mathbb{C}^r \mid \prod_{i \text{ s.t. } \rho_i \not\subseteq \sigma} z_i = 0\}.$$

Let  $U_{\Sigma}$  be equal to the open subset  $\mathbb{C}^r - \bigcup_{\sigma \in \Sigma} Z_{\sigma}$ . Note that  $U_{\Sigma}$  is invariant under the action of  $G_{\Sigma}$ .

**Definition 3.2.** The toric orbifold  $\mathcal{X}_{\Sigma}$  associated to the stacky fan  $\Sigma$  is the quotient stack  $[U_{\Sigma}/G_{\Sigma}]$ .

The fact that  $\mathcal{X}_{\Sigma}$  is a DM stack is proved in Proposition 3.2 of [BCS]. Also, it follows from Definition 3.1 that  $\mathcal{X}_{\Sigma}$  is an orbifold, i.e. it has trivial generic stabilizer.

**Remark 3.3.** The coarse moduli space of  $\mathcal{X}_{\Sigma}$  is the toric variety associated to the underlying ordinary fan  $\Sigma$ ,  $X_{\Sigma}$ , see [BCS] Proposition 3.7. If  $\Sigma$  is smooth and all the vectors  $b_i = \beta(e_i)$  are primitive, then  $\mathcal{X}_{\Sigma}$  is a variety and is isomorphic to  $X_{\Sigma}$ . This is not the case if the  $b_i$ -s are primitive vectors but  $\Sigma$  is not smooth: indeed  $\mathcal{X}_{\Sigma}$  is always smooth.

**Remark 3.4.** As explained in [FMN] toric DM stacks admit an intrinsic definition as compactifications of DM tori (see [FMN] Definition 3.2) carrying a DM torus action, much in the same way as ordinary toric varieties. The case of toric orbifolds is especially simple. Indeed, if the generic stabilizer is trivial the DM torus of [FMN] coincides with the ordinary algebraic torus  $T = N \otimes_{\mathbb{Z}} \mathbb{C}^*$  and there is an open embedding  $T \hookrightarrow \mathcal{X}_{\Sigma}$ .

Next we state some results which extend to toric orbifolds some familiar properties of toric varieties. In the sequel when referring to objects in  $\text{Perf}(\mathcal{X}_{\Sigma})$  admitting an equivariant structure, this will always be with respect to the natural action by the open dense torus  $T \hookrightarrow \mathcal{X}_{\Sigma}$ .

**Proposition 3.5** ([BH]). *Let  $\mathcal{X}$  be a toric orbifold, then any coherent sheaf  $\mathcal{F}$  on  $\mathcal{X}$  admits a finite resolution  $\mathcal{P}^{\bullet} = (\dots \rightarrow \mathcal{P}^1 \rightarrow \mathcal{P}^0) \rightarrow \mathcal{F}$  such that, for all  $m$ ,  $\mathcal{P}^m$  is isomorphic to a direct sum of line bundles:  $\mathcal{P}^m \cong \bigoplus_{i=1}^{i=r} \mathcal{L}_i$ .*

*Proof.* This is proved in [BH], see Corollary 4.8. □

**Remark 3.6.** It will be useful to think of the resolution  $\mathcal{P}^{\bullet}$  using the language of Postnikov systems. Indeed, we can regard  $\mathcal{P}^i$  as objects in  $\text{Perf}(\mathcal{X})$  placed in degree zero. Then Proposition 3.5 can be restated as saying that  $\mathcal{F}$  is a left convolution of  $\mathcal{P}^{\bullet}$ .

**Proposition 3.7.** *Any line bundle  $\mathcal{L}$  on a toric orbifold  $\mathcal{X}$  admits a  $T$ -equivariant structure.*

*Proof.* Line bundles on  $\mathcal{X}_\Sigma = [U_\Sigma/G_\Sigma]$  are given by  $G_\Sigma$ -equivariant line bundles on  $U_\Sigma$ . The statement then follows because  $U_\Sigma$  is an ordinary toric variety and line bundles on ordinary toric varieties always admit an equivariant structure.  $\square$

**Corollary 3.8.** • *Let  $L$  be the set of line bundles on  $\mathcal{X}$ , then  $\langle L \rangle = \text{Perf}(\mathcal{X})$ .*  
 • *Let  $E_T$  be the set of objects in  $\text{Perf}(\mathcal{X})$  admitting an equivariant structure, then  $\langle E_T \rangle = \text{Perf}(\mathcal{X})$ .*

#### 4. CONSTRUCTIBLE SHEAVES AND CRAGGED LAGRANGIANS

In this section we prove a few general results about constructible sheaves which will play a key role in our argument. We start by giving some more explanations on the notion of singular support. For the rest of this section we assume that  $M$  is a real analytic manifold. Let  $\mathcal{F}$  be an object in  $Sh_c(M)$ :

**Definition 4.1** ([KS] Definition 5.1.2). The singular support  $SS(\mathcal{F})$  of  $\mathcal{F}$  is the subset of  $T^*M$  defined by the following condition: if  $p$  is a point in  $T^*M$ ,  $p \notin SS(\mathcal{F})$  if and only if there exists an open neighborhood  $U$  of  $p$  having the property that for all  $x_0 \in M$ , and for all 1-differentiable function  $\psi$  defined in a neighborhood of  $x_0$  and such that  $\psi(x_0) = 0$  and  $d\psi(x_0) \in U$ , we have an isomorphism  $(R\Gamma_{\{x \in M \text{ s. t. } \psi(x) \geq 0\}}(\mathcal{F}))_{x_0} \cong 0$ .

**Definition 4.2.** Let  $M$  be a manifold. A *stratification*  $\mathcal{S}$  of  $M$  is the datum of a preorder  $I$ , and a locally finite collection of locally closed subsets of  $M$  indexed by  $I$ ,  $\{Z_i\}_{i \in I}$ , with the following properties

- for all  $i, j$ , if  $i \neq j$  then  $Z_i \cap Z_j = \emptyset$ ,
- $Z_i \cap \overline{Z_j} \neq \emptyset$  if and only if  $Z_i$  is contained in  $\overline{Z_j}$ , and this holds if and only if  $i \leq j$ .
- The subsets  $\{Z_i\}_{i \in I}$  are the strata of a Whitney stratification of  $M$ .

**Definition 4.3.** If  $\mathcal{S}$  is a stratification, we denote  $Sh_c(M, \mathcal{S}) \subset Sh_c(M)$  the triangulated dg category of complexes that have cohomology sheaves which are constructible with respect to  $\{Z_i\}_{i \in I}$ .

In Theorem 4.4 we gather a few important properties of the singular support, referring to [KS] Chapter V and VI for proofs. We say that a subset  $A \subset T^*M$  is *conical* if it is invariant under fiber-wise dilation by positive real numbers, that is if for all  $z \in M$  and  $\lambda \in \mathbb{R}_{\geq 0}$   $\lambda(A \cap T_z^*M) \subset A$ .

**Theorem 4.4.** *Let  $\mathcal{F} \in Sh_c(M)$ , then*

- (1)  $SS(\mathcal{F}) \subset T^*M$  is a conical Lagrangian subset.
- (2) The support of  $\mathcal{F}$  is equal to  $SS(\mathcal{F}) \cap T_M^*M$ .
- (3) If  $\mathcal{S} = \{Z_i\}_{i \in I}$  is a stratification of  $M$  and  $\mathcal{F}$  is constructible with respect to  $\mathcal{S}$ , then  $SS(\mathcal{F}) \subset \bigcup_{i \in I} T_{Z_i}^*M$ .
- (4) If  $\mathcal{F}_1 \rightarrow \mathcal{F}_2 \rightarrow \mathcal{F}_3$  is a cofiber sequence in  $Sh_c(M)$ ,  $SS(\mathcal{F}_2) \subset SS(\mathcal{F}_1) \cup SS(\mathcal{F}_3)$ .

As we mentioned in Section 2.2 if  $\Lambda \subset T^*M$  is a conical Lagrangian set, we can consider the subcategory  $Sh_c(M, \Lambda)$  of constructible sheaves  $\mathcal{F}$  such that  $SS(\mathcal{F}) \subset \Lambda$ : Theorem 4.4 (4) shows that  $Sh_c(M, \Lambda)$  is triangulated.

**Remark 4.5.** Let  $\mathcal{S} = \{Z_i\}_{i \in I}$  be a stratification, and let  $\Lambda_{\mathcal{S}} = \bigcup_{i \in I} T_{Z_i}^* M$ . Then by Theorem 4.4 (3),  $Sh_c(M, \mathcal{S}) \subset Sh_c(M, \Lambda_{\mathcal{S}})$ . It is easy to see that this inclusion is an equivalence.

The category  $Sh_c(M)$  has two additional structures which will be of interest for us: it is a symmetric monoidal category under tensor product of sheaves; also, it comes equipped with a tautological t-structure whose heart is given by sheaves placed in degree 0. However for a general conical Lagrangian  $\Lambda$  the tensor product does not restrict to the subcategory  $Sh_c(M, \Lambda) \subset Sh_c(M)$ , and similarly for the t-structure. In the rest of this Section we introduce a class of Lagrangians, called *cragged*, having the property that both the t-structure and the tensor product restrict to  $Sh_c(M, \Lambda)$ .

**Remark 4.6.** Let  $\mathcal{S} = \{Z_i\}_{i \in I}$  be a stratification, and let  $\Lambda_{\mathcal{S}} = \bigcup_{i \in I} T_{Z_i}^* M$ . It is easy to see that  $Sh_c(M, \mathcal{S})$  is closed under tensor product, and that the tautological t-structure of  $Sh_c(M)$  restricts as well. Lagrangians of the form  $\Lambda_{\mathcal{S}}$  are encompassed by our notion of cragged Lagrangian.

**Definition 4.7.** (1) Let  $U$  be an open subset of a vector space  $E$ . We say that a closed subset  $A \subset T^*U$  is *globally cragged* if there is a stratification  $\mathcal{S} = \{Z_i\}_{i \in I}$  of  $U$ , and a collection of closed convex cones  $\{V_i\}_{i \in I}$  in  $E^*$  such that:  $A = \bigcup_{i \in I} Z_i \times V_i$ .  
 (2) More generally, suppose that  $M$  is a manifold and that  $A \subset T^*M$  is a closed subset. We say that  $A$  is *cragged* if there is an atlas  $(U_{\alpha}, \phi_{\alpha})_{\alpha \in J}$  of  $M$  such that, for all  $\alpha \in J$ ,  $(\phi_{\alpha}, d\phi_{\alpha})$  is a trivialization of the cotangent bundle that maps  $A_{\alpha} = A \cap T^*U_{\alpha}$  onto a globally cragged set.

**Lemma 4.8.** *Under the assumptions of Definition 4.7 (1), we also have that if  $i \leq j$  then  $V_i \supset V_j$ .*

*Proof.* If  $z$  lies in  $Z_i$ , there is a sequence  $(x_n)_{n \in \mathbb{N}}$  in  $Z_j$  having the property that  $x_n$  converges to  $z$ . Let  $\xi$  be in  $V_j$ , and consider the sequence  $(x_n, \xi)_{n \in \mathbb{N}} \rightarrow (z, \xi)$ . As  $A$  is closed and  $(x_n, \xi)_n$  belongs to  $A$  for all  $n$ , the limit  $(z, \xi)$  belongs to  $A$  as well: thus  $\xi$  lies in  $V_i$ .  $\square$

Denote  $+: T^*M \times_M T^*M \rightarrow T^*M$  the map of fiberwise addition. If  $A$  and  $B$  are subsets of  $T^*M$ , write  $A + B$  for the image of  $A \times_M B$  under  $+$ . Following [KS] denote  $A \hat{+} B$  the subset of  $T^*M$  given in coordinates as follows:  $(z, \zeta) \in T^*M$  lies in  $A \hat{+} B$  if and only if there are sequences  $(x_n, \xi_n)_{n \in \mathbb{N}}$  in  $A$  and  $(y_n, \eta_n)_{n \in \mathbb{N}}$  in  $B$  such that

- (1)  $x_n \xrightarrow{n} z$ ,  $y_n \xrightarrow{n} z$ , and  $\xi_n + \eta_n \xrightarrow{n} \zeta$ , and
- (2)  $|x_n - y_n| |\xi_n| \xrightarrow{n} 0$ .

**Definition 4.9.** A subset  $A \subset T^*M$  is called *pre-additive* (resp. *additive*) if it is conical, and  $A + A = A$  (resp.  $A \hat{+} A = A$ ).

Note that an additive set is also pre-additive.

**Lemma 4.10.** *Let  $M$  be a manifold. If  $A \subset T^*M$  is a cragged set, then  $A$  is additive.*

*Proof.* Since additivity is a local property, we can assume that  $M = U$  where  $U$  is an open subset in a vector space  $E$ , and that  $A$  is globally cragged. We use the notation of Definition 4.7 (1). If  $p$  lies in  $M$  we denote  $i_p$  the element of  $I$  such that  $p$  belongs to  $Z_{i_p}$ .

Fix  $z \in M$ . We need to show that we have an inclusion  $(A \hat{+} A)_z \subset A_z$ . Note that for all points  $p$  in a sufficiently small neighborhood of  $z$ , we have that  $i_z \leq i_p$ . Let  $(z, \xi)$  be in  $(A \hat{+} A)_z$ . Then there are sequences  $(x_n, \xi_n)_{n \in \mathbb{N}}$  and  $(y_n, \eta_n)_{n \in \mathbb{N}}$  in  $A$  such that  $x_n \xrightarrow{n} z$ ,  $y_n \xrightarrow{n} z$ , and  $\xi_n + \eta_n \xrightarrow{n} \xi$ . We can assume that  $i_z \leq i_{x_n}$  and  $i_z \leq i_{y_n}$  for all  $n$ . Since  $A$  is cragged, this implies that for all  $n$  both  $(z, \xi_n)$  and  $(z, \eta_n)$  belong to  $A_z$ . Also  $A_z$  is a closed convex cone: since the sequence  $(z, \xi_n + \eta_n)_{n \in \mathbb{N}}$  lies in  $A_z$ , the limit  $(z, \xi)$  to which this sequence converges belongs to  $A_z$  as well. That is,  $(z, \xi) \in A_z$  as we wanted to prove.  $\square$

Suppose that  $M, A \subset T^*M$  and  $\{U_\alpha, \phi_\alpha\}_{\alpha \in J}$  satisfy the conditions described in Definition 4.7 (2) in the following weaker form. The chart  $(\phi_\alpha, d\phi_\alpha)$  maps  $A_\alpha = T^*U_\alpha \cap A$  onto a set of the form  $\bigcup_{i \in I} P_i \times V_i$ , where  $\{V_i\}_{i \in I}$  are convex cones and  $\bigcup_{i \in I} P_i = \phi_\alpha(U_\alpha)$ . We do not assume that  $\{P_i\}_{i \in I}$  is a stratification (in particular the  $P_i$ -s might be non pairwise disjoint): we assume however that there exists a stratification that refines  $\{P_i\}_{i \in I}$ . If  $A$  has these properties we say that  $A$  satisfies condition (2'). The following holds:

**Lemma 4.11.** *Let  $A \subset T^*M$  be a closed subset. The following are equivalent:*

- *$A$  is cragged.*
- *$A$  is pre-additive and satisfies condition (2').*

*Proof.* Lemma 4.10 shows that cragged implies pre-additive, so we only have to prove the other implication. Let  $\alpha \in J$ : using  $\phi_\alpha$  we identify  $T^*U_\alpha$  and  $U_\alpha \times E^*$ , and we set  $U := U_\alpha$  and  $A := A \cap T^*U$ . By pre-additivity the fibers of  $A$  over the points of  $U$  are cones: also, they can be written as finite unions of elements in  $\{V_i\}_{i \in I}$ . By shrinking  $U$  we can assume that, regarded as subsets of  $E^*$ , there are only finitely many fibers of  $A$  over  $U$ . We denote them  $W_1, \dots, W_n$ . By further shrinking  $U$  we can also assume that the intersection of  $\{P_i\}_{i \in I}$  with  $U$  is a finite collection of closed subsets, which we denote  $P_1, \dots, P_m$ .

Now consider the partition  $Q_1, \dots, Q_n$  of  $U$  defined by the following rule:  $p \in Q_j$  if  $A_p = W_j$ . Note that the  $Q_j$ -s are obtained from the closed subsets  $P_1, \dots, P_m$  by taking unions, intersections, and complements: since by assumption  $\{P_i\}_{i \in I}$  can be refined to a stratification, the same is true of  $Q_1, \dots, Q_n$ . Let  $\{Z_i\}_{i \in I'}$  be a stratification refining the partition  $\{Q_1 \dots Q_n\}$ . By construction, we can write  $A = \bigcup_{i \in I'} Z_i \times W_i$ , where  $W_i \in \{W_1, \dots, W_n\}$  are convex cones. That is,  $A$  is cragged, as we wanted to show.  $\square$

**Lemma 4.12.** *Let  $\Lambda \subset T^*M$  be a conical Lagrangian subset and assume further that  $\Lambda$  is additive. Then  $Sh_c(M, \Lambda)$  is closed under tensor product.*

*Proof.* This follows from results contained in Chapter 5 of [KS] (see also Theorem 1.1 of [GS]): in particular, we have an inclusion  $SS(F \otimes G) \subset SS(F) \hat{+} SS(G)$ . If  $\Lambda$  is additive, then  $SS(F) \hat{+} SS(G) \subset \Lambda \hat{+} \Lambda = \Lambda$ .  $\square$

**Corollary 4.13.** *If  $\Lambda \subset T^*M$  is a cragged conical Lagrangian, then  $Sh_c(M, \Lambda)$  is closed under tensor product.*

Next, we prove that if  $\Lambda$  is a cragged conical Lagrangian the tautological  $t$ -structure on  $Sh(M)$  restricts to a  $t$ -structure on  $Sh(M, \Lambda)$ .

**Lemma 4.14.** *Let  $i : U \hookrightarrow M$  be an open subset, and let  $\mathcal{F}$  and  $\mathcal{G}$  be two objects in  $Sh(M)$ . Then:*

- $SS(i^*\mathcal{F}) = SS(\mathcal{F}) \cap T^*U$ .
- $H^j(i^*\mathcal{F}) \cong i^*(H^j(\mathcal{F}))$ .

*Proof.* As the definition of singular support is local in nature, the first claim follows immediately. The second is a consequence of the fact that  $i^*$  is exact.  $\square$

**Lemma 4.15.** *Let  $U$  be an open subset of a vector space  $E$ , and let  $V$  be a closed convex cone in  $E^*$ . If  $\mathcal{F} \in Sh(M)$  has the property that  $SS(\mathcal{F})$  is contained in  $U \times V$ , then  $SS(H^j(\mathcal{F})) \subset U \times V$  for all  $j$ .*

*Proof.* See [KS], Exercise VI.1.  $\square$

**Proposition 4.16.** *Let  $M$  be a manifold, and let  $\Lambda \subset T^*M$  be a cragged conical lagrangian. Then the  $t$ -structure on  $Sh(M)$  restricts to a  $t$ -structure on  $Sh(M, \Lambda)$ .*

*Proof.* We have to prove that for all objects  $\mathcal{F}$  in  $Sh(M, \Lambda)$ , and for all  $i \in \mathbb{Z}$ ,  $H^i(\mathcal{F})$  lies in  $Sh(M, \Lambda)$ , or equivalently that  $SS(H^i(\mathcal{F})) \subset \Lambda$ . By Lemma 4.14 we can assume that  $M$  is an open subset  $U$  in a vector space  $E$ , and that  $\Lambda$  is globally cragged. We use the notations of Definition 4.7 (1).

If  $p$  is a point in  $U$  we denote  $i_p \in I$  the index such that  $p \in Z_{i_p}$ . It is sufficient to show that for all  $p$ , the fiber  $SS(H^i(\mathcal{F}))_p$  is contained in  $V_{i_p}$ . We can consider an open neighborhood  $U'$  of  $p$  having the property that for all  $q \in U'$ ,  $i_p \leq i_q$ . Let  $i : U' \subset U$  be the inclusion. By Lemma 4.14  $SS(i^*\mathcal{F}) = SS(\mathcal{F}) \cap T^*U' \subset U' \times V_{i_p}$ . Applying Lemma 4.14 and then Lemma 4.15 we obtain the inclusion

$$SS(H^j(\mathcal{F})) \cap T^*U' = SS(i^*H^j(\mathcal{F})) = SS(H^j(i^*\mathcal{F})) \subset U' \times V_{i_p},$$

This implies in particular that  $SS(H^j(\mathcal{F}))_p \subset V_{i_p}$ .  $\square$

If  $\mathcal{S}$  is a stratification satisfying some extra conditions then  $Sh_c(M, \mathcal{S})$  can be described very explicitly as a category of quiver representations. We make use of the convenient recent reference [Ba], but remark that this circle of ideas goes back to MacPherson and others: for more information see for instance [Ka], and the discussion in the first Section of [Tr1].

Let us briefly summarize the results in [Ba] that we will need in the sequel. Let  $\mathcal{S}$  be an *acyclic stratification* of  $M$  in the sense of [Ba] Definition 17. For our purposes it is sufficient to remark that any triangulation of  $M$  yields such a stratification, and that any stratification in the sense of 4.2 admits a refinement that is acyclic. Section 4 of [Ba] contains a prescription that attaches to  $\mathcal{S}$  a directed quiver with relations  $(Q_{\mathcal{S}}, R_{\mathcal{S}})$ . Let  $A_{\mathcal{S}}$  be the path algebra of  $(Q_{\mathcal{S}}, R_{\mathcal{S}})$ , and let  $D_{dg}^b(\text{mod } A_{\mathcal{S}})$  be the triangulated dg category of finite  $A_{\mathcal{S}}$ -modules.

**Remark 4.17.** Recall that a directed quiver is a quiver which has no oriented cycles and loops. Note that as the quiver  $Q_{\mathcal{S}}$  is directed,  $A_{\mathcal{S}}$  has finite global dimension.

**Theorem 4.18** ([Ba] Theorem 21). *There is a quasi-equivalence*

$$\Phi : Sh_c(M, \mathcal{S}) \cong D_{dg}^b(\text{mod } A_{\mathcal{S}}).$$

Also,  $\Phi$  preserves the tautological  $t$ -structures.

## 5. THE COHERENT-CONSTRUCTIBLE CORRESPONDENCE

We start by reviewing the statement of the equivariant coherent-constructible (CC) correspondence of [FLTZ1, FLTZ2]. Next we extend to toric orbifolds partial results of Treumann [Tr] on the non-equivariant CC-correspondence. Building on these results in the next Section we prove one of our main theorems: we establish a complete statement of the non-equivariant CC-correspondence for a class of toric orbifolds satisfying some special combinatorial conditions.

We use the notations introduced in Section 3. Let  $\Sigma = (N, \Sigma, \beta)$  be a stacky fan. Henceforth, we assume that all stacky fans  $\Sigma$  are complete: that is, that their underlying ordinary fan  $\Sigma$  is complete. Recall that we write  $M = \text{Hom}(N, \mathbb{Z})$ ,  $M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}$ , and that  $\beta$  is a homomorphism of the form  $\beta : \mathbb{Z}^r = \oplus_{i=1}^r \mathbb{Z}e_i \rightarrow N$ . We set  $b_i = \beta(e_i)$  and  $B = \{b_1, \dots, b_r\}$ . For all  $\sigma \in \Sigma$ , we denote  $N_{\sigma} \subset N$  the sublattice generated by  $B \cap \sigma = \{w_1, \dots, w_d\}$ , and we denote  $M_{\sigma}$  the dual lattice  $M_{\sigma} = \text{Hom}(N_{\sigma}, \mathbb{Z})$ . If  $\sigma \in \Sigma$ , and  $\chi \in M_{\sigma}$ , we make the following definitions:

$$\sigma_{\chi}^{\perp} := \{x \in M_{\mathbb{R}} \mid \langle x, w_i \rangle = \langle \chi, w_i \rangle_{\sigma}\}, \quad \sigma_{\chi}^{\vee} := \{x \in M_{\mathbb{R}} \mid \langle x, w_i \rangle \geq \langle \chi, w_i \rangle_{\sigma}\},$$

where  $\langle \bullet, \bullet \rangle$  and  $\langle \bullet, \bullet \rangle_{\sigma}$  are the natural pairings. Following [FLTZ2] we define the conical Lagrangian subset  $\tilde{\Lambda}_{\Sigma} \subset M_{\mathbb{R}} \times N_{\mathbb{R}} = T^*M_{\mathbb{R}}$  by the formula:

$$\tilde{\Lambda}_{\Sigma} = \bigcup_{\tau \in \Sigma} \bigcup_{\chi \in M_{\tau}} \tau_{\chi}^{\perp} \times -\tau \subset M_{\mathbb{R}} \times N_{\mathbb{R}} = T^*M_{\mathbb{R}}.$$

Let  $Sh_{cc}(M, \tilde{\Lambda}_{\Sigma}) \subset Sh_c(M, \tilde{\Lambda}_{\Sigma})$  be the subcategory of sheaves with compact support. Denote  $T = N \otimes_{\mathbb{Z}} \mathbb{C}^*$  the torus acting on  $\mathcal{X}$ . In [FLTZ1] and [FLTZ2] Fang, Liu, Treumann and Zaslow prove the following theorem:

**Theorem 5.1** (Equivariant CC-correspondence, [FLTZ2] Theorem 7.6). *There is a quasi-equivalence of dg categories  $\tilde{\kappa} : \text{Perf}_T(\mathcal{X}_{\Sigma}) \rightarrow Sh_{cc}(M, \tilde{\Lambda}_{\Sigma})$ .*

The purpose of this Section is to extend work of Treumann [Tr] on the non-equivariant CC-correspondence for toric varieties to the larger class of toric orbifolds. We follow very closely the arguments in [Tr]. Let  $\mathbb{T}$  be the torus obtained as the quotient  $M_{\mathbb{R}}/M$  and let  $p$  be the quotient map,  $p : M_{\mathbb{R}} \rightarrow \mathbb{T}$ . Note that  $\tilde{\Lambda}_{\Sigma}$  is invariant under translation by elements in the integral lattice  $M$ . Denote  $\Lambda_{\Sigma}$  its quotient under the  $M$ -action:  $\Lambda_{\Sigma}$  is a conical Lagrangian subset of  $T^*\mathbb{T}$ . We denote  $\mathcal{F}org$  both the forgetful functor  $\mathcal{QCoh}_T(\mathcal{X}_{\Sigma}) \rightarrow \mathcal{QCoh}(\mathcal{X}_{\Sigma})$ , and its restriction to perfect complexes  $\text{Perf}_T(\mathcal{X}_{\Sigma}) \rightarrow \text{Perf}(\mathcal{X}_{\Sigma})$ . The following holds:

**Theorem 5.2.** *There is a quasi fully-faithful functor of triangulated dg categories*

$$\kappa : \text{Perf}(\mathcal{X}_{\Sigma}) \rightarrow Sh_c(\mathbb{T}, \Lambda_{\Sigma}),$$

*which makes the following diagram commute:*

$$\begin{array}{ccc} \text{Perf}_T(\mathcal{X}_{\Sigma}) & \xrightarrow{\tilde{\kappa}} & Sh_{cc}(M_{\mathbb{R}}, \tilde{\Lambda}_{\Sigma}) \\ \mathcal{F}org \downarrow & & \downarrow p! \\ \text{Perf}(\mathcal{X}_{\Sigma}) & \xrightarrow{\kappa} & Sh_c(\mathbb{T}, \Lambda_{\Sigma}). \end{array}$$

We prove Theorem 5.2 by constructing  $\kappa$  explicitly on a set of generators. Let us review briefly the main steps of the Theorem 5.1. In [FLTZ2] the authors attach to all pairs  $(\sigma, \chi)$ ,  $\sigma \in \Sigma$  and  $\chi \in M_\sigma$ , a quasi-coherent sheaf  $\Theta'_T(\sigma, \chi) \in \mathcal{Q}Coh_T(\mathcal{X}_\Sigma)$  and a constructible sheaf  $\Theta_T(\sigma, \chi) \in Sh_c(M_\mathbb{R})$  as follows:

$$\begin{aligned}\Theta_T(\sigma, \chi) &:= ((\sigma_\chi^\vee)^\circ \hookrightarrow M_\mathbb{R})! \mathbf{or}[\dim(M_\mathbb{R})], \\ \Theta'_T(\sigma, \chi) &:= (\mathcal{X}_\sigma \hookrightarrow \mathcal{X}_\Sigma)_* \mathcal{O}_{\mathcal{X}_\sigma}(\chi),\end{aligned}$$

where  $\mathcal{X}_\sigma \subset \mathcal{X}_\Sigma$  is the open substack associated to the cone  $\sigma$ , and  $(\sigma_\chi^\vee)^\circ \subset M_\mathbb{R}$  denotes the interior of  $\sigma_\chi^\vee$ . The functor  $\tilde{\kappa}$  is first defined as a functor between the larger categories  $\mathcal{Q}Coh(\mathcal{X}_\Sigma) \rightarrow Sh_c(M_\mathbb{R}, \tilde{\Lambda}_\Sigma)$ :  $\tilde{\kappa}$  sends  $\Theta'_T(\sigma, \chi)$  to  $\Theta_T(\sigma, \chi)$  and this assignment determines  $\tilde{\kappa}$  uniquely. The key fact is that  $\tilde{\kappa}$  restricts to the subcategories  $\mathcal{P}erf(\mathcal{X}_\Sigma)$  and  $Sh_{cc}(M_\mathbb{R}, \tilde{\Lambda}_\Sigma)$ , and defines a quasi-equivalence between them.

In the non-equivariant setting we consider the sheaves

$$\begin{aligned}\Theta(\sigma, \chi) &:= p_! \Theta_T(\sigma, \chi), \\ \Theta'(\sigma, \chi) &:= \mathcal{F}org(\Theta'_T(\sigma, \chi)).\end{aligned}$$

We start showing that the prescription  $\kappa(\Theta'(\sigma, \chi)) = \Theta(\sigma, \chi)$  defines a quasi fully-faithful embedding of dg categories  $\kappa : \mathcal{Q}Coh(\mathcal{X}_\Sigma) \rightarrow Sh_{qc}(\mathbb{T}, \Lambda_\Sigma)$ . Note that we have a natural action of  $M$  on  $M_\sigma$ : with slight abuse of notation, if  $m \in M$  and  $\chi \in M_\sigma$  we write  $m + \chi$  for  $(m)|_{N_\sigma} + \chi$ . More generally, let  $\tau$  and  $\sigma$  be two cones such that  $\tau \subset \sigma$ , and thus also  $\sigma^\vee \subset \tau^\vee$ . We have a canonical map  $M_\sigma \rightarrow M_\tau$  given by restriction to  $N_\tau$ . As before, if  $\chi_1 \in M_\sigma$  and  $\chi_2 \in M_\tau$  we simply write  $\chi_1 + \chi_2$  instead of  $(\chi_1)|_{N_\tau} + \chi_2$ . We remark that the sheaves  $\Theta(\sigma, \chi)$  and  $\Theta'(\sigma, \chi)$  depend on  $\chi$  only up to the action of  $M$ .

**Lemma 5.3.** *Let  $\sigma$  and  $\tau$  be cones in  $\Sigma$ . There is a  $\xi \in M$  such that  $\sigma_{\chi_1}^\vee \subset \tau_{\chi_2 + \xi}^\vee$  if and only if  $\tau \subset \sigma$  and  $\xi \in \tau_{\chi_2 - \chi_1}^\vee$ .*

*Proof.* Let us assume first that  $\tau \subset \sigma$ . The condition  $\xi \in \tau_{\chi_2 - \chi_1}^\vee$  implies that  $\xi - \chi_1 + \chi_2 \in \tau^\vee$  and hence  $\sigma^\vee - \chi_1 + \chi_2 + \xi \subset \tau^\vee$ . We conclude that  $\sigma_{\chi_1}^\vee \subset \tau_{\chi_2 + \xi}^\vee$ . Conversely, if  $\sigma_{\chi_1}^\vee \subset \tau_{\chi_2 + \xi}^\vee$ , then  $\sigma^\vee - \chi_1 + \chi_2 + \xi \subset \tau^\vee$ . It follows that  $\sigma^\vee \subset \tau^\vee$ , hence  $\tau \subset \sigma$  and  $-\chi_1 + \chi_2 + \xi \in \tau^\vee$ . The second condition is equivalent to the fact that  $\xi \in \tau_{\chi_2 - \chi_1}^\vee$ .  $\square$

**Proposition 5.4.** *We have*

$$\mathrm{Ext}^i(\Theta(\sigma, \chi_1), \Theta(\tau, \chi_2)) \cong \begin{cases} \mathbb{C}[\tau_{\chi_2 - \chi_1}^\vee \cap M] & \text{if } i = 0 \text{ and } \tau \subset \sigma, \\ 0 & \text{otherwise} \end{cases}$$

and

$$\mathrm{Ext}^i(\Theta'(\sigma, \chi_1), \Theta'(\tau, \chi_2)) \cong \begin{cases} \mathbb{C}[\tau_{\chi_2 - \chi_1}^\vee \cap M] & \text{if } i = 0 \text{ and } \tau \subset \sigma, \\ 0 & \text{otherwise.} \end{cases}$$

*Proof.* By adjunction we can write

$$\mathrm{hom}(\Theta(\sigma, \chi_1), \Theta(\tau, \chi_2)) = \mathrm{hom}(p_! \Theta_T(\sigma, \chi_1), \Theta(\tau, \chi_2)) \cong \mathrm{hom}(\Theta_T(\sigma, \chi_1), p^! \Theta(\tau, \chi_2)).$$

Also,  $p^! \Theta(\tau, \chi_2)$  decomposes as the direct sum  $\bigoplus_{\xi \in M} \Theta_T(\tau, \chi_2 + \xi)$ . Thus we have that

$$\mathrm{hom}(\Theta(\sigma, \chi_1), \Theta(\tau, \chi_2)) \cong \bigoplus_{\xi \in M} \mathrm{hom}(\Theta_T(\sigma, \chi_1), \Theta_T(\tau, \chi_2 + \xi)).$$

Then the first equality is a consequence of [FLTZ2] Proposition 5.5 and Lemma 5.3. The second equality follows in a similar way.  $\square$

Let  $\langle \Theta_T \rangle \subset Sh_c(M_{\mathbb{R}})$  and  $\langle \Theta'_T \rangle \subset QCoh_T(\mathcal{X}_{\Sigma})$  denote the full triangulated dg subcategories generated respectively by the objects  $\Theta_T(\sigma, \chi)$  and  $\Theta'_T(\sigma, \chi)$ . In the non-equivariant setting, we write  $\langle \Theta \rangle \subset Sh_{qc}(\mathbb{T})$  for the full triangulated dg subcategory generated by the objects  $\Theta(\sigma, \chi)$ , and  $\langle \Theta' \rangle \subset QCoh(\mathcal{X}_{\Sigma})$  the one generated by the objects  $\Theta'(\sigma, \chi)$ . The assignment  $\Theta'(\sigma, \chi) \mapsto \Theta(\sigma, \chi)$  determines uniquely a functor  $\kappa : \langle \Theta' \rangle \rightarrow \langle \Theta \rangle$ .

**Theorem 5.5.** *The functor  $\kappa : \langle \Theta' \rangle \rightarrow \langle \Theta \rangle$  is a quasi-equivalence, and fits in the following commutative diagram of dg categories*

$$\begin{array}{ccc} \langle \Theta'_T \rangle & \xrightarrow{\tilde{\kappa}} & \langle \Theta_T \rangle \\ \text{Forg} \downarrow & & \downarrow p_! \\ \langle \Theta' \rangle & \xrightarrow{\kappa} & \langle \Theta \rangle. \end{array}$$

*Proof.* The functor  $\kappa$  is a quasi-equivalence by Proposition 5.4. The commutativity of the diagram follows from the definition of  $\kappa$ .  $\square$

**Proposition 5.6.** *The dg category  $\mathcal{P}erf(\mathcal{X}_{\Sigma})$  is contained in  $\langle \Theta' \rangle$ .*

*Proof.* The equivariant CC-correspondence implies that  $\mathcal{P}erf_T(\mathcal{X}_{\Sigma}) \subset \langle \Theta'_T \rangle$ . The image of  $\mathcal{P}erf_T(\mathcal{X}_{\Sigma})$  under the functor  $\text{Forg}$  is by definition the set  $E_T$  of perfect complexes on  $\mathcal{X}$  admitting an equivariant structure. It follows that  $E_T \subset \langle \Theta' \rangle$ . Since  $\langle \Theta' \rangle$  is triangulated it must contain also the triangulated hull of  $E_T$ ,  $\langle E_T \rangle$ . By Corollary 3.8  $\langle E_T \rangle = \mathcal{P}erf(\mathcal{X}_{\Sigma})$ , and therefore  $\mathcal{P}erf(\mathcal{X}_{\Sigma})$  is contained in  $\langle \Theta' \rangle$ : this concludes the proof.  $\square$

In view of the previous proposition we can restrict  $\kappa$  to  $\mathcal{P}erf(\mathcal{X}_{\Sigma})$ . In order to prove Theorem 5.2 it remains only to show that  $\kappa$  sends  $\mathcal{P}erf(\mathcal{X}_{\Sigma})$  to  $Sh_c(\mathbb{T}, \Lambda_{\Sigma})$ : this is the analogue of Theorem 7.1 of [FLTZ2] in the non-equivariant setting.

*The proof of Theorem 5.2.* Corollary 3.8 shows that  $\mathcal{P}erf(\mathcal{X}_{\Sigma}) = \langle L \rangle$  where  $L$  is the set of line bundles on  $\mathcal{X}_{\Sigma}$ . It follows that the image of  $\mathcal{P}erf(\mathcal{X}_{\Sigma})$  under  $\kappa$  is equal to  $\langle \kappa(L) \rangle$ , where  $\kappa(L) := \{\kappa(\mathcal{L}), \mathcal{L} \in L\}$ . By Proposition 3.7 if  $\mathcal{L}$  is in  $L$  we can choose an equivariant representative  $\tilde{\mathcal{L}} \in \mathcal{P}erf_T(\mathcal{X}_{\Sigma})$  such that  $\mathcal{L} \cong \text{Forg}(\tilde{\mathcal{L}})$ . Also, using Theorem 5.5 we can write  $\kappa(\mathcal{L}) \cong p_!(\tilde{\kappa}(\tilde{\mathcal{L}}))$ . One consequence of Theorem 7.1 of [FLTZ2] is that  $\kappa(\tilde{\mathcal{L}})$  belongs to  $Sh_{cc}(M_{\mathbb{R}}, \tilde{\Lambda}_{\Sigma})$ . This implies that  $p_!(\tilde{\kappa}(\tilde{\mathcal{L}}))$  is a constructible sheaf and also that its singular support is contained in  $p(\tilde{\Lambda})$ :  $p_!(\tilde{\kappa}(\tilde{\mathcal{L}})) \in Sh_c(\mathbb{T}, \Lambda_{\Sigma})$ . Since  $Sh_c(\mathbb{T}, \Lambda_{\Sigma})$  is triangulated we conclude that

$$\kappa(\mathcal{P}erf(\mathcal{X}_{\Sigma})) \subset \langle \kappa(L) \rangle \subset Sh_c(\mathbb{T}, \Lambda_{\Sigma})$$

as we wanted to prove.  $\square$

**Remark 5.7.** We believe that there is an inclusion  $Sh_c(\mathbb{T}, \Lambda_{\Sigma}) \subset \langle \Theta \rangle$  but we do not know how to prove this in full generality. Indeed, this statement is equivalent to the fact that the quasi-fully faithful embedding  $\kappa : \mathcal{P}erf(\mathcal{X}_{\Sigma}) \rightarrow Sh_c(\mathbb{T}, \Lambda_{\Sigma})$  is also essentially surjective. In Section 6 we give a proof of the essential surjectivity of  $\kappa$  but only under additional assumptions on the combinatorics of  $\Sigma$ , see Theorem 6.11.

**5.1. Functoriality of  $\kappa$  and tensor products.** For completeness we prove that  $\kappa$  is functorial with respect to maps of stacky fans, and that  $\kappa$  intertwines the ordinary tensor product on  $\mathcal{P}\text{erf}(\mathcal{X}_\Sigma)$  and the convolution tensor product on  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$ . This is the analogue in the non-equivariant setting of Theorem 5.16 and Corollary 5.18 of [FLTZ2]. We remark that these results will not be used in the rest of the paper.

The next Definition is due to Borisov, Chen and Smith, see [BCS] Remark 4.5.

**Definition 5.8.** A *map of stacky fans* between  $\Sigma_1 = (N_1, \Sigma_1, \beta_1)$  and  $\Sigma_2 = (N_2, \Sigma_2, \beta_2)$  is a group homomorphism  $f : N_1 \rightarrow N_2$  with the following properties:

- (1) Denote  $f_\mathbb{R} : (N_1)_\mathbb{R} \rightarrow (N_2)_\mathbb{R}$  the homomorphism obtained by extension of scalars. Then  $f_\mathbb{R}$  maps cones in  $\Sigma_1$  to cones in  $\Sigma_2$ : that is, if  $\sigma_1 \in \Sigma_1$ , there is  $\sigma_2 \in \Sigma_2$  such that  $f_\mathbb{R}(\sigma_1) \subset \sigma_2$ .
- (2) If  $\sigma_1 \in \Sigma_1$  and  $\sigma_2 \in \Sigma_2$  are as in (1), then  $f(N_{\sigma_1}) \subset N_{\sigma_2}$ .

To a map of stacky fans  $f$  we can attach (see [FLTZ2] Section 5.6 for additional details):

- A map of vector spaces  $\tilde{v}_f : M_{2,\mathbb{R}} \rightarrow M_{1,\mathbb{R}}$ . As  $\tilde{v}_f^{-1}(M_{1,\mathbb{Z}})$  is contained in  $M_{2,\mathbb{Z}}$ ,  $\tilde{v}_f$  descends to a map  $v_f : \mathbb{T}_1 := M_{1,\mathbb{R}}/M_{1,\mathbb{Z}} \rightarrow \mathbb{T}_2 := M_{2,\mathbb{R}}/M_{2,\mathbb{Z}}$ .
- A map of stacks  $u_f : \mathcal{X}_{\Sigma_1} \rightarrow \mathcal{X}_{\Sigma_2}$  which is equivariant with respect to the torus actions: that is,  $u_f$  fits in the commutative diagram

$$\begin{array}{ccc} T_1 \times \mathcal{X}_{\Sigma_1} & \xrightarrow{u_f|_{T_1} \times u_f} & T_2 \times \mathcal{X}_{\Sigma_2} \\ m_1 \downarrow & & \downarrow m_2 \\ \mathcal{X}_{\Sigma_1} & \xrightarrow{u_f} & \mathcal{X}_{\Sigma_2}, \end{array}$$

where  $T_1 \subset \mathcal{X}_{\Sigma_1}$  and  $T_2 \subset \mathcal{X}_{\Sigma_2}$  are the open torus orbits, and  $m_1$  and  $m_2$  denote the respective actions on  $\mathcal{X}_{\Sigma_1}$  and  $\mathcal{X}_{\Sigma_2}$ .

**Theorem 5.9.** *Let  $f : \Sigma_1 \rightarrow \Sigma_2$  be a map of stacky fans. Assume that the underlying homomorphism  $f : N_1 \rightarrow N_2$  is injective. Then we have a commutative diagram of dg categories:*

$$\begin{array}{ccc} \langle \Theta'_{\Sigma_2} \rangle & \xrightarrow{\kappa_2} & \langle \Theta_{\Sigma_2} \rangle \\ u^* \downarrow & & \downarrow v_! \\ \langle \Theta'_{\Sigma_1} \rangle & \xrightarrow{\kappa_1} & \langle \Theta_{\Sigma_1} \rangle. \end{array}$$

*Proof.* We define a natural transformation  $\iota : v_! \circ \kappa_2 \rightarrow \kappa_1 \circ u^*$  on the generators  $\Theta'(\sigma, \chi)$ . Note that there are natural quasi-isomorphisms

- $v_! \circ \kappa_2(\Theta'_{\Sigma_2}(\sigma, \chi)) \cong v_! \Theta_{\Sigma_2}(\sigma, \chi) = v_! p_{2!} \Theta_{T_2, \Sigma_2}(\sigma, \chi) \cong p_{1!} \tilde{v}_! \Theta_{T_2, \Sigma_2}(\sigma, \chi) \cong p_{1!}(\tilde{v}_! \tilde{\kappa}_2(\Theta'_{\Sigma_2}(\sigma, \chi))),$
- $\kappa_1 \circ u^*(\Theta'_{\Sigma_2}(\sigma, \chi)) \cong \kappa_1 \circ \mathcal{F}org(u^* \Theta'_{T_2, \Sigma_2}(\sigma, \chi)) \cong p_{1!}(\tilde{\kappa}_1 u^* \Theta'_{T_2, \Sigma_2}(\sigma, \chi)).$

The natural transformation  $\iota$  can therefore be defined by applying  $p_{1!}$  to the natural transformation  $\tilde{\iota} : v_! \circ \tilde{\kappa}_2 \rightarrow \tilde{\kappa}_1 \circ u^*$  which is defined in [FLTZ2] Theorem 5.16. As  $\tilde{\iota}$  is a quasi-isomorphism,  $\iota$  is as well, and this concludes the proof.  $\square$

**Corollary 5.10.** *The functor  $\kappa : \mathcal{P}\text{erf}(\mathcal{X}_\Sigma) \rightarrow \text{Sh}_c(\mathbb{T}, \Lambda_\Sigma)$  is monoidal with respect to the usual tensor product of perfect complexes, and the convolution product on constructible sheaves.*

*Proof.* Let  $\Sigma \times \Sigma$  be the stacky fan of the product  $\mathcal{X}_\Sigma \times \mathcal{X}_\Sigma$ . The diagonal  $\Delta : N \rightarrow N \times N$  defines a injective morphism of stacky fans between  $\Sigma$  and  $\Sigma \times \Sigma$ . Note that  $u_f$  is the diagonal embedding  $\mathcal{X}_\Sigma \rightarrow \mathcal{X}_\Sigma \times \mathcal{X}_\Sigma$ , while  $v_f$  is the addition map:  $v_f : \mathbb{T} \times \mathbb{T} \rightarrow \mathbb{T}, v_f(a, b) = a + b$ . Let  $\mathcal{F}, \mathcal{G}$  be objects in  $\langle \Theta' \rangle_\Sigma$  (resp., in  $\langle \Theta \rangle_\Sigma$ ): the *box product* of  $\mathcal{F}$  and  $\mathcal{G}$  is defined by the formula  $\mathcal{F} \boxtimes \mathcal{G} := p_1^* \mathcal{F} \otimes p_2^* \mathcal{G} \in \langle \Theta' \rangle_{\Sigma \times \Sigma}$  (resp.  $\mathcal{F} \boxtimes \mathcal{G} \in \langle \Theta \rangle_{\Sigma \times \Sigma}$ ), where  $p_1, p_2$  are the projections,  $p_1, p_2 : \mathcal{X}_\Sigma \times \mathcal{X}_\Sigma \rightarrow \mathcal{X}_\Sigma$  (resp.  $p_1, p_2 : \mathbb{T} \times \mathbb{T} \rightarrow \mathbb{T}$ ). It follows immediately from the definition of  $\kappa$  that there are natural isomorphisms  $\kappa(\Theta'(\sigma_1, \chi_1) \boxtimes \Theta'(\sigma_2, \chi_2)) \cong \kappa(\Theta'(\sigma_1, \chi_1)) \boxtimes \kappa(\Theta'(\sigma_2, \chi_2))$ .

Denote  $\otimes^\circ$  the ordinary tensor product on  $\langle \Theta' \rangle$ , and  $\otimes^c$  the convolution tensor product on  $\langle \Theta \rangle$ . Applying Theorem 5.9 to  $\Delta$  we obtain natural isomorphisms

$$\kappa(\Theta'(\sigma_1, \chi_1) \otimes^\circ \Theta'(\sigma_2, \chi_2)) \cong \kappa(u_f^*(\Theta'(\sigma_1, \chi_1) \boxtimes \Theta'(\sigma_2, \chi_2))) \cong$$

$$v_{f!} \kappa(\Theta'(\sigma_1, \chi_1) \boxtimes \Theta'(\sigma_2, \chi_2)) = v_{f!}(\Theta(\sigma_1, \chi_1) \boxtimes \Theta(\sigma_2, \chi_2)) \cong \Theta(\sigma_1, \chi_1) \otimes^c \Theta(\sigma_2, \chi_2).$$

This implies that  $\kappa : \langle \Theta' \rangle \rightarrow \langle \Theta \rangle$  intertwines  $\otimes^\circ$  and  $\otimes^c$ . As these tensor structures restrict to the subcategories  $\mathcal{P}\text{erf}(\mathcal{X}_\Sigma)$  and  $\text{Sh}_c(\mathbb{T}, \Lambda_\Sigma)$  this proves the statement.  $\square$

## 6. CRAGGED STACKY FANS AND THE NON EQUIVARIANT CC-CORRESPONDENCE

In this Section we prove that the quasi fully-faithful functor  $\kappa : \mathcal{P}\text{erf}(\mathcal{X}_\Sigma) \rightarrow \text{Sh}_c(\mathbb{T}, \Lambda_\Sigma)$  is essentially surjective provided  $\Sigma$  satisfies some special properties, that is, it is *cragged* in the sense of Definition 6.3.

**Remark 6.1.** The essential surjectivity of  $\kappa$  is equivalent to the statement that the image of  $p_!$  generates  $\text{Sh}_c(\mathbb{T}, \Lambda_\Sigma)$ . Indeed Corollary 3.8 shows that the image of  $\mathcal{F} \text{org}$  generates  $\mathcal{P}\text{erf}(\mathcal{X}_\Sigma)$ , and  $\tilde{\kappa}$  is an equivalence by the equivariant CC-correspondence of [FLTZ2].

**Lemma 6.2.** *Let  $\mathcal{X}$  a toric DM stack, consider the point  $1 \in T \subset \mathcal{X}$ , and let  $k(1)$  be the skyscraper sheaf at 1. Then,  $\kappa(k(1)) \cong \mathbb{C}_\mathbb{T}$ .*

*Proof.* Denote  $j : T \hookrightarrow \mathcal{X}$  the inclusion of the open torus orbit. Computing the Koszul resolution of  $k(1)$  gives a complex

$$(1) \quad (j_* \mathcal{O}_T)^{\binom{n}{n}} \rightarrow \cdots \rightarrow (j_* \mathcal{O}_T)^{\binom{n}{2}} \rightarrow (j_* \mathcal{O}_T)^n \rightarrow j_* \mathcal{O}_T,$$

which is isomorphic to  $k(1)$  in the derived category of quasi-coherent sheaves on  $\mathcal{X}$ . Recall that  $\kappa$  is defined precisely by assigning quasi-constructible sheaves to quasi-coherent sheaves obtained as push-forward of structure sheaves on torus invariant open subsets. Thus, we can evaluate  $\kappa$  on the complex (1).

The functor  $\kappa$  admits a simple geometric interpretation on complexes of  $j_* \mathcal{O}_T$ -modules. Indeed, we can regard a complex of  $j_* \mathcal{O}_T$ -modules as a complex of modules over the group algebra  $\mathbb{C}[\pi_1(\mathbb{T})] = \mathbb{C}[T_1, T_1^{-1}, \dots, T_m, T_m^{-1}]$ . Further, the abelian category of modules over  $\mathbb{C}[\pi_1(\mathbb{T})]$  is naturally equivalent to the category of locally constant quasi-constructible sheaves on  $\mathbb{T}$ :  $\kappa$  maps a complex of  $j_* \mathcal{O}_T$ -modules to the corresponding complex of locally constant quasi-constructible sheaves on  $\mathbb{T}$ .

As a complex of  $\mathbb{C}[\pi_1(\mathbb{T})]$ -modules, (1) is the Koszul resolution of the trivial representation of  $\mathbb{C}[\pi_1(\mathbb{T})]$ ,

$$(2) \quad (\mathbb{C}[\pi_1(\mathbb{T})])^{\binom{n}{n}} \rightarrow \cdots \rightarrow (\mathbb{C}[\pi_1(\mathbb{T})])^{\binom{n}{2}} \rightarrow (\mathbb{C}[\pi_1(\mathbb{T})])^n \rightarrow \mathbb{C}[\pi_1(\mathbb{T})].$$

Thus  $\kappa(k(1))$  is the locally constant sheaf given by the trivial  $\mathbb{C}[\pi_1(\mathbb{T})]$ -module, i.e. the constant sheaf  $\mathbb{C}_{\mathbb{T}}$ . This concludes the proof.  $\square$

Let  $\Sigma = (N, \Sigma, \beta)$  be a stacky fan. Let  $\beta$  be of the form  $\beta : \mathbb{Z}^r = \bigoplus_{i=1}^r \mathbb{Z}e_i \rightarrow N$ . As in Section 5 we denote  $b_i = \beta(e_i)$  and  $B = \{b_1, \dots, b_r\}$ .

**Definition 6.3.** We say that  $\Sigma$  is *cragged* if the following two conditions are satisfied:

- (1) (exhaustiveness) Let  $S$  be a subset of  $\Sigma$ . Denote  $\langle S \rangle \subset N_{\mathbb{R}}$  the smallest convex cone containing all cones  $\sigma$  in  $S$ . Then there is a subset  $T \subset \Sigma$  such that  $\langle S \rangle = \bigcup_{\tau \in T} \tau$ .
- (2) (unimodularity) Let  $b_1, \dots, b_l$  be a set of linearly independent vectors in  $B$ , and let  $\rho_i = \mathbb{R}_{\geq 0} b_i$ . Exhaustiveness implies that  $\langle \rho_1, \dots, \rho_l \rangle = \bigcup_{\tau \in T} \tau$ . Let  $N_T \subset N$  be the sublattice generated by  $B \cap \bigcup_{\tau \in T} \tau$ . Then  $b_1, \dots, b_l$  form a  $\mathbb{Z}$ -basis for  $N_T$ .

**Proposition 6.4.** A stacky fan  $\Sigma$  is cragged if and only if  $\tilde{\Lambda}_{\Sigma}$  is a cragged Lagrangian.

*Proof.* We start by proving that if  $\Sigma$  is cragged then  $\Lambda_{\Sigma}$  is a cragged Lagrangian. Note that for all stacky fans  $\Sigma$ ,  $\tilde{\Lambda}_{\Sigma}$  satisfies the assumptions of Lemma 4.11. Thus it is sufficient to prove that if  $\Sigma$  is exhaustive and unimodular then  $\tilde{\Lambda}_{\Sigma}$  is pre-additive. We have to show that for all  $\phi \in M_{\mathbb{R}}$ ,  $(\tilde{\Lambda}_{\Sigma})_{\phi}$  is a convex cone. Denote  $S_{\phi}$  the subset of  $\Sigma$  consisting of all cones  $\tau$  such that  $\phi(N_{\tau})$  takes integral values, that is such that  $\phi \in M_{\tau}$ .

By exhaustiveness, there is a  $T_{\phi} \subset \Sigma$  such that  $\langle S_{\phi} \rangle = \bigcup_{\tau \in T_{\phi}} \tau$ . Using unimodularity we can find a subset  $B_{\phi} \subset B$  with the following properties:

- (1) the vectors in  $B_{\phi}$  form a basis of the sublattice generated by  $B \cap T_{\phi}$ ,
- (2) for each  $b \in B_{\phi}$  there is a  $\sigma \in S_{\phi}$  such that  $b$  belongs to  $\sigma$ .

This implies that  $T_{\phi} \subset S_{\phi}$ . Indeed, let  $\tau \in T_{\phi}$  and set  $B^{\tau} = B \cap \tau$ . The cone  $\tau$  lies in  $S_{\phi}$  if and only if  $\phi$  takes integral values on the elements of  $B^{\tau}$ . Since  $B^{\tau}$  is contained in the sublattice generated by  $B \cap \bigcup_{\tau \in T_{\phi}} \tau$ , all its elements can be written as integral linear combinations of vectors in  $B_{\phi}$ . By definition  $\phi$  takes integral values on  $B_{\phi}$ , therefore the same is true for  $B^{\tau}$  as well. These considerations give us an inclusion

$$-\langle S_{\phi} \rangle = \bigcup_{\tau \in T_{\phi}} -\tau \subseteq \bigcup_{\sigma \in S_{\phi}} -\sigma = (\tilde{\Lambda}_{\Sigma})_{\phi},$$

and the reverse inclusion is clear. Thus  $(\tilde{\Lambda}_{\Sigma})_{\phi} = -\langle S_{\phi} \rangle$ , and therefore  $(\tilde{\Lambda}_{\Sigma})_{\phi}$  is a convex cone as we needed to show.

Let us prove the other implication: if  $\tilde{\Lambda}_{\Sigma}$  is pre-additive then  $\Sigma$  is exhaustive and unimodular. Let  $S$  be a subset of  $\Sigma$ , denote  $N_S$  the sublattice of  $N$  generated by  $\langle S \rangle \cap B$  and denote  $M_S$  the dual lattice. Pick a minimal set of generators  $\phi_1^S, \dots, \phi_l^S$  of  $M_S$ . Choose preimages  $\phi_1, \dots, \phi_l$  in  $M_{\mathbb{R}}$  such that  $\phi_i$  is sent to  $\phi_i^S$  under the quotient map  $M_{\mathbb{R}} \rightarrow M_S \otimes_{\mathbb{Z}} \mathbb{R}$ . The fibers of  $\tilde{\Lambda}_{\Sigma}$  are naturally subsets of  $N_{\mathbb{R}}$ . Thus it makes sense to consider the intersection of the fibers  $(\tilde{\Lambda}_{\Sigma})_{\phi_i}$  as a subset of  $N_{\mathbb{R}}$ . It is immediate to see that  $\cap_i (\tilde{\Lambda}_{\Sigma})_{\phi_i} = \bigcup_{\sigma \in \langle S \rangle} -\sigma$ . Since  $\tilde{\Lambda}_{\Sigma}$  is pre-additive,  $(\tilde{\Lambda}_{\Sigma})_{\phi_i}$  is a convex cone for all  $i$ , and therefore  $\cap_i (\tilde{\Lambda}_{\Sigma})_{\phi_i} = \bigcup_{\sigma \in \langle S \rangle} -\sigma$  is

a convex cone as well. There is an inclusion  $\bigcup_{\sigma \subset \langle S \rangle} \sigma \subset \langle S \rangle$ . The reverse inclusion holds as well since  $\bigcup_{\sigma \in S} \sigma \subset \bigcup_{\sigma \subset \langle S \rangle} \sigma$ , and by definition  $\langle S \rangle$  is the smallest convex cone containing  $\bigcup_{\sigma \in S} \sigma$ . Thus  $\langle S \rangle = \bigcup_{\sigma \subset \langle S \rangle} \sigma$ : this proves that  $\Sigma$  is exhaustive.


Next we show that unimodularity holds as well. Suppose to the contrary that unimodularity fails: in the notation of the statement this means that there are one dimensional cones  $\rho_1, \dots, \rho_m$  such that  $\langle \rho_1, \dots, \rho_m \rangle = \bigcup_{\tau \in T} \tau$  but  $\{b_1, \dots, b_m\}$  is not a basis of the sublattice  $N_T$  generated by  $B \cap \bigcup_{\tau \in T} \tau$ . We define a homomorphism  $\phi : N \rightarrow \mathbb{R}$  having the following two properties:

- (1)  $\phi$  takes integer values on  $\{b_1, \dots, b_m\}$ ,
- (2) there is an element  $b_0 \in B \cap \bigcup_{\tau \in T} \tau$  such that  $\phi(b_0)$  is not an integer.

Denote  $\rho_0$  the one dimensional cone such that  $b_0 \in \rho_0$ . The conditions on  $\phi$  imply that  $-\rho_i$  is contained in  $(\tilde{\Lambda}_\Sigma)_\phi$  for all  $\rho \in \{1, \dots, n\}$ , while  $-\rho_0$  is not:  $-\rho_0 \notin (\tilde{\Lambda}_\Sigma)_\phi$ . Since  $\Lambda_\Sigma$  is pre-additive  $(\tilde{\Lambda}_\Sigma)_\phi$  is a convex cone and therefore contains  $-\langle \rho_1, \dots, \rho_m \rangle$  as a convex sub-cone. This gives a contradiction because then we would have  $-\rho_0 \subset -\langle \rho_1, \dots, \rho_m \rangle \subset (\tilde{\Lambda}_\Sigma)_\phi$ .  $\square$


We give below two examples of fans where either unimodularity or exhaustiveness fails.

**Example 6.5.** • Consider the fan  $\Sigma$  in  $\mathbb{R}^3$  given by  $\langle 0 \rangle$ , the one-dimensional cones  $\rho_i$ , the four faces  $\langle \rho_1, \rho_2 \rangle, \langle \rho_1, \rho_3 \rangle, \langle \rho_2, \rho_4 \rangle, \langle \rho_3, \rho_4 \rangle$  and the cone spanned by all the one-dimensional cones  $\langle \rho_1, \rho_2, \rho_3, \rho_4 \rangle$ .



Then  $\Sigma$  is not exhaustive: indeed the cone  $\langle \rho_1, \rho_2, \rho_3 \rangle$  cannot be written as a union of cones of  $\Sigma$ .

- All complete fans in  $\mathbb{R}^2$  are exhaustive, but they might fail to be unimodular: in fact unimodularity is a more restrictive condition than smoothness. Consider the smooth complete fan  $\Sigma$  in  $\mathbb{R}^2$  spanned by the one-dimensional cones  $\rho_1 := \mathbb{R}_{>0}(1, 0)$ ,  $\rho_2 := \mathbb{R}_{>0}(1, 1)$ ,  $\rho_3 := \mathbb{R}_{>0}(1, -1)$  and  $\rho_4 := \mathbb{R}_{>0}(-1, 0)$ :



If  $i, j$  are in  $\{1, \dots, 4\}$  denote  $\sigma_{ij}$  the two dimensional cone of  $\Sigma$  given by  $\sigma_{ij} = \langle \rho_i, \rho_j \rangle$ . Then  $\langle \rho_1, \rho_3 \rangle = \sigma_{12} \cup \sigma_{23}$  but the  $\mathbb{Z}$ -lattice spanned by  $N \cap (\sigma_{12} \cup \sigma_{23})$  is generated by  $(1, 1)$ ,  $(1, -1)$  and  $(1, 0)$ . Hence  $\Sigma$  is not unimodular.

**Corollary 6.6.**  $\Sigma$  is cagged if and only if  $\Lambda_\Sigma$  is a cagged Lagrangian.

*Proof.* For Lagrangians being cragged is a local condition, thus  $\Lambda_\Sigma$  is cragged if and only if  $\tilde{\Lambda}_\Sigma$  is cragged. This together with Proposition 6.4 proves the statement.  $\square$

The conditions of Definitions 6.3 are quite restrictive but they are verified in a number of interesting examples. For instance, all toric Fano surfaces have cragged fans. Also Propositions 6.9 and 6.10 below imply, in particular, that in all dimensions there are infinitely many examples of toric orbifolds with cragged fans.

**Definition 6.7.** Let  $\Sigma = (N, \Sigma, \beta)$  be a stacky fan such that  $N$  has rank  $n$  and  $\Sigma$  contains  $n + 1$  one dimensional cones. We say that  $\mathcal{X}_\Sigma$  is a *generalized weighted projective space*.

**Remark 6.8.** Equivalently, generalized weighted projective spaces are toric orbifolds with Picard number one. Generalized weighted projective spaces contain ordinary weighted projective spaces but form a strictly larger class, see [Bu] for several concrete examples. Note that the author of [Bu] calls these varieties “fake weighted projective spaces.” We follow a different convention here because the name “fake projective space” is most commonly used in algebraic geometry to refer to a different class of objects, see e.g. [PY].

**Proposition 6.9.** *If  $\Sigma = (N, \Sigma, \beta)$  is the stacky fan of a generalized weighted projective space then  $\Sigma$  is cragged.*

*Proof.* Let  $N = \mathbb{Z}^n$ . Then  $\Sigma$  is fully determined by the assignment of a set of  $n + 1$  vectors  $B = \{b_1 \dots b_{n+1}\}$  in  $N = \mathbb{Z}^n$ : they satisfy an equation of the form  $m_1 b_1 + \dots + m_{n+1} b_{n+1} = 0$  where all the coefficients are strictly positive integers. The one dimensional cones of  $\Sigma$  are given by  $\rho_i = \mathbb{R}_{\geq 0} b_i$ ,  $\beta : \bigoplus_{i=1}^{i=n+1} \mathbb{Z} e_i \rightarrow N$  maps  $e_i$  to  $b_i$ , and the fan  $\Sigma$  is the collection of all the cones spanned by proper subsets of  $\{\rho_1 \dots \rho_{n+1}\}$ . This implies in particular that  $\Sigma$  is exhaustive. Now let  $b_{i_1} \dots b_{i_l}$  be a set of linearly independent vectors in  $B$ . By definition  $\langle \rho_{i_1} \dots \rho_{i_l} \rangle$  is a cone in  $\Sigma$ . Thus the fact that  $b_{i_1} \dots b_{i_l}$  are a  $\mathbb{Z}$ -basis of the sublattice generated by  $B \cap \langle \rho_{i_1} \dots \rho_{i_l} \rangle = \{b_{i_1} \dots b_{i_l}\}$  is tautological:  $\Sigma$  is therefore unimodular as well.  $\square$

**Proposition 6.10.** *Let  $\mathcal{X}_\Sigma$  be a toric orbifold with cragged stacky fan  $\Sigma = (N, \Sigma, \beta)$ , and let  $G$  be a finite group acting on  $\mathcal{X}$  in a way that is compatible with the torus action. Then the stacky fan of the quotient  $[\mathcal{X}_\Sigma/G]$  is cragged as well.*

*Proof.* Note that  $G$  can be embedded as a finite subgroup of the torus  $T = N \otimes_{\mathbb{Z}} \mathbb{C}^*$  in such a way that the action of  $G$  on  $\mathcal{X}_\Sigma$  is induced from the action of  $T$  by restriction. The subgroup  $G$  determines an overlattice  $M'$  of  $M = \text{Hom}(N, \mathbb{Z})$ :  $M'$  is the lattice of characters of  $T$  which restrict to the trivial representation of  $G$ . By dualizing we get a surjective map  $q : N \rightarrow N'$ . The stacky fan of  $[\mathcal{X}_\Sigma/G]$  is given by  $\Sigma' = (N', \Sigma', q \circ \beta)$ , where  $\Sigma'$  is the set of cones  $\{q(\sigma) | \sigma \in \Sigma\}$ . Exhaustiveness and unimodularity of  $\Sigma'$  follow immediately from the same properties of  $\Sigma$ .  $\square$

The next Theorem is one of our main results.

**Theorem 6.11.** *If  $\mathcal{X}_\Sigma$  is a smooth toric orbifold such that  $\Sigma$  is cragged, then*

$$\kappa : \text{Perf}(\mathcal{X}_\Sigma) \rightarrow \text{Sh}_c(\mathbb{T}, \Lambda_\Sigma)$$

*is an equivalence of categories.*

*Proof.* As explained in Remark 6.1 it is sufficient to prove that the image of

$$p_! : Sh_c(M, \tilde{\Lambda}_\Sigma) \rightarrow Sh_c(\mathbb{T}, \Lambda_\Sigma)$$

generates  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$ . Informally, our argument depends on the availability of a kind of categorification of partitions of unity: we explain briefly this analogy before proceeding with the proof. The role of the unit function is played by the constant sheaf  $\mathbb{C}_\mathbb{T} \in Sh_c(\mathbb{T}, \Lambda_\Sigma)$ , which is the unit for the tensor product<sup>5</sup> in  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$ . We regard the image under  $\kappa$  of a line bundle  $\mathcal{L}$  as corresponding to a bump function (or rather, to the indicator function of an open subset): as a justification recall that if  $\tilde{\mathcal{L}}$  is an equivariant ample line bundle then  $\tilde{\kappa}(\mathcal{L}) = i_! \mathbb{C}_{\Delta_\mathcal{L}^\circ}$ , where  $\mathbb{C}_{\Delta_\mathcal{L}^\circ}$  is the constant sheaf on the interior of the lattice polytope  $\Delta_\mathcal{L}$  (see [FLTZ2]). When  $\Sigma$  is cragged we can construct a complex  $X^\bullet$  in  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$  with the property that  $X^i$  are direct sums of objects of the form  $\kappa(\mathcal{L})$  and  $\mathbb{C}_\mathbb{T}$  is a left convolution of  $X^\bullet$ : this is the key step in our argument, and we think of this as an analogue of a partition of unity.

Lemma 6.2 implies that there is a resolution

$$\mathcal{P}^\bullet = (\mathcal{P}^m \rightarrow \mathcal{P}^{m-1} \rightarrow \dots \rightarrow \mathcal{P}^0) \rightarrow k(1)$$

of the skyscraper sheaf  $k(1)$  such that  $\mathcal{P}^i$  is a direct sum of line bundles for all  $i$ . We regard  $\mathcal{P}^\bullet$  as a complex of objects in  $\text{Perf}(\mathcal{X}_\Sigma)$  and  $k(1)$  as the left convolution of  $\mathcal{P}^\bullet$ . Evaluating  $\kappa$  on  $\mathcal{P}^\bullet$  we obtain a complex in  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$ ,  $\kappa(\mathcal{P}^\bullet) = (\kappa(\mathcal{P}^m) \rightarrow \dots \rightarrow \kappa(\mathcal{P}^0))$  such that  $\kappa(k(1)) \cong \mathbb{C}_\mathbb{T}$  is its left convolution. We can use this to realize all objects of  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$  as iterated cones of objects lying in the image of  $p_!$  in the following manner.

Let  $\mathcal{F} \in Sh(\mathbb{T}, \Lambda_\Sigma)$ . We tensor  $\kappa(\mathcal{P}^\bullet)$  with  $\mathcal{F}$  and we get a complex

$$\kappa(\mathcal{P}^m) \otimes \mathcal{F} \rightarrow \dots \rightarrow \kappa(\mathcal{P}^0) \otimes \mathcal{F},$$

such that  $\mathcal{F} = \mathbb{C}_\mathbb{T} \otimes \mathcal{F}$  is its left convolution. This implies that  $\mathcal{F}$  lies in the subcategory

$$\langle \kappa(\mathcal{P}^m) \otimes \mathcal{F}, \dots, \kappa(\mathcal{P}^0) \otimes \mathcal{F} \rangle \subset Sh_c(\mathbb{T}).$$

Note that  $\kappa(\mathcal{P}^i) \otimes \mathcal{F}$  is an object of  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$  for all  $i$ : in fact, since  $\Lambda_\Sigma$  is cragged,  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$  is closed under tensor product by Corollary 4.13. We prove next that  $\kappa(\mathcal{P}^i) \otimes \mathcal{F}$  lies in the image of  $p_!$ . Write  $\mathcal{P}^i = \bigoplus_{j \in J} L_j$  where  $L_j$  are line bundles, and choose equivariant representatives  $\tilde{L}_j \in \text{Perf}_T(\mathcal{X}_\Sigma)$ . There are natural isomorphisms  $\kappa(\bigoplus_{j \in J} L_j) \cong \bigoplus_{j \in J} \kappa(L_j) \cong \bigoplus_{j \in J} p_!(\tilde{\kappa}(\tilde{L}_j))$ . Thus we have that

$$\mathcal{P}^i \otimes \mathcal{F} \cong \bigoplus_{j \in J} (p_!(\tilde{\kappa}(\tilde{L}_j)) \otimes \mathcal{F}) \cong \bigoplus_{j \in J} p_!(\tilde{\kappa}(\tilde{L}_j) \otimes p^! \mathcal{F}),$$

where the last isomorphism is given by the projection formula.<sup>6</sup> This completes the proof.  $\square$

<sup>5</sup>We stress that here we are working with the ordinary tensor product of constructible sheaves, as opposed to the *convolution* product which was discussed in Section 5.1.

<sup>6</sup>Note that  $p^! \mathcal{F} \in Sh_c(M, \tilde{\Lambda}_\Sigma)$  is not compactly supported: however the tensor product  $\tilde{\kappa}(\tilde{L}_j) \otimes p^! \mathcal{F}$  is, and therefore lies in  $Sh_{cc}(M, \tilde{\Lambda}_\Sigma)$  as required.

## 7. TILTING COMPLEXES

As an application of Theorem 6.11 we prove that if  $\Sigma$  is a cragged stacky fan, then  $\mathcal{P}\text{erf}(\mathcal{X}_\Sigma)$  has a tilting complex. Let  $A$  be a finite-dimensional algebra over  $\mathbb{C}$ .<sup>7</sup> Let us recall some notations:  $\text{mod}A$  is the abelian category of finite dimensional  $A$ -modules,  $D_{dg}^b(A)$  is the triangulated dg category of finite dimensional  $A$ -modules, and  $D^b(A)$  the homotopy category  $D^b(A) = D(D_{dg}^b(A))$ .  $D_{dg}^b(A)$  has a tautological  $t$ -structure with heart equal to  $\text{mod}A$ .

**Theorem 7.1.** *Let  $A$  be a finite dimensional algebra of finite global dimension. Let  $\mathcal{C}$  be a triangulated dg category. Assume that there is a quasi fully-faithful embedding of  $\mathcal{C}$  into  $D_{dg}^b(A)$  that satisfies the following conditions:*

- (1) *The restriction of the tautological  $t$ -structure of  $D_{dg}^b(A)$  is a well defined  $t$ -structure on  $\mathcal{C}$ .*
- (2) *The embedding  $\mathcal{C} \rightarrow D_{dg}^b(A)$  admits a left adjoint.*

*Then there exists a finite-dimensional algebra  $B$  having global dimension smaller or equal than the global dimension of  $A$ , and a quasi-equivalence  $\mathcal{C} \cong D_{dg}^b(B)$ .*

Let  $A$  and  $\mathcal{C}$  be as in the statement of the theorem. We denote  $\mathcal{C}^0 = \mathcal{C} \cap \text{mod}A$ :  $\mathcal{C}^0$  is the heart of the induced  $t$ -structure on  $\mathcal{C}$ . The homotopy category  $D(\mathcal{C})$  is a thick subcategory of  $D^b(A)$ , and the inclusion  $\mathcal{C} \rightarrow D_{dg}^b(A)$  admits a left adjoint which we denote  $L$ .

**Lemma 7.2.** *An object  $X \in D_{dg}^b(A)$  belongs to  $\mathcal{C}$  if and only if  $H^i(X) \in \mathcal{C}^0$  for all  $i \in \mathbb{Z}$ .*

*Proof.* Clearly if  $X \in \mathcal{C}$  then  $H^i(X) \in \mathcal{C}^0$ . We prove the other implication by induction on the number of indexes  $i$  for which  $H^i(X)$  is different from zero. Suppose that  $H^i(X)$  vanishes for all but one  $i$ . Then  $X \cong H^i(X)$  and as the objects of  $\mathcal{C}^0$  are contained in  $\mathcal{C}$  we have that  $X \in \mathcal{C}$ . Suppose now that the set of indices  $i$  such that  $H^i(X) \neq 0$  has cardinality  $n$ . Take  $j$  to be the maximal index such that  $H^j(X) \neq 0$ . In  $D_{dg}^b(A)$  we have a cofiber sequence  $Y \rightarrow X \rightarrow H^j(X)$ , where  $Y$  has non-vanishing cohomology in a strictly smaller set of degrees than  $X$ . By the inductive hypothesis  $Y \in \mathcal{C}$ . As  $\mathcal{C}$  is triangulated,  $X$  belongs to  $\mathcal{C}$  as well.  $\square$

Let  $\mathcal{A}$  and  $\mathcal{B}$  be two abelian categories. We say that an exact functor  $F : \mathcal{B} \rightarrow \mathcal{A}$  is *degree-wise fully faithful* if  $\text{Ext}_{\mathcal{B}}^i(A, B) \cong \text{Ext}_{\mathcal{A}}^i(FA, FB)$  for all  $i \in \mathbb{N}$ .

**Lemma 7.3.** *Let  $A$  and  $B$  be finite-dimensional algebras. Suppose that  $A$  has finite global dimension. If  $F : \text{mod}B \rightarrow \text{mod}A$  is a degree-wise fully faithful functor, then its extension to the dg enhancements  $F : D_{dg}^b(B) \rightarrow D_{dg}^b(A)$  is also quasi fully faithful. Further the global dimension of  $B$  is smaller or equal than the global dimension of  $A$ .*

*Proof.* Since  $F$  is degree-wise fully faithful, the  $A$ -module  $F(B)$  satisfies  $\text{Ext}_A^i(F(B), F(B)) = 0$  for all  $i \neq 0$ . It follows from [Ri] Theorem 2.12 and Propositions 3.1 and 3.2 that  $F$  induces a fully faithful embedding into  $D^b(A)$  of the triangulated category of perfect  $B$ -modules,  $\mathcal{P}\text{erf}(\text{mod}B)$ . Note that the maximum range of non vanishing  $\text{Ext}$ -groups between  $B$ -modules is bounded by that of  $A$ -modules. That is, the global dimension of  $B$  is bounded by the global dimension of  $A$ , and  $\mathcal{P}\text{erf}(\text{mod}B) \cong D^b(B)$ . Thus, there is a

<sup>7</sup>All algebras appearing in this paper are always assumed to be associative and unital.

fully faithful embedding  $D^b(B) \rightarrow D^b(A)$ . By the uniqueness of cg enhancements (see [LO][Proposition 2.6]) for  $D^b(B)$  and  $D^b(A)$ , the embedding lifts to a quasi fully faithful embedding  $F : D_{dg}^b(B) \rightarrow D_{dg}^b(A)$ . This completes the proof.  $\square$

Let  $(D_{\leq 0}, D_{> 0})$  be the aisles of the tautological  $t$ -structure on  $D^b(A)$ , and  $\tau_{\leq 0} : D^b(A) \rightarrow D_{\leq 0}$  and  $\tau_{> 0} : D^b(A) \rightarrow D_{> 0}$  be the truncation functors.

**Proposition 7.4.** *Let  $D(\mathcal{C})$  and  $D^b(A)$  be as in the statement of Theorem 7.1. Suppose now that  $M \cong \tau_{\leq 0}M$  for some  $M \in \mathcal{C}$ , that is  $M$  has zero homology only in positive degree. Then the left adjoint  $L(M)$  has zero homology in positive degree, that is  $L(M) \cong \tau_{\leq 0}L(M)$ .*

*Proof.* If  $M$  is an object in  $D^b(A)$  there is a unique object  $M^\perp \in D(\mathcal{C})^\perp$  such that  $M^\perp \rightarrow M \rightarrow L(M) \rightarrow M^\perp[1]$  is a distinguished triangle. We have the following commutative diagram whose middle row and columns are distinguished triangles:

$$\begin{array}{ccccc} \tau_{\leq 0}M^\perp & \longrightarrow & \tau_{\leq 0}M & \longrightarrow & \tau_{\leq 0}L(M) \\ \downarrow & & \downarrow & & \downarrow \\ M^\perp & \longrightarrow & M & \longrightarrow & L(M) \\ \downarrow & & \downarrow & & \downarrow \\ \tau_{> 0}M^\perp & \longrightarrow & \tau_{> 0}M & \longrightarrow & \tau_{> 0}L(M). \end{array}$$

Suppose now that  $M \cong \tau_{\leq 0}M$ . Then  $\tau_{> 0}M$  vanishes and  $\text{Hom}_{D^b(A)}(M, X) = 0$  for all  $X$  in  $D_{> 0}$ . As  $\tau_{> 0}L(M)$  lies in  $\mathcal{C}$ , it follows that

$$\text{Hom}_{D^b(A)}(M, \tau_{> 0}L(M)) \cong \text{Hom}_{D^b(A)}(L(M), \tau_{> 0}L(M)) = 0.$$

Thus  $\tau_{> 0}L(M) = 0$ , and we have that  $L(M) \cong \tau_{\leq 0}L(M)$ .  $\square$

We can now proceed with the proof the main theorem of this section.

*The proof of Theorem 7.1.* Denote by  $j$  the embedding of  $\text{mod}A$  into  $D^b(A)$  as the heart of the tautological  $t$ -structure. Consider the following commutative diagram:

$$\begin{array}{ccc} D(\mathcal{C}) & \xrightleftharpoons[i]{L} & D^b(A) \\ H^0 \downarrow & & H^0 \downarrow \uparrow j \\ \mathcal{C}^0 & \xrightarrow{i_0} & \text{mod}A. \end{array}$$

We denote  $L^0 : \text{mod}A \rightarrow \mathcal{C}^0$  the composition  $L^0 = H^0 L j$ . We prove first that  $L^0$  is a left adjoint of  $i_0$ . Let  $N \in \text{mod}A$  and  $M \in \mathcal{C}^0$ . We have that  $H^0 L(N) = \tau_{> -1} \tau_{\leq 0} L(N)$ . Since  $N$  is concentrated in degree zero by Remark 7.4,  $\tau_{\leq 0} L(N) \cong L(N)$ . Thus

$$\text{Hom}_{\mathcal{C}^0}(L^0 N, M) \cong \text{Hom}_{D_{dg}^b(A)}(\tau_{> -1} L(N), M).$$

We apply  $\text{Hom}(-, M)$  to the distinguished triangle

$$\tau_{\leq -1} L(N) \rightarrow L(N) \rightarrow \tau_{> -1} L(N) \rightarrow \tau_{\leq -1} L(N)[1].$$

As  $M$  is concentrated in degree zero, we deduce that

$$\mathrm{Hom}_{D^b(A)}(\tau_{>-1}LN, M) \cong \mathrm{Hom}(LN, M) \cong \mathrm{Hom}(N, M),$$

that is,  $L^0$  is a left adjoint of  $i_0$ .

Left adjoints of exact functors preserve projective generators. Thus  $L^0(A)$  is a projective generator of  $\mathcal{C}^0$ , and  $\mathcal{C}^0$  is equivalent to  $\mathrm{mod}B$  where  $B$  is equal to  $\mathrm{End}_{\mathrm{mod}A}(L^0(A))$ . The equivalence  $\mathrm{mod}B \rightarrow \mathcal{C}^0$  is given by the functor  $- \otimes_B L^0(A)$ . Consider the composition  $F := i_0(- \otimes_B L^0(A))$ ,  $F$  is a degree-wise fully faithful functor from  $\mathrm{mod}B$  to  $\mathrm{mod}A$ . By Lemma 7.3,  $F$  extends to a fully faithful embedding  $F : D_{dg}^b(B) \rightarrow D_{dg}^b(A)$  whose image consists of complexes with cohomology in  $\mathcal{C}^0$ . By Lemma 7.2, the image of  $F$  lies in  $\mathcal{C}$ . Conversely  $D(\mathcal{C})$  is generated by  $\mathcal{C}^0$  as a triangulated category. This shows that there is an equivalence  $\mathcal{C} \cong D_{dg}^b(B)$ .  $\square$

**Theorem 7.5.** *Let  $\mathcal{X}_\Sigma$  be a smooth toric orbifold such that  $\Sigma$  is cragged. Then there is a finite-dimensional algebra  $B$  of finite global dimension such that  $\mathrm{Perf}(\mathcal{X}_\Sigma)$  is quasi-equivalent to  $D_{dg}^b(B)$ . In particular  $\mathrm{Perf}(\mathcal{X}_\Sigma)$  contains a tilting complex.*

*Proof.* Since  $\Lambda_\Sigma$  is cragged, the tautological  $t$ -structure of  $Sh_c(\mathbb{T})$  restricts to  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$ . Let  $\mathcal{S}$  be a stratification of  $\mathbb{T}$  that is acyclic and is such that all complexes in  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$  have cohomology that is constructible with respect to  $\mathcal{S}$ : this gives an embedding of  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$  into  $Sh_c(\mathbb{T}, \mathcal{S})$ . By Theorem 4.18 there is a quasi-equivalence  $Sh_c(\mathbb{T}, \mathcal{S}) \cong D_{dg}^b(A_\mathcal{S})$  that preserves the  $t$ -structures. Thus we obtain a quasi fully-faithful functor

$$J : Sh_c(\mathbb{T}, \Lambda_\Sigma) \rightarrow D_{dg}^b(A_\mathcal{S}),$$

that is compatible with the  $t$ -structures. Also,  $J$  has a left adjoint. Indeed, since  $\mathcal{X}_\Sigma$  is a smooth and proper DM stack,  $\mathrm{Perf}(\mathcal{X}_\Sigma)$  is a saturated dg category in the sense of [To]. In particular, by Proposition 13 of [To] all quasi fully-faithful functors with source  $\mathrm{Perf}(\mathcal{X}_\Sigma)$  admit both a left and a right adjoint, and the same holds for the quasi-equivalent category  $Sh_c(\mathbb{T}, \Lambda_\Sigma)$ . This completes the proof, as we can apply Theorem 7.1 and obtain a chain of quasi-equivalences

$$\mathrm{Perf}(\mathcal{X}_\Sigma) \cong Sh_c(\mathbb{T}, \Lambda_\Sigma) \cong D_{dg}^b(B).$$

$\square$

**Remark 7.6.** For weighted projective spaces stronger results are known. Borisov and Hua [BH] prove that they admit a full strong exceptional collection of line bundles, this generalizes Beilinson's famous theorem [Be] on the derived category of ordinary projective spaces.

## REFERENCES

- [A1] M. Abouzaid, “Homogeneous coordinate rings and mirror symmetry for toric varieties,” *Geom. Topol.* **10** 1097–1156 (2006).
- [A2] M. Abouzaid, “Morse homology, tropical geometry, and homological mirror symmetry for toric varieties,” *Selecta Math. (N.S.)* **15** (2) 189–270 (2009).
- [AKO1] D. Auroux, L. Katzarkov, D. Orlov, “Mirror symmetry for del Pezzo surfaces: vanishing cycles and coherent sheaves,” *Invent. Math.* **166** (3) 537–582 (2006).
- [AKO2] D. Auroux, L. Katzarkov, D. Orlov, “Mirror symmetry for weighted projective planes and their noncommutative deformations,” *Ann. of Math. (Second Series)* **167** (3) 867–943 (2008).

- [Bal] M. Ballard “Meet homological mirror symmetry,” in *Modular forms and string duality*, Fields Inst. Commun., 54, Amer. Math. Soc., Providence, RI, 191–224, (2008).
- [Ba] A. Balthazar, “Formality of the Constructible Derived Category for Spheres: A Combinatorial and a Geometric Approach,” *Mediterranean Journal of Mathematics*, **6** (4), 403–430 (2009).
- [Be] A. Beilinson, “Coherent sheaves on  $\mathbb{P}^n$  and problems in linear algebra,” *Funct. An. and Appl.* **12**(3) (1978) 68–69.
- [Bo] A. Bondal, “Derived categories of toric varieties,” in *Convex and Algebraic geometry, Oberwolfach conference reports, EMS Publishing House* **3** 284–286 (2006).
- [BR] A. Bondal, “Integrable systems related to triangulated categories,” talk at the MSRI workshop on Generalized McKay Correspondences and Representation Theory on 3/21/2006. The video is available at [www.msri.org/communications/vmath/VMathVideos/VideoInfo/2469/show\\_video](http://www.msri.org/communications/vmath/VMathVideos/VideoInfo/2469/show_video)
- [BCS] L. A. Borisov, L. Chen, G. G. Smith, “The orbifold Chow ring of toric Deligne-Mumford stacks,” *J. Am. Math. Soc.* **18** (2005), no. 1, 193–215.
- [BH] L. A. Borisov and R. P. Horja, “On the  $K$ -theory of smooth toric DM stacks,” *Snowbird lectures on string geometry*, 21–42, *Contemp. Math.* **401**, Amer. Math. Soc., Providence, RI, 2006.
- [BHu] L. A. Borisov, Z. Hua, “On the conjecture of King for smooth toric Deligne–Mumford stacks,” *Adv. in Math.* **221**, Issue 1, 277–301(2009).
- [Bu] W. Buczyńska, “Fake weighted projective spaces,” [arXiv:0805.1211](https://arxiv.org/abs/0805.1211).
- [CM] L. Costa, R. M. Miró-Roig, “Tilting sheaves on toric varieties,” *Math. Zeit.* **248** Issue 4, 849–865 (2004).
- [Dr] V. Drinfeld, “DG quotients of DG categories,” *J. Algebra* **272** no. 2, 643–691 (2004).
- [Ef] A. I. Efimov, “Maximal lengths of exceptional collections of line bundles,” [arXiv:1010.3755](https://arxiv.org/abs/1010.3755).
- [Fa] B. Fang, “Homological mirror symmetry is T-duality for  $\mathbb{P}^n$ ,” *Commun. Number Theory Phys.* **2**, 719–742 (2008).
- [FLTZ1] B. Fang, D. Treumann, C.-C. Liu and E. Zaslow, “A categorification of Morelli’s theorem,” *Invent. Math.* **186**(1) 179–214 (2011).
- [FLTZ2] B. Fang, D. Treumann, C.-C. Liu and E. Zaslow, “The Coherent-Constructible Correspondence for toric Deligne-Mumford stacks,” [arXiv:0911.4711](https://arxiv.org/abs/0911.4711).
- [FLTZ3] B. Fang, C.-C. Liu, D. Treumann, E. Zaslow, “T-duality and homological mirror symmetry for toric varieties” *Adv. Math.* **229**(3), 1875–1911(2012).
- [FMN] B. Fantechi, E. Mann and F. Nironi “Smooth toric DM stacks,” *J. Reine Angew. Math.* 648 (2010), 201–244.
- [FOOO] K. Fukaya, Y.-G. Oh, H. Ohta, K. Ono, “Lagrangian intersection Floer theory: anomaly and obstruction.” Part I and II. *AMS/IP Studies in Advanced Mathematics*, International Press, Somerville, MA, (2009).
- [GM] S. I. Gelfánd, Y. I. Manin, “Methods of homological algebra,” *Springer Monographs in Mathematics*, Springer-Verlag, 2nd ed. 2002.
- [GS] S. Guillermou, and P. Schapira, “Microlocal theory of sheaves and Tamarkin’s non displaceability theorem,” [arXiv 1106.1576](https://arxiv.org/abs/1106.1576)
- [HP] L. Hille, and M. Perling “A counterexample to King’s conjecture,” *Compos. Math.* **142** no. 6, 1507–1521(2006).
- [Iw] I. Iwanari, “The category of toric stacks,” *Comp. Math.* **145** no. 3, 718–746 (2009).
- [Ka] M. Kapranov, “Mutations and Serre functors in constructible sheaves,” *Funct. Anal. and Appl.* 24, 155–156 (1990).
- [Kaw] Y. Kawamata, “Derived categories of toric varieties II,” *Michigan Math. J.* **62** (2) 225–448 (2013).
- [Ke] B. Keller, “On differential graded categories,” *International Congress of Mathematicians. Vol. II*, Eur. Math. Soc., Zürich, 151–190 (2006).
- [Ki] A. D. King, “Tilting bundles on some rational surfaces,” preprint available at <http://www.maths.bath.ac.uk/~masadk/papers/tilt.ps>
- [KS] M. Kashiwara and P. Schapira, “Sheaves on Manifolds,” *Grundlehren der Mathematischen Wissenschaften* **292**, Springer-Verlag, 1994.

- [Ko] M. Kontsevich, “Lectures at ENS,” Paris, Spring 1998, notes taken by J. Bellaïche, J.-F. Dat, I. Marin, G. Racinet and H. Randriambololona.
- [LO] V. Lunts and D. Orlov, “Uniqueness of enhancement for triangulated categories,” *J. Amer. Math. Soc.* **23** (2010), 853–908.
- [Lu] J. Lurie, “Higher algebra,” available at <http://www.math.harvard.edu/~lurie/papers/HigherAlgebra.pdf>
- [N] D. Nadler, “Microlocal Branes are Constructible Sheaves,” *Sel. Math.* **15** (4), 563–619 (2009).
- [NZ] D. Nadler and E. Zaslow, “Constructible sheaves and the Fukaya category,” *J. Amer. Math. Soc.* **22** 233–286 (2009).
- [Po] A. Polishchuk, “Perverse sheaves on a triangulated space,” *Math. Res. Lett.*, Vol. 4, 191–199 (1997).
- [PY] G. Prasad, S.-K. Yeung, “Fake projective planes,” *Invent. Math.*, **168** (2), 321–370 (2007).
- [Ri] J. Rickard, “Morita theory for derived categories” *J. London Math. Soc.* **2** 39, 436–456 (1989).
- [Ro] M. Romagny, “Group actions on stacks and applications,” *Michigan Math. J.*, **53**(1) 209–236 (2005).
- [S] P. Seidel, “Fukaya categories and Picard-Lefschetz theory,” *Zürich Lectures in Advanced Mathematics*. European Mathematical Society (EMS), Zürich, (2008).
- [SYZ] A. Strominger, S.-T. Yau, and E. Zaslow “Mirror symmetry is T-duality,” *Nuclear Physics B* **479** (12) (1996), 243–259.
- [To] Bertrand Toën, *Lectures on DG-categories*, available at [www.math.univ-montp2.fr/~toen](http://www.math.univ-montp2.fr/~toen).
- [Tr1] D. Treumann, “Exit paths and constructible stacks,” *Comp. Math.* **145** 1504–1532 (2009).
- [Tr] D. Treumann, “Remarks on the nonequivariant coherent-constructible correspondence for toric varieties,” *arXiv* 1006.5756.
- [Vi] A. Vistoli, “Intersection theory on algebraic stacks and their moduli spaces,” *Invent. Math.*, **97** 613–670 (1987).
- [Vy] M. Vybornov, “Constructible sheaves on simplicial complexes and Koszul duality,” *Math. Res. Lett.*, Vol. 5, 675–683 (1998).
- [Wh] H. Whitney, “Tangents to an analytic variety,” *Ann. of Math. Second Series*, **81** (3), 496–549, (1965).

SARAH SCHEROTZKE, MATHEMATICAL INSTITUTE OF THE UNIVERSITY OF BONN, ENDENICHER ALLEE 60, 53115 BONN, GERMANY

*Email address:* [sarah@math.uni-bonn.de](mailto:sarah@math.uni-bonn.de)

NICOLÒ SIBILLA, MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111 BONN, GERMANY

*Email address:* [sibilla@mpim-bonn.mpg.de](mailto:sibilla@mpim-bonn.mpg.de)