



SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI

SISSA Digital Library

Isomonodromic deformations along a stratum of the coalescence locus

Original

Isomonodromic deformations along a stratum of the coalescence locus / Guzzetti, Davide. - In: JOURNAL OF PHYSICS. A, MATHEMATICAL AND THEORETICAL. - ISSN 1751-8113. - 55:45(2022), pp. 1-59. [10.1088/1751-8121/ac9ba8]

Availability:

This version is available at: 20.500.11767/130130 since: 2022-11-15T18:30:54Z

Publisher:

Published

DOI:10.1088/1751-8121/ac9ba8

Terms of use:

Testo definito dall'ateneo relativo alle clausole di concessione d'uso

Publisher copyright

IOP- Institute of Physics

This version is available for education and non-commercial purposes.

note finali coverpage

(Article begins on next page)

Isomonodromic deformations along a stratum of the coalescence locus

Davide Guzzetti

SISSA, Via Bonomea, 265, 34136 Trieste & INFN, Sezione di Trieste, Italy.

E-mail: guzzetti@sissa.it

ORCID ID: 0000-0002-6103-6563

Abstract: We consider deformations of a differential system with Poincaré rank 1 at infinity and Fuchsian singularity at zero along a stratum of a coalescence locus. We give necessary and sufficient conditions for the deformation to be strongly isomonodromic, both as an explicit Pfaffian system (integrable deformation) and as a non linear system of PDEs on the residue matrix A at the Fuchsian singularity. This construction is complementary to that of [13]. For the specific system here considered, the results generalize those of [26], by giving up the generic conditions, and those of [3], by giving up the Lidskii generic assumption. The importance of the case here considered originates from its applications in the study of strata of Dubrovin-Frobenius manifolds and F -manifolds.

Keywords: Non generic Isomonodromy Deformations, Stokes phenomenon, Resonant Irregular Singularity, Stokes matrices, Monodromy data

Contents

1 Introduction	2
2 Preliminaries	8
3 Weak isomonodromic deformations	9
3.1 Proof of Theorem 3.1	12
4 Canonical solutions of (1.1) at $z = \infty$	14
5 More on the Levelt form	17
6 Strong Isomonodromy Deformations	20
7 The 3-dimensional case	33
8 A few words on the 4-dimensional case	37
9 Example: the Caustic of a Dubrovin-Frobenius manifold in a generic case	39
10 Appendix. Proof of Lemma 6.1	53

Notation. For an $n \times n$ matrix A_k , we denote the matrix entries by either $(A_k)_{ij}$ or $A_{ij}^{(k)}$, where $i, j \in \{1, \dots, n\}$. We can partition A_k into s^2 blocks of dimension $p_a \times p_b$, where $a, b = 1, \dots, s$ and $p_1 + \dots + p_s = n$. The block labelled by a, b , of dimension $p_a \times p_b$ will be denoted by $A_{[a,b]}^{(k)}$.

1 Introduction

In the work [13], and in the related [12, 19, 20, 21, 22], we have studied an $n \times n$ matrix differential system of the shape (1.1) below, with an irregular singularity at $z = \infty$ and a Fuchsian one at $z = 0$, with leading term at ∞ given by a diagonal matrix $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$, whose eigenvalues $\lambda = (\lambda_1, \dots, \lambda_n)$ vary in a polydisc of \mathbb{C}^n . The polydisc contains a *coalescence locus*, where some eigenvalues merge, namely $\lambda_j - \lambda_k \rightarrow 0$ for some $1 \leq j \neq k \leq n$. For this system, we have proved that a monodromy preserving deformation theory can be well defined (in an analytic way) with constant monodromy data on the whole polydisc, including the coalescence locus. This result, which generalizes the theory of Jimbo, Miwa and Ueno [26], is possible if the vanishing conditions $A_{jk}(\lambda) \rightarrow 0$ hold when $\lambda_j - \lambda_k \rightarrow 0$.

In this paper, we consider an n -dimensional differential system

$$\frac{dY}{dz} = \left(\Lambda + \frac{A(\lambda)}{z} \right) Y, \quad \lambda = (\lambda_1, \dots, \lambda_s) \in \mathbb{D} \subset \mathbb{C}^s, \quad (1.1)$$

where \mathbb{D} is a polydisc and

$$\Lambda = \lambda_1 I_{p_1} \oplus \dots \oplus \lambda_s I_{p_s} := \text{diag}(\underbrace{\lambda_1, \dots, \lambda_1}_{p_1}, \underbrace{\lambda_2, \dots, \lambda_2}_{p_2}, \dots, \underbrace{\lambda_s, \dots, \lambda_s}_{p_s})$$

$$I_{p_j} = p_j\text{-dimensional identity matrix, } p_1 + \dots + p_s = n.$$

We can think of λ as the parameters varying *within a stratum* of a coalescence locus, specified by p_1, \dots, p_s . We would like to establish the full isomonodromy deformation theory within this stratum.

The deformation considered here is complementary to that of [13], because it occurs within the prefixed stratum, while in [13] the deformation takes place in a domain containing the coalescence set, the latter being included in the range of deformation under specific vanishing conditions on A . The problem of the present paper is therefore different from [13]: here A will be any matrix and we do not suppose that the entries of A corresponding to equal eigenvalues of Λ are zero. The deformation theory that we will develop cannot be deduced either from [13] or [26]. This theory is realized in an important geometric setting, namely at the nilpotent locus of a Dubrovin-Frobenius manifold [16], called *caustic* [24]. This application will be discussed in Section 9, for the type of caustics geometrically described in [28].

In the sequel, it will be convenient to partition A into blocks $A_{[i,j]}$, $i, j = 1, \dots, s$, of dimension $p_i \times p_j$, inherited from Λ . We will work in the following analytic setting.

Assumption 1.

- The polydisc \mathbb{D} is sufficiently small so that, as λ varies in \mathbb{D} , the Stokes rays defined in (1.2) below do not cross the half-lines $\arg z = \tau + k\pi$, $k \in \mathbb{Z}$, where $\tau \in \mathbb{R}$ is fixed, and called an **admissible direction**.
- $A(\lambda)$ is holomorphic in \mathbb{D} .

The Stokes rays in the assumption are the rays in the universal covering \mathcal{R} of $\overline{\mathbb{C}} \setminus \{0, \infty\}$ defined by

$$\Re((\lambda_i - \lambda_j)z) = 0, \quad \Im((\lambda_i - \lambda_j)z) < 0, \quad z \in \mathcal{R}. \quad (1.2)$$

In this paper, we establish the necessary and sufficient conditions for (1.1) to be strongly isomonodromic on the polydisc \mathbb{D} . The notion of “strong isomonodromy” is implicit in the seminal paper [26], meaning that *all essential monodromy data* (monodromy exponents, connection matrices, Stokes matrices, see Definition 6.1) are independent of λ . The adjective “strong” was introduced in [19], to point out that a system may just be “weakly” isomonodromic, namely with constant monodromy matrices, but with possibly non-constant essential monodromy data.

In the isomonodromy theory of [26], several assumptions are made to assure that the differential system is generic (the deformation is called admissible). In case of (1.1), the eigenvalues of A are not allowed to differ by integers (so A is in particular diagonalizable), and Λ has pairwise distinct eigenvalues.

In the paper [3], the isomonodromy deformation theory has been extended to rational connection with both Fuchsian and irregular singularities of any Poincaré rank, without several of the assumptions of [26]. The residue matrices at the Fuchsian singularities are not subject to restrictions, while the leading matrix at an irregular singularity can have any Jordan form, but with a *prefixed* Jordan type $\lambda_1^{n_1} \lambda_2^{n_2} \dots \lambda_K^{n_K}$ (in a notation due to Arnol’d [2]). For example, in case of (1.1), the Jordan type is prefixed to be

$$\underbrace{\lambda_1 \dots \lambda_1}_{p_1} \underbrace{\lambda_2 \dots \lambda_2}_{p_2} \dots \underbrace{\lambda_s \dots \lambda_s}_{p_s}. \quad (1.3)$$

Besides the prefixed Jordan type, another important assumption of [3] is that the next sub-leading matrix (in our case A) at an irregular singularity must be *Lidskii generic*, according to definition 2.1 in [3]. In our case, this means that each diagonal block of A (with block partition inherited from Λ) must have distinct and nonzero eigenvalues. Theorem 5.3 of [3] states that the deformation is isomonodromic (preserving a set of monodromy data, which include the Stokes matrices) if and only if a class of fundamental matrix solutions satisfy a certain Pfaffian system. This result generalizes theorem 3.1 of [26]. Moreover, [3] studies the generalization of the isomonodromic τ -function.

Remark 1.1. Given a differential system $\frac{dY}{dz} = \mathfrak{A}(z, \lambda)Y$ such that the deformation λ does not satisfying some admissibility conditions of [26], by *generalization* of [26] we mean: find necessary and sufficient conditions ensuring that *all essential monodromy data* (strong isomonodromy) of the differential system are constant. These conditions are of the type: constant data if and only if all the canonical solutions satisfy a Pfaffian system $dY = \omega Y$, with a very specific ω (such as (1.4) below); or if and only if the coefficients in $\mathfrak{A}(z, \lambda)$ satisfy certain non linear PDEs, called isomonodromy deformation equations (like the equations (1.6) below).

As mentioned in the beginning, an extension of the isomonodromy deformation theory has been achieved in [13] for a system such as (1.1) when the Jordan type of the leading matrix Λ changes within a polydisc of \mathbb{C}^n . In [13], A is any, while

$$\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$$

has n eigenvalues $\lambda = (\lambda_1, \dots, \lambda_n)$ varying in a polydisc containing a *coalescence locus* where $\lambda_j = \lambda_k$ for some $j \neq k$. The extension of [13] can be done under the condition that the entries $A_{jk}(\lambda) \rightarrow 0$ for $\lambda_j - \lambda_k \rightarrow 0$.

In this paper, the deformation parameters $\lambda = (\lambda_1, \dots, \lambda_s)$ are assumed to vary inside a stratum of the coalescence locus, namely Λ has $s < n$ eigenvalues varying in the polydisc \mathbb{D} specified before. In this sense, the Jordan type (1.3) of Λ is fixed, as in [3]. Nevertheless, we drop any assumption on A , and we do not assume Lidskii generic conditions.

We mention that in [11] isomonodromy deformations are defined for a system of type (1.1), with coefficients in the Lie algebra of an arbitrary complex algebraic group G . In the specific case we consider here the coefficients are $n \times n$ complex matrices, so that $G = GL_n(\mathbb{C})$. In this case, the assumptions of [11] require that Λ is diagonal¹ with a prefixed Jordan type (1.3), invariant by the deformation λ , and the corresponding diagonal blocks of A are zero. It is to be noticed that the assumption $A_{[k,k]} = 0, \forall k = 1, \dots, s$ implies that theorem 3.3. and remark 3.3 of [11] for $G = GL_n(\mathbb{C})$ are immediately deducible from the main theorem of [13], starting from $\Lambda = \text{diag}(\tilde{\lambda}_1, \dots, \tilde{\lambda}_n)$ and considering the coalescence

$$(\tilde{\lambda}_1, \dots, \tilde{\lambda}_n) \mapsto (\underbrace{\lambda_1 \dots \lambda_1}_{p_1} \underbrace{\lambda_2 \dots \lambda_2}_{p_2} \dots \underbrace{\lambda_s \dots \lambda_s}_{p_s}).$$

In the present paper, no assumptions on the diagonal blocks $A_{[k,k]}$ will be made, so that our results are not deducible from or reducible to theorem 3.3. and remark 3.3 of [11] (which, as said, are obtainable from the results of [13], which are complementary to the present paper). It is also to be mentioned that the notion of isomonodromy in definition 3.2 of [11] requires that only the Stokes matrices are constant, while here we require constancy of a more stringent set of monodromy data, including the monodromy exponents and the central connection matrix.

Isomonodromic deformation equations preserving G -valued Stokes matrices were first defined in [4] for meromorphic connections on principal G -bundles, with G a complex reductive group. The leading term at an irregular singularity is assumed to be regular semisimple in the Lie algebras \mathfrak{g} . In case $G = GL(n, \mathbb{C})$ and $\mathfrak{g} = Mat(n, \mathbb{C})$, this means that its Jordan form has a single Jordan block for each eigenvalue, and in particular this implies pairwise distinct eigenvalues in the diagonalizable case. A generalization of this assumption was then given in [7], where the Jordan type is fixed (no further coalescences allowed).

Results

Our goal is a the generalization of [26], in the sense of Remark 1.1, for system (1.1).

The main results of the paper are Theorems 6.1 and 6.2. Preliminarily to them, in Theorem 3.1 some results in the weakly isomonodromic case are given: system (1.1) is weakly isomonodromic with an isomonodromic fundamental matrix solution in Levelt form at $z = 0$ if and only if the latter satisfies a Pfaffian system whose λ -components are holomorphic in $\mathbb{C} \times \mathbb{D}$. In this case, the monodromy exponents at $z = 0$ are constant. This fact is mainly based on [34].

Theorem 6.1 states that system (1.1) is strongly isomonodromic if and only if a fundamental matrix solution at $z = 0$ in Levelt form and the canonical solutions at $z = \infty$ (defined in the

¹[11] requires that Λ is diagonalizable, so one can work in the base where it is diagonal.

paper) all satisfy the integrable Pfaffian system

$$dY = \left[\left(\Lambda + \frac{A}{z} \right) dz + \sum_{j=1}^s \left(z E_{p_j} + \tilde{\omega}_j(\lambda) \right) d\lambda_j \right] Y, \quad (1.4)$$

with

$$\tilde{\omega}_j(\lambda) = \omega_j(\lambda) + \frac{\partial \mathcal{T}(\lambda)}{\partial \lambda_j} \cdot \mathcal{T}(\lambda)^{-1},$$

where $E_{p_j} := \partial \Lambda / \partial \lambda_j$ is the matrix with blocks $E_{[a,b]}^{(p_j)} = \delta_{aj} \delta_{bj} I_{p_j}$, for $a, b, j = 1, \dots, s$ (all entries are zero, except for diagonal block I_{p_j}), the matrices $\omega_j(\lambda)$ are holomorphic in $\mathbb{C} \times \mathbb{D}$ and univocally given in formula (6.2), while the block-diagonal matrix $\mathcal{T} = \mathcal{T}_1 \oplus \dots \oplus \mathcal{T}_s$ is holomorphic invertible in \mathbb{D} and reduces to Jordan form the block-diagonal part of $A(\lambda)$, namely

$$\mathcal{T}_k(\lambda)^{-1} A_{[k,k]}(\lambda) \mathcal{T}_k(\lambda) = J_k. \quad (1.5)$$

The second part of Theorem 6.1 also says that in the strong isomonodromic case, A satisfies the non-linear system

$$dA = \left[\sum_{j=1}^n \tilde{\omega}_j(\lambda) d\lambda_j, A \right]. \quad (1.6)$$

The above (1.6) predicts that the block-diagonal part

$$A_{[1,1]} \oplus \dots \oplus A_{[s,s]}$$

is constant. In particular, it has a constant Jordan form $J_1 \oplus \dots \oplus J_s$. This fact is not immediately obvious and will be proved in the paper. It implies that isomonodromy deformations with constant $\mathcal{T} = \mathcal{T}_0$ are always allowed, and in such case

$$\tilde{\omega}_j(\lambda) = \omega_j(\lambda).$$

All the other possible isomonodromic deformations are obtained with

$$\mathcal{T}(\lambda) = \mathcal{T}_0 \mathfrak{B}(\lambda),$$

for any $\mathfrak{B}(\lambda) = \mathfrak{B}_1(\lambda) \oplus \dots \oplus \mathfrak{B}_s(\lambda)$ satisfying

$$[\mathfrak{B}_k(\lambda), J_k] = 0, \quad k = 1, \dots, s.$$

Therefore, for a given A_0 at $\lambda = \lambda_0 \in \mathbb{D}$, let \mathcal{T}_0 be the block-diagonal matrix such that

$$\mathcal{T}_0^{-1} \cdot \left(A_{[1,1]}^{(0)} \oplus \dots \oplus A_{[s,s]}^{(0)} \right) \cdot \mathcal{T}_0 = J_1 \oplus \dots \oplus J_s.$$

Then, there are several possible strong isomonodromy deformations $A(\lambda)$, having the same constant block-diagonal part $A_{[1,1]}^{(0)} \oplus \dots \oplus A_{[s,s]}^{(0)}$, but different off-diagonal blocks $A_{[i,j]}(\lambda)$, $1 \leq i \neq j \leq s$, obeying equation (1.6) with the different possible choices of $\mathcal{T}(\lambda) = \mathcal{T}_0 \mathfrak{B}(\lambda)$. These different deformations are related by a λ -dependent gauge transformation, as will be explained after Theorem 6.1.

In the 3-dimensional case, when the only non-trivial case is $\Lambda = \text{diag}(\lambda_1, \lambda_2, \lambda_2)$ (up to permutations), the above freedom in the coefficients $\tilde{\omega}_j$ implies the existence of the particular isomonodromy deformation with constant A and non constant $\mathcal{T}(\lambda)$, and conversely of the deformation with non constant $A(\lambda)$ and constant \mathcal{T} . Since a gauge transformation can reduce to constant A , the 3-dimensional case is rigid. See Section 7 for details. The first non-trivial case occurs for $n = 4$ and $s = 3$. Already in simplified situations, we will show in Section 8 that the isomonodromic problem is at least as transcendental as a Painlevé equation.

Remark 1.2 (Important Remark). That strong isomonodromy deformations of (1.1) exist with constant \mathcal{T} , and that they can always be reached by a gauge transformation, does not at all mean that these deformations are trivial. Indeed, even in case \mathcal{T} is constant, $A = A(\lambda)$ is in general a *highly non trivial* and *highly transcendental* matrix function, which must satisfy the extremely difficult system (1.6) with the $\tilde{\omega}_j(\lambda) = \omega_j(\lambda)$. Only in case $n = 3$ (and $s = 1, 2$) one can reduce to the deformations with constant A , but starting from $n \geq 4$, solutions to the deformation equations (1.6) are transcendental. See Section 8 for a 4-dimensional example.

Theorem 6.2 is the converse to the second part of Theorem 6.1. It says that system (1.1) is strongly isomonodromic if A is not partially resonant (Definition 4.1) and satisfies the Frobenius integrable system (1.6), with

$$\tilde{\omega}_j(\lambda) = \omega_j(\lambda) + \mathcal{D}_j(\lambda),$$

where the $\mathcal{D}_j = \mathcal{D}_{[1,1]}^{(j)} \oplus \cdots \oplus \mathcal{D}_{[s,s]}^{(j)}$ are holomorphic block-diagonal matrices, arbitrary² except for the differential constraint

$$\partial_j \mathcal{D}_k - \partial_k \mathcal{D}_j = [\mathcal{D}_j(\lambda), \mathcal{D}_k(\lambda)],$$

which is required by the integrability of (1.6). Hence,

$$dT = \left(\sum_j \mathcal{D}_j(\lambda) d\lambda_j \right) T.$$

is integrable. It admits holomorphic fundamental matrix solutions $\mathcal{T}(\lambda) = \mathcal{T}_1 \oplus \cdots \oplus \mathcal{T}_s$ such that (1.5) holds.

The above theorems 6.1 and 6.2 generalize to the non-generic case (1.1) the strategy and the results of [26].

Remark 1.3. As a corollary, if Λ has pairwise distinct eigenvalues, (1.1) is strongly isomonodromic if and only if (1.6) holds with $\tilde{\omega}_j(\lambda) = \omega_j(\lambda) + \mathcal{D}_j(\lambda)$, with $\mathcal{D}_j(\lambda)$ diagonal satisfying $\partial_j \mathcal{D}_k - \partial_k \mathcal{D}_j = 0$ (here $j, k = 1, \dots, n$). In this case $\mathcal{T}(\lambda)$ above is *any* diagonal matrix.

An important application of the isomonodromy deformation theory here developed will be given in Section 9 for the *caustic* of a semisimple Dubrovin-Frobenius manifold M [16, 17] of dimension n . The caustic is a hypersurface $\mathcal{K} \subset M$ of codimension 1, such that the multiplication defined on the tangent bundle is nilpotent. Following [24] and [28], a generic point of \mathcal{K} has a neighbourhood in M where local coordinates $(t_1, t_2, u_3, \dots, u_n)$ are defined,

²So the case $\mathcal{D}_j = 0$ for all $j = 1, \dots, s$ is possible.

such that \mathcal{K} corresponds to $t_2 = 0$ and the vector $\partial/\partial t_2|_{t_2=0}$ is nilpotent. The flat sections of the Dubrovin deformed connection (defined in Section 9), expressed in these coordinates and restricted at the caustic, are solutions of a Pfaffian system exactly of the type (1.4) (see system (9.27)), with deformation parameters given by the coordinates (t_1, u_3, \dots, u_n) on \mathcal{K} . It has non trivial \mathcal{T} depending on the flat metric defined of M , and

$$\Lambda := \text{diag}(\underbrace{t_1, t_1}_{n-2}, \underbrace{u_3, \dots, u_n}_{\text{distinct}}).$$

Its z -component is always strongly isomonodromic in the sense described in this paper. Indeed, we will show that in case of a caustic the deformation equations (1.6), concretely realized by system (9.28), are sufficient conditions for strong isomonodromy. The results are summarized in Proposition 9.1 and Remark 9.4. Moreover, the deformation theory developed in this paper allows us to predict some properties of the caustic (see Corollary 9.1 and point 3) of Remark 9.3).

Some further remarks

The integrability conditions of a Pfaffian system (1.4) with *given* $\tilde{\omega}_j(\lambda) = \omega_j(\lambda)$ of the specific form (6.2) are the non-linear “deformation equations” (1.6) and their compatibility conditions. This is an elementary computation and is not new (see also the proof of part II of Theorem 6.1 here). For distinct eigenvalues these deformation equations are a particular case of the JMMS equations introduced in [25] (in particular section 4 and the appendix A. See also [23]), while in case of repeated eigenvalues, but no further coalescences, (1.4) and (1.6) fit into the more general deformations equations studied in [6] (see also [7]).

The purpose of the present paper is not to give deformations equations (integrability conditions) for a Pfaffian system, but to derive the Pfaffian system (1.4) as the necessary and sufficient condition for *all* the essential monodromy data of (1.1) to be constant; (1.6) is consequently the integrability condition of (1.4). Moreover, in case there are no partial resonances, we show that (1.6) is also a sufficient condition for all the essential monodromy data to be constant. This, in the spirit of Remark 1.1.

To conclude, we make two more general comments. The first is that the main difficulty to generalize [26] to non generic cases (in the sense of Remark 1.1) is to find a suitable canonical representations for a class of fundamental matrix solutions (like the Levelt form at a Fuchsian singularity and the solutions having a canonical asymptotics in Stokes sectors), and to deal with *the change of those representations* when the Jordan type of the leading matrix at an irregular singularity changes, namely some eigenvalues merge. To our knowledge, this is an extremely difficult problem, which is far from being solved. In the literature, we either find attempts to deal with coalescences of eigenvalues with a change of Jordan type, but with suitable analyticity and semisimplicity assumptions, such as in the work [13], or the Jordan types are fixed such as in [3] and in the present paper.

The second comment – which in a sense expands Remark 1.1 – is that there are two approaches in order to describe, *from the analytic viewpoint*, the isomonodromy deformations of a differential system

$$\frac{dY}{dz} = \mathfrak{A}(z, \lambda)Y, \tag{1.7}$$

where $\mathfrak{A}(z, \lambda)$ is rational in z and analytic in a domain of λ (such as \mathbb{D} here). One approach starts by proving the existence of fundamental matrix solutions of (1.7), holomorphic in the deformation parameters (under certain assumptions) in the λ -domain, and characterized by a certain canonical form, such as Levelt form at Fuchsian singularities and canonical asymptotics at irregular ones. Then, one must show that these solutions satisfy a Pfaffian system

$$dY = \omega Y, \tag{1.8}$$

with a specific $\omega(z, \lambda)$, possibly determined by $\mathfrak{A}(z, \lambda)$, if and only if the deformation is isomonodromic, namely preserves a certain class of monodromy data (such as monodromy matrices or essential monodromy data of the above mentioned solutions). This is the approach of [26] and the approach we mainly follow in our paper.

The other approach starts by assuming that we are given a Pfaffian system (1.8), satisfying the Frobenius integrability condition $d\omega = \omega \wedge \omega$, and such that the dz component of ω gives a differential system (1.7), namely

$$\omega(z, \lambda) \Big|_{\lambda \text{ fixed}} = \mathfrak{A}(z, \lambda) dz.$$

This implies that the monodromy matrices of a fundamental matrix solution $Y(z, \lambda)$ of (1.8) are constant, so that system (1.7) is weakly isomonodromic. Then, this approach proceeds by showing if, depending on the specific $\omega(z, \lambda)$, the Pfaffian system admits fundamental matrix solutions with a canonical structure, whose corresponding essential monodromy data are constant. This is, for example, the approach of [8, 9, 10, 34].

Acknowledgments. I thank P. Boalch for several useful remarks and for pointing out some references. The author is member of the European Union's H2020 research and innovation programme under the Marie Skłodowska-Curie grant No. 778010 *IPaDEGAN*. He is also a member of the Project 'Mathematical Methods in Non Linear Physics' (MMNLP), Commissione Scientifica Nazionale 4 - Fisica Teorica (CNS4) of the Istituto Nazionale di Fisica Nucleare (INFN).

2 Preliminaries

It is a standard result [33, 1] that, for each fixed λ , system (1.1) admits a fundamental solution with *Levelt form*³ at $z = 0$:

$$Y^{(0)}(z, \lambda) := G^{(0)}(\lambda) \widehat{Y}^{(0)}(z, \lambda) z^{D^{(0)}} z^{L^{(0)}(\lambda)}, \tag{2.1}$$

$$J^{(0)} = D^{(0)} + S^{(0)}, \quad L^{(0)} = S^{(0)} + R^{(0)}, \tag{2.2}$$

where:

$$\widehat{Y}^{(0)}(z, \lambda) = I + \sum_{k=1}^{\infty} F_k^{(0)}(\lambda) z^k \quad \text{is convergent for finite } |z|; \tag{2.3}$$

³This is an improper Levelt form, obtained by a permutation $Y \mapsto YP$, P a suitable permutation matrix, from a proper Levelt form (5.8) of Section 5.

the matrix $J^{(0)}(\lambda)$ is a Jordan form of $A(\lambda)$; the eigenvalues $\mu_1(\lambda), \dots, \mu_n(\lambda)$ of $A(\lambda)$ are uniquely decomposed as

$$\mu_j(\lambda) = d_j^{(0)} + \rho_j^{(0)}(\lambda), \quad \text{with } 0 \leq \Re \rho_j^{(0)} < 1 \text{ and } d_j^{(0)} \in \mathbb{Z};$$

and

$$\text{diag}(S^{(0)}) = \text{diag}(\rho_1^{(0)}, \dots, \rho_n^{(0)}), \quad D^{(0)} = \text{diag}(d_1^{(0)}, \dots, d_n^{(0)}).$$

The matrix $R^{(0)}$ is nilpotent, with entries

$$(R^{(0)}(\lambda))_{ij} \neq 0 \text{ only if } \mu_i(\lambda) - \mu_j(\lambda) \in \mathbb{N} \setminus \{0\}.$$

The invertible matrix $G^{(0)}$ puts A in Jordan form. In general, the dependence of $Y^{(0)}$ on λ is not holomorphic in \mathbb{D} .

Assumption 2. $A(\lambda)$ is holomorphically similar to a Jordan form $J^{(0)}(\lambda)$ in \mathbb{D} , namely there is $G^{(0)} = G^{(0)}(\lambda)$ holomorphically invertible in \mathbb{D} such that

$$J^{(0)}(\lambda) = G^{(0)}(\lambda)^{-1} A(\lambda) G^{(0)}(\lambda).$$

The matrix $A(\lambda)$ is said to be *resonant* at $\lambda \in \mathbb{D}$ if there exist $i \neq j \in \{1, \dots, n\}$ such that $\mu_i(\lambda) - \mu_j(\lambda) \in \mathbb{Z} \setminus \{0\}$. If the eigenvalues do not depend on λ in \mathbb{D} , we simply say that A is resonant.

If there are no resonances, Assumption 2 guarantees that $Y^{(0)}(z, \lambda)$ can be taken holomorphic on $\mathcal{R} \times \mathbb{D}$. Otherwise, in addition to Assumption 2 we need to require that if $\mu_i(\lambda) - \mu_j(\lambda) = \ell_{ij} \in \mathbb{Z} \setminus \{0\}$ for some λ , then the *resonance persists all over* \mathbb{D} , namely

$$\mu_i(\lambda) - \mu_j(\lambda) = \ell_{ij} \in \mathbb{Z} \setminus \{0\} \quad \forall \lambda \in \mathbb{D}. \quad (2.4)$$

Then, holomorphy of (2.1) follows from its standard formal computation (see [33]). The reason for (2.4) is that if it does not hold, then $R^{(0)}(\lambda)$ may have an extremely wild behaviour in λ .

3 Weak isomonodromic deformations

Lemma 3.1 (Isospectrality). *Let $A(\lambda)$ be holomorphic on \mathbb{D} . If (1.1) has for each $\lambda \in \mathbb{D}$ a fundamental matrix solution $Y(z; \lambda)$ whose monodromy*

$$Y(z; \lambda) \longmapsto Y(ze^{2\pi i}; \lambda) = Y(z; \lambda)M,$$

is the same for all $\lambda \in \mathbb{D}$ (i.e. the monodromy matrix M is constant), then the eigenvalues of $A(\lambda)$ are constant on \mathbb{D} . In particular, (2.4) holds in case of resonances. If moreover Assumption 2 holds, then a Levelt form $Y^{(0)}$ is holomorphic on $\mathcal{R} \times \mathbb{D}$.

Notice that in Lemma 3.1 it is not assumed that $Y(z; \lambda)$ depends holomorphically on λ .

Proof. There exists for each λ an invertible connection matrix $C(\lambda)$ such that

$$Y(z; \lambda) = Y^{(0)}(z, \lambda)C(\lambda) = G^{(0)}(\lambda)\widehat{Y}^{(0)}(z, \lambda)z^{D^{(0)}(\lambda)}z^{L^{(0)}(\lambda)}C(\lambda).$$

By assumption, $M = C(\lambda)^{-1}e^{2\pi iL^{(0)}(\lambda)}C(\lambda)$ does not depend on λ , so the eigenvalues $\rho_j^{(0)}$ of $L^{(0)}$ are constant. Since $A(\lambda)$ is holomorphic, its eigenvalues $\mu_j(\lambda) = d_j^{(0)} + \rho_j^{(0)}$ are continuous. It follows that both the integers $d_j^{(0)}$ and the eigenvalues μ_j are constant. Clearly, (2.4) holds, so that holomorphy follows from the formal computation (see [33]) of (2.1) and (2.3). \square

Definition 3.1. *System (1.1) is weakly isomonodromic in \mathbb{D} if there exists a fundamental matrix solution $Y^{\text{hol}}(z, \lambda)$ depending holomorphically on $(z, \lambda) \in \mathcal{R} \times \mathbb{D}$, with λ -independent monodromy matrix M , defined by*

$$Y^{\text{hol}}(z, \lambda) \mapsto Y^{\text{hol}}(z, \lambda)M, \quad z \mapsto ze^{2\pi i}.$$

Proposition 3.1. *Let $A(\lambda)$ be holomorphic in \mathbb{D} . System (1.1) is weakly isomonodromic in \mathbb{D} if and only if it is the z -component of an integrable Pfaffian system*

$$dY = \omega(z, \lambda)Y, \quad \omega(z, \lambda) = \left(\Lambda + \frac{A}{z} \right) dz + \sum_{j=1}^s \omega_j(z, \lambda)d\lambda_j. \quad (3.1)$$

with $\omega(z, \lambda)$ holomorphic in $\overline{\mathbb{C}} \setminus \{0, \infty\} \times \mathbb{D}$, satisfying the integrability condition

$$d\omega = \omega \wedge \omega.$$

The proof is standard.

Theorem 3.1. *Let $A(\lambda)$ be holomorphic in \mathbb{D} . System (1.1) is weakly isomonodromic in \mathbb{D} with holomorphic fundamental matrix solution Y^{hol} coinciding with a fundamental solution in Levelt form $Y^{(0)}$, if and only*

the coefficients $\omega_j(z, \lambda)$ are holomorphic in $\mathbb{C} \times \mathbb{D}$.

In this case, the following facts hold.

- $D^{(0)}$ and $L^{(0)}$ are constant, or equivalently $J^{(0)}$ and $R^{(0)}$ are constant.
- $A(\lambda)$ is holomorphically similar to $J^{(0)}$ through a fundamental matrix solution $G^{(0)}(\lambda)$ of

$$dG = \sum_{j=1}^s \omega_j(0, \lambda)d\lambda_j G.$$

The matrices $\omega_j(z, \lambda)$ in Theorem 3.1 may have isolated singularity in $z = \infty$. The requirement $Y^{\text{hol}} = Y^{(0)}$ is equivalent to the requirement that $Y^{\text{hol}} = Y^{(0)}C$, for C constant invertible matrix.

Assumption 2 is not explicitly written in the statement of Theorem 3.1. If (1.1) is weakly isomonodromic in \mathbb{D} and we assume that there is a holomorphic fundamental matrix solution in Levelt form $Y^{(0)}$, then Assumption 2 is a necessary condition, so it is automatically assumed by requiring that $Y^{(0)}$ is holomorphic. Conversely, if the coefficients $\omega_j(z, \lambda)$ are holomorphic in $\mathbb{C} \times \mathbb{D}$, then it follows from Proposition 3.2 below that $A(\lambda)$ is holomorphically similar to $J^{(0)}$, namely Assumption 2 is satisfied.

Theorem 3.1 holds also for

$$\begin{aligned} \frac{dY}{dz} &= \mathfrak{A}(z, t)Y, & \mathfrak{A}(z, t) &:= \Lambda + \frac{A(t)}{z-a} \\ t &:= (\lambda, a), & \lambda &= (\lambda_1, \dots, \lambda_s) \in \mathbb{D} \subset \mathbb{C}^s, \quad a \in \mathbb{C}. \end{aligned} \quad (3.2)$$

where one deformation parameter is the pole $z = a$. In this case, (3.1) is replaced by

$$\omega(z, t) = \mathfrak{A}(z, t)dz + \sum_{j=1}^s \omega_j(z, t)d\lambda_j + \left(\omega_0(z, t) - \frac{A(t)}{z-a} \right) da. \quad (3.3)$$

The coefficients $\omega_j(z, \lambda)$ and $\omega_0(z, t)$ are holomorphic in $\mathbb{C} \times \mathbb{D}$. We stress that the above holds for any $A(t)$, including the non-diagonalizable and resonant cases. Moreover, $G^{(0)}(t)$ is a fundamental matrix solution of

$$dG = \left(\sum_{j=1}^s \omega_j(a, t)d\lambda_j + \omega_0(a, t)da + \varphi(t)da \right) G, \quad (3.4)$$

with

$$\varphi = G^{(0)} \left(F_1^{(0)} + [F_1^{(0)}, J^{(0)}] + R_1^{(0)} \right) (G^{(0)})^{-1}. \quad (3.5)$$

Here, $F_1^{(0)}$ appears in the Taylor expansion (2.3), $J^{(0)}$ in (2.2), and $R_1^{(0)}$ is the first term in the decomposition of $R^{(0)} = \sum_{\ell=1}^m R_\ell^{(0)}$, where

$$(R_\ell^{(0)})_{ij} \neq 0 \text{ only if } \mu_i - \mu_j = \ell \in \mathbb{N} \setminus \{0\}.$$

Notice that (3.4) is linear, because $F_1^{(0)}$ and $R_1^{(0)}$ are obtained by the standard formal computation yielding (2.1), which is done using the differential system $d\tilde{Y}/dz = (G^{(0)})^{-1}\mathfrak{A}(z, t)G^{(0)}\tilde{Y}$, after the gauge transformation $Y = G^{(0)}\tilde{Y}$. Explicit computation shows that $G^{(0)}$ cancels in (3.5), namely φ is only determined by $\mathfrak{A}(z, t)$.

Expressions (3.3), (3.4), (3.5) can be obtained following the same steps of the proofs of Propositions 3.2 and 3.3 below. Just notice that for example in (3.8) one has

$$\omega = d(G^{(0)}\hat{Y}^{(0)}) \cdot (G^{(0)}\hat{Y}^{(0)})^{-1} + G^{(0)}\hat{Y}^{(0)} \frac{D^{(0)} + (z-a)^{D^{(0)}}L^{(0)}(z-a)^{-D^{(0)}}}{(z-a)} (G^{(0)}\hat{Y}^{(0)})^{-1} dz$$

$$-G^{(0)}\widehat{Y}^{(0)}\frac{D^{(0)}+(z-a)^{D^{(0)}}L^{(0)}(z-a)^{-D^{(0)}}}{z-a}(G^{(0)}\widehat{Y}^{(0)})^{-1}da.$$

Then use the definitions of the monodromy exponents (2.2), which imply that

$$D^{(0)}+(z-a)^{D^{(0)}}L^{(0)}(z-a)^{-D^{(0)}}=J^{(0)}+\sum_{\ell=1}^m R_{\ell}^{(0)}(z-a)^{\ell}.$$

3.1 Proof of Theorem 3.1

Theorem 3.1 follows from Propositions 3.2 and 3.3 below. These propositions apply also to system (3.2), with form (3.3) and (3.4).

By its definition $D^{(0)}$ is locally constant on subsets of \mathbb{D} . It may have jump discontinuities on \mathbb{D} , so that $dD^{(0)}$ is not well defined. We will sometimes write $dD^{(0)}=0$ with abuse of notation when we want to indicate that $D^{(0)}$ is constant on the whole \mathbb{D} .

Proposition 3.2. *Let $A(\lambda)$ be holomorphic in \mathbb{D} and let (1.1) be the z -component of an integrable Pfaffian system (3.1) whose coefficients $\omega_j(z, \lambda)$ are holomorphic in $\mathbb{C} \times \mathbb{D}$.*

Then, there is a fundamental matrix solution $Y^{(0)}(z, \lambda)$ of (3.1) in Levelt form (2.1), with

$$dD^{(0)}=dL^{(0)}=0, \quad \text{or equivalently} \quad dJ^{(0)}=dR^{(0)}=0 \quad \text{on } \mathbb{D}. \quad (3.6)$$

Moreover, the matrix $A(\lambda)$ is holomorphically similar to $J^{(0)}$ through a fundamental matrix solution $G^{(0)}(\lambda)$ of

$$dG = \sum_{j=1}^s \omega_j(0, \lambda) d\lambda_j G.$$

Proof. Proposition 3.2 is a particular case of the main results of [34] on fundamental matrix solutions of Pfaffian systems at a logarithmic (Fuchsian) singularity. \square

The converse of the above is the following

Proposition 3.3. *Let $A(\lambda)$ be holomorphic in \mathbb{D} and let Assumption 2 hold.*

a) Suppose that (1.1) is weakly isomonodromic. If there is a fundamental matrix solution of (3.1) in Levelt form $Y^{(0)}(z, \lambda)$, then the coefficients $\omega_j(z, \lambda)$ of (3.1) are holomorphic in $\mathbb{C} \times \mathbb{D}$ and (3.6) holds.

b) Conversely, if system (1.1) has a fundamental solution $Y^{(0)}(z, \lambda)$ in Levelt form (2.1) such that (3.6) holds, then the system is weakly isomonodromic, and the corresponding Pfaffian system (3.1) has coefficients $\omega_j(z, \lambda)$ holomorphic in $\mathbb{C} \times \mathbb{D}$.

In both cases a) and b), $G^{(0)}(\lambda)$ is a fundamental matrix solution of

$$dG = \sum_{j=1}^s \omega_j(0, \lambda) d\lambda_j G.$$

Proof. a) Being (1.1) weakly isomonodromic, there is an isomonodromic $Y^{\text{hol}}(z, \lambda)$, with constant monodromy matrix M , satisfying (3.1). By Lemma 3.1, a solution $Y^{(0)}(z, \lambda)$ of (1.1) exists holomorphic in $\mathcal{R} \times \mathbb{D}$, with

$$D^{(0)} \text{ constant.}$$

By the assumption in a), $Y^{(0)}$ also satisfies (3.1). Being solutions of (1.1), Y^{hol} and $Y^{(0)}$ are related by a holomorphic connection matrix $C(\lambda)$:

$$Y^{\text{hol}}(z, \lambda) = Y^{(0)}(z, \lambda)C(\lambda).$$

Since both $dY^{\text{hol}} = \omega Y^{\text{hol}}$ and $dY^{(0)} = \omega Y^{(0)}$ hold, then

$$dC = 0.$$

Let us rewrite

$$\begin{aligned} Y^{\text{hol}}(z, \lambda) &= Y^{(0)}(z, \lambda)C = G^{(0)}(\lambda)\widehat{Y}^{(0)}(z, \lambda)z^{D^{(0)}}Cz^{\mathcal{L}^{(0)}}, \\ \mathcal{L}^{(0)}(\lambda) &:= C^{-1}L^{(0)}(\lambda)C. \end{aligned}$$

Since $dM = d(\exp\{2\pi i\mathcal{L}^{(0)}\}) = 0$ by assumption, we have $d\mathcal{L}^{(0)} = 0$, and then

$$dL^{(0)} = 0.$$

Therefore, we find

$$\omega = dY^{\text{hol}} \cdot (Y^{\text{hol}})^{-1} = d(Y^{(0)} \cdot (Y^{(0)})^{-1}) \quad (3.7)$$

$$= d(G^{(0)}\widehat{Y}^{(0)}) \cdot (G^{(0)}\widehat{Y}^{(0)})^{-1} + G^{(0)}\widehat{Y}^{(0)} \frac{D^{(0)} + z^{D^{(0)}}L^{(0)}z^{-D^{(0)}}}{z} (G^{(0)}\widehat{Y}^{(0)})^{-1} dz \quad (3.8)$$

Now, the definition of $D^{(0)}$ and $L^{(0)}$ implies that $D^{(0)} + z^{D^{(0)}}L^{(0)}z^{-D^{(0)}}$ is holomorphic at $z = 0$ and

$$\lim_{z \rightarrow 0} (D^{(0)} + z^{D^{(0)}}L^{(0)}z^{-D^{(0)}}) = J^{(0)}, \quad (3.9)$$

so that

$$\omega = dG^{(0)} \cdot (G^{(0)})^{-1} + \text{reg}(z, \lambda) + \left(\frac{A(\lambda)}{z} + \text{reg}_1(z, \lambda) \right) dz.$$

Here

$$\text{reg}(z, \lambda) = O(z), \text{ for } z \rightarrow 0$$

is a 1-form in dz and $d\lambda_1, \dots, d\lambda_s$, holomorphic in $\mathbb{C} \times \mathbb{D}$. Moreover, $\text{reg}_1(z, \lambda)$ is a holomorphic matrix in $\mathbb{C} \times \mathbb{D}$ with behaviour

$$\text{reg}_1(z, \lambda) = O(1), \text{ for } z \rightarrow 0.$$

We conclude that

$$\omega = \sum_{j=1}^s \omega_j(z, \lambda) d\lambda_j + \left(\frac{A(\lambda)}{z} + \text{reg}_1(z, \lambda) \right) dz,$$

where both $\text{reg}_1(z, \lambda)$ and the matrices $\omega_j(z, \lambda)$ are holomorphic in $\mathbb{C} \times \mathbb{D}$, of order $O(1)$ for $z \rightarrow 0$. In particular,

$$dG^{(0)} \cdot (G^{(0)})^{-1} = \sum_{j=1}^s \omega_j(0, z) d\lambda_j.$$

b) Suppose that $dD^{(0)} = dL^{(0)} = 0$, so that for $z \mapsto ze^{2\pi i}$ the monodromy $Y^{(0)} \mapsto Y^{(0)} e^{2\pi i L^{(0)}}$ is constant. This implies that Lemma 3.1 holds, so that $Y^{(0)}(z, \lambda)$ is holomorphic in $\mathcal{R} \times \mathbb{D}$. We prove that $Y^{(0)}$ satisfies a Pfaffian system. We define

$$\omega(z, \lambda) := dY^{(0)}(z, \lambda) \cdot (Y^{(0)}(z, \lambda))^{-1}.$$

This is single valued with respect to z , because the monodromy of $Y^{(0)}$ is constant. The structure of $\omega(z, \lambda) = \omega_0(z, \lambda) dz + \sum_{j=1}^s \omega_j(z, \lambda) d\lambda_j$ is computable from (2.1):

$$\begin{aligned} dY^{(0)} \cdot (Y^{(0)})^{-1} &= \\ &= dG^{(0)} \cdot G^{(0)} + G^{(0)} d\hat{Y}^{(0)} \cdot (G^{(0)} \hat{Y}^{(0)})^{-1} + G^{(0)} \hat{Y}^{(0)} \frac{D^{(0)} + z^{D^{(0)}} L^{(0)} z^{-D^{(0)}}}{z} (G^{(0)} \hat{Y}^{(0)})^{-1} \\ &= dG^{(0)} \cdot G^{(0)} + \text{reg}(z, \lambda) + \left(\frac{A_1}{z} + \text{reg}_1(z, \lambda) \right) dz. \end{aligned}$$

In the last step, we have used (2.3) and (3.9). Here $\text{reg}(z, \lambda)$ stands for a matrix valued 1-form in the $d\lambda_j$'s, holomorphic in $\mathbb{C} \times \mathbb{D}$, and of order $O(z) \rightarrow 0$ as $z \rightarrow 0$, while $\text{reg}_1(z, \lambda)$ is a matrix holomorphic in $\mathbb{C} \times \mathbb{D}$ of order $O(1)$ as $z \rightarrow 0$ (and we know that it must be Λ). In conclusion, we have found that

$$\sum_{j=1}^s \omega_j(z, \lambda) d\lambda_j := dG^{(0)} \cdot G^{(0)} + \text{reg}(z, \lambda),$$

and in particular $dG^{(0)} \cdot (G^{(0)})^{-1} = \sum_{j=1}^s \omega_j(0, \lambda) d\lambda_j$. \square

4 Canonical solutions of (1.1) at $z = \infty$

Let us partition A in blocks $A_{[i,j]}$, $i, j = 1, \dots, s$, of dimension $p_i \times p_j$, inherited from Λ . Let

$$\mathcal{T}(\lambda) = \mathcal{T}_1(\lambda) \oplus \dots \oplus \mathcal{T}_s(\lambda),$$

be a block diagonal matrix such that

$$\mathcal{T}_k(\lambda)^{-1} A_{[k,k]}(\lambda) \mathcal{T}_k(\lambda) = J_k(\lambda) \quad \text{Jordan form,} \quad k = 1, \dots, s. \quad (4.1)$$

It has structure $\mathcal{T}_k(\lambda) = \mathcal{T}_k^0(\lambda)\mathfrak{B}_k(\lambda)$, where $\mathcal{T}_k^0(\lambda)$ is a chosen matrix satisfying (4.1) and $\mathfrak{B}_k(\lambda)$ is any matrix such that $[\mathfrak{B}_k, J_k] = 0$.

Assumption 3. $A_{[1,1]}(\lambda) \oplus \cdots \oplus A_{[s,s]}(\lambda)$ is holomorphically reducible to Jordan form

$$J(\lambda) = J_1(\lambda) \oplus \cdots \oplus J_s(\lambda).$$

This means that each $\mathcal{T}_k(\lambda)$ is holomorphic on \mathbb{D} , and so is each $J_k(\lambda)$.

We can arrange each J_k into $h_k \leq p_k$ Jordan blocks $J_1^{(k)}, \dots, J_{h_k}^{(k)}$

$$J_k = J_1^{(k)} \oplus \cdots \oplus J_{h_k}^{(k)}. \quad (4.2)$$

Each block $J_j^{(k)}$, $1 \leq j \leq h_k$, has dimension $r_j \times r_j$, with $r_j \geq 1$, $r_1 + \cdots + r_{h_k} = p_k$. Each $J_j^{(k)}$ has only one eigenvalue $\mu_j^{(k)}$, with structure,

$$J_j^{(k)}(\lambda) = \mu_j^{(k)}(\lambda)I_{r_j} + H_{r_j}, \quad I_{r_j} = r_j \times r_j \text{ identity matrix,}$$

$$H_{r_j} = \begin{cases} 0 & \text{if } r_j = 1, \\ \begin{bmatrix} 0 & 1 & & \\ & 0 & 1 & \\ & & \ddots & \ddots \\ & & & 0 & 1 \\ & & & & 0 \end{bmatrix} & \text{if } r_j \geq 2. \end{cases}$$

Note that $\mu_1^{(k)}, \dots, \mu_{h_k}^{(k)}$ are not necessarily distinct. The decomposition $\mu_j^{(k)} = d_j^{(k)} + \rho_j^{(k)}$, with $d_j^{(k)} \in \mathbb{Z}$ and $0 \leq \Re \rho_j^{(k)} < 1$, induces the decomposition

$$J_k = D_k + S_k, \quad k = 1, \dots, s. \quad (4.3)$$

where D_k is diagonal with eigenvalues $d_j^{(k)}$ and S_k is Jordan with eigenvalues $\rho_j^{(k)}$. We let

$$D := D_1 \oplus \cdots \oplus D_s, \quad S := S_1 \oplus \cdots \oplus S_s \quad \text{so that } J = D + S.$$

If Assumption 3 holds, the gauge

$$Y(z, \lambda) = \mathcal{T}(\lambda)\widehat{X}(z, \lambda)$$

transforms system (1.1) into

$$\frac{d\widehat{X}}{dz} = \left(\Lambda + \frac{\mathcal{A}(\lambda)}{z} \right) \widehat{X}, \quad \mathcal{A} := \mathcal{T}^{-1}A\mathcal{T} \equiv \begin{pmatrix} J_1 & \mathcal{A}_{[1,2]} & \cdots & \mathcal{A}_{[1,s]} \\ \mathcal{A}_{[2,1]} & J_2 & \cdots & \mathcal{A}_{[2,s]} \\ \vdots & \vdots & \ddots & \vdots \\ \mathcal{A}_{[s,1]} & \mathcal{A}_{[s,2]} & \cdots & J_s \end{pmatrix} \quad (4.4)$$

We can then apply to (4.4) the computations of section 4.1 of [13], which allow to find formal solutions of (1.1) depending holomorphically on $\lambda \in \mathbb{D}$, with structure

$$Y_F(z, \lambda) = \mathcal{T}(\lambda) \left(I + \sum_{j=1}^{\infty} F_j(\lambda) z^{-j} \right) z^{D(\lambda)} z^{L(\lambda)} e^{\Lambda z}. \quad (4.5)$$

Here

$$L := L_1 \oplus \cdots \oplus L_s, \quad L_k := S_k + R_k, \quad R_k \text{ is nilpotent.} \quad (4.6)$$

Each R_k has possibly non zero blocks

$$[R_k]_{\text{block } a,b} \neq 0 \quad \text{only if} \quad \mu_b^{(k)}(\lambda) - \mu_a^{(k)}(\lambda) = \ell_{ba} \in \mathbb{N} \setminus \{0\}, \quad a \neq b = 1, \dots, h_k. \quad (4.7)$$

The diagonal matrix $D(\lambda)$ is locally constant, from its very definition, and may have discrete jumps as λ varies in \mathbb{D} . The computation of the $F_k(\lambda)$ and $R = R_1 \oplus \cdots \oplus R_s$ follows exactly the procedure of proposition 4.1 of [13].

In case there are no resonances in $A_{[k,k]}(\lambda)$, then $R_k(\lambda) = 0$. If there are no resonances in all the blocks $A_{[k,k]}$, $\forall k = 1, \dots, s$, then

$$R = 0 \quad \implies \quad L(\lambda) = S(\lambda),$$

and

$$Y_F(z, \lambda) = \mathcal{T}(\lambda) \left(I + \sum_{j=1}^{\infty} F_j(\lambda) z^{-j} \right) z^{D(\lambda)} z^{S(\lambda)} e^{\Lambda z} = \mathcal{T}(\lambda) \left(I + \sum_{j=1}^{\infty} F_j(\lambda) z^{-j} \right) z^{J(\lambda)} e^{\Lambda z}$$

Then, by Assumption 3, $Y_F(z, \lambda)$ depends holomorphically on λ .

In case of resonance of some $A_{[k,k]}(\lambda)$, a sufficient condition for the $F_j(\lambda)$'s and $L(\lambda)$ to depend holomorphically on λ is that when it happens that $\mu_b^{(k)}(\lambda) - \mu_a^{(k)}(\lambda) = \ell_{ba} \in \mathbb{N} \setminus \{0\}$ for some value of λ , then the resonance persists all over \mathbb{D} , namely

$$\mu_b^{(k)}(\lambda) - \mu_a^{(k)}(\lambda) = \ell_{ba} \in \mathbb{N} \setminus \{0\} \quad \forall \lambda \in \mathbb{D}. \quad (4.8)$$

In this case, being D locally constant in \mathbb{D} , $Y_F(z, \lambda)$ locally depends holomorphically on λ .

Definition 4.1. *In the terminology introduced in [29], if there exists k such that $A_{[k,k]}$ is resonant, we say that A has a **partial resonance**.*

A formal solution (4.5) with given \mathcal{T} , L , D and Λ is uniquely determined only if all the matrices $A_{[1,1]}(\lambda), \dots, A_{[s,s]}(\lambda)$ are non-resonant (see corollary 4.1 of [13]).

Remark 4.1. In case $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$ has pairwise distinct eigenvalues, then

$$Y_F(z, \lambda) = \mathcal{T}(\lambda) \left(I + \sum_{j=1}^{\infty} F_j(\lambda) z^{-j} \right) z^{\text{diag}(A(\lambda))} e^{\Lambda z},$$

and $\mathcal{T}(\lambda)$ is an arbitrary invertible diagonal matrix. One can choose it to be the identity matrix.

Stokes Matrices

Consider an admissible direction τ as in Assumption 1 and the following λ -independent sectors in \mathcal{R} of central angular opening $\pi + 2\delta$:

$$\mathcal{S}_\nu : \quad (\tau + (\nu - 1)\pi) - \delta < \arg z < (\tau + \nu\pi) + \delta, \quad \nu \in \mathbb{Z}, \quad \delta > 0.$$

If Assumption 1 holds, there is a sufficiently small δ such that $\mathcal{S}_\nu \cap \mathcal{S}_{\nu+1}$ does not contain Stokes rays as λ varies in \mathbb{D} . From [30], we know that to a prefixed formal solution (4.5) there correspond actual solutions satisfying

$$Y_\nu(z, \lambda) = \mathcal{T}(\lambda) \widehat{Y}_\nu(z, \lambda) z^{D(\lambda)} z^{L(\lambda)} e^{\Lambda z}, \quad (4.9)$$

$$\widehat{Y}_\nu(z, \lambda) \sim I + \sum_{j=1}^{\infty} F_j(\lambda) z^{-j}, \quad z \rightarrow \infty \text{ in } \mathcal{S}_\nu. \quad (4.10)$$

For short, we will improperly write

$$Y_\nu(z, \lambda) \sim Y_F(z, \lambda), \quad z \rightarrow \infty \text{ in } \mathcal{S}_\nu.$$

They are uniquely determined by the above asymptotic behaviour (as proved in theorem 6.2 of [13]). When Assumption 3 and (4.8) hold, they are holomorphic in $\mathcal{R} \times \mathbb{D}$. In this case, the holomorphic **Stokes matrices** $\mathbb{S}_\nu(\lambda)$ are defined by

$$Y_{\nu+1}(z, \lambda) = Y_\nu(z, \lambda) \mathbb{S}_\nu(\lambda).$$

5 More on the Levelt form

This technical section can be skipped at first reading. It introduces details needed especially in the proof in the Appendix. The reader not interested in the Appendix may just read the last sentence of this section, starting with “In conclusion,...”.

Consider a $N \times N$ system $Y' = \mathfrak{A}(z)Y$, such that $\mathfrak{A}(z)$ has a Fuchsian singularity in $z = a$, for $a \in \mathbb{C}$, or in $z = \infty$. The residue matrix of $\mathfrak{A}(z)$ at $z = a$ (or $z = \infty$) has a Jordan form

$$J = J_1 \oplus \cdots \oplus J_r,$$

with

$$J_j = \mu_j I_{m_j} + H_{m_j}, \quad m_1 + \dots + m_r = N,$$

$$H_{m_j} = \begin{cases} 0 & \text{if } m_j = 1, \\ \begin{bmatrix} 0 & 1 & & & \\ & 0 & 1 & & \\ & & \ddots & \ddots & \\ & & & 0 & 1 \\ & & & & 0 \end{bmatrix} & \text{if } m_j \geq 2. \end{cases}$$

We can arrange the Jordan form so that the eigenvalues μ_1, \dots, μ_r of J have real parts forming a non increasing sequence if $z = a$ is the singularity:

$$\Re \mu_1 \geq \Re \mu_2 \geq \cdots \geq \Re \mu_r; \quad (5.1)$$

or a non decreasing sequence in case $z = \infty$ is the singularity:

$$\Re \mu_1 \leq \Re \mu_2 \leq \cdots \leq \Re \mu_r. \quad (5.2)$$

We also write $\mu_j = \rho_j + d_j$, with $0 \leq \Re \rho_i < 1$ and $d_j \in \mathbb{Z}$, and

$$J = D + S,$$

where D is the diagonal matrix of integers d_j .

The differential system can be reduced to normal form by a standard procedure [33], and this allows to find a fundamental matrix solution in Levelt form

$$Y(z) = \mathcal{G}(\zeta)\zeta^D\zeta^L, \quad (5.3)$$

where $\zeta = z - a$ if a is the singularity, or $\zeta = z$ if ∞ is the singularity. Here, $\mathcal{G}(\zeta)$ is holomorphic at $z = a$ (or at $z = \infty$). In case the matrix coefficient $\mathfrak{A}(z)$ is holomorphic only in a sector centered at the singularity, and admits there an asymptotic expansion, then $\mathcal{G}(\zeta)$ is holomorphic in that sector, with asymptotic expansion there [33]. Moreover, the monodromy exponent L is

$$L = S + R,$$

where the matrix R is nilpotent and obtained by the formal computation of the normal form.

Consider the block partition of R inherited from J . For the singularity $z = a$, it possibly has a non-trivial block in position (i, j) , with $1 \leq i \neq j \leq r$, if $\mu_i - \mu_j = d_i - d_j \geq 1$ is integer. For the singularity $z = \infty$, R possibly has a non-trivial block in position (j, i) if $\mu_i - \mu_j = d_i - d_j \geq 1$ is integer. It follows from the ordering (5.1) or (5.2) that R only has possibly non zero blocks in the upper triangular part of its block partition (R is upper triangular if J is diagonal). The diagonal blocks of R are zero (the diagonal is zero if J is diagonal).

Examples. The solution (2.1) is an example for $a = 0$. The solutions $Y_\nu(z, \lambda)$ in (4.9) contains the matrix factor $\mathcal{T}(\lambda)\widehat{Y}_\nu(z, \lambda)z^{D(\lambda)}z^{L(\lambda)}$, which is an example with $\zeta = z$ and $\mathcal{G}(\zeta)$ holomorphic at $z = \infty$ in a sector \mathcal{S}_ν : indeed, it is a fundamental solution in Levelt form for the Fuchsian system (4.1) at $z = \infty$ of the paper [13].

Notice once more that, with the given ordering (5.1) or (5.2), for $1 \leq i < j \leq r$ we have $\mu_i \neq \mu_j$ and $\rho_i = \rho_j$ whenever $\mu_i - \mu_j \equiv d_i - d_j \neq 0$ is a non-zero integer, and correspondingly R possibly has a non-zero block in position (i, j) . Therefore, possibly acting by a permutation $L \mapsto P^{-1}LP$ if necessary (which means changing $Y \mapsto YP$ by a permutation matrix P), we can always do the above construction in such a way that L admits another partition into blocks

$$L = L_1 \oplus \cdots \oplus L_\ell, \quad \text{with } \ell \leq r,$$

where each block L_q is upper triangular, it has only one eigenvalue σ_q equal to some $\rho_i = \rho_j = \dots$ from the set $\{\rho_1, \dots, \rho_r\}$, satisfying $0 \leq \Re\sigma_q < 1$, and $\sigma_p \neq \sigma_q$ for $1 \leq p \neq q \leq \ell$, and the corresponding diagonal matrix D of integer parts of the eigenvalues of J is split into blocks $D = D_1 \oplus \cdots \oplus D_\ell$, with

$$D_q = \text{diag}(d_{q,1}, d_{q,2}, \dots), \quad q = 1, \dots, \ell,$$

where for each q the integers form a non-increasing finite sequence

$$d_{q,1} \geq d_{q,2} \geq \dots$$

in case the singularity is $z = a$; or a non-decreasing finite sequence

$$d_{q,1} \leq d_{q,2} \leq \dots$$

in case $z = \infty$ is the singularity. We can therefore rewrite

$$L = S + R, \quad \text{with} \quad S = S_1 \oplus \cdots \oplus S_\ell, \quad R = R_1 \oplus \cdots \oplus R_\ell.$$

Each block L_q , $1 \leq q \leq \ell$, consists of sub-blocks, according to the structure (for some integer k_q):

$$L_q = S_q + R_q, \quad S_q = \begin{pmatrix} S_1^{(q)} & & & \\ & S_2^{(q)} & & \\ & & \ddots & \\ & & & S_{k_q}^{(q)} \end{pmatrix}, \quad R_q = \begin{pmatrix} \mathbf{0} & * & * & * \\ & \mathbf{0} & * & * \\ & & \ddots & * \\ & & & \mathbf{0} \end{pmatrix}, \quad (5.4)$$

where each matrix $S_i^{(q)}$ is a Jordan matrix with the same eigenvalue σ_q on the diagonal and 1's on the second upper diagonal:

$$S_i^{(q)} = \begin{pmatrix} \sigma_q & 1 & 0 & 0 \\ & \sigma_q & 1 & 0 \\ & & \ddots & 1 \\ & & & \sigma_q \end{pmatrix}, \quad i = 1, 2, \dots, k_q;$$

while in R_q the $\mathbf{0}$ are zero diagonal blocks (corresponding to the blocks $S_i^{(q)}$), and each $*$ is an off-diagonal block which is possibly non zero (now the block partition of R_q in (5.4) is inherited from that of S_q).

One can also decompose the above L as

$$L = \Sigma + N, \quad \Sigma \text{ diagonal and } N \text{ nilpotent}, \quad (5.5)$$

with

$$\Sigma = \sigma_1 I_1 \oplus \cdots \oplus \sigma_\ell I_\ell, \quad N = N_1 \oplus \cdots \oplus N_\ell.$$

Here I_1, \dots, I_ℓ are identity matrices, each I_q having the dimension of L_q . It follows that

$$[\Sigma, N] = 0. \quad (5.6)$$

Therefore,

$$z^D z^L = z^D z^\Sigma z^N = z^\Delta z^N,$$

where

$$\Delta := D + \Sigma \quad (5.7)$$

is a diagonal matrix, whose eigenvalues are the eigenvalues of J . The above properties allow to write

$$z^\Delta z^N = z^\Delta \sum_{k=1}^{\bar{k}} \frac{N^k}{k!} (\ln z)^k \quad \text{finite sum,}$$

where \bar{k} depends on the order of nilpotency of N .

In conclusion, a Levelt form (5.3) can be always chosen so that D and L (and so S and R) satisfy the above properties (5.5), (5.6) and (5.7), namely

$$Y(\zeta) = \mathcal{G}(\zeta)\zeta^D\zeta^L = \mathcal{G}(\zeta)\zeta^\Delta\zeta^N. \quad (5.8)$$

This can always be achieved by a permutation matrix P , by changing a fundamental matrix solution in Levelt form Y to another fundamental solution YP with Levelt form having the desired properties. In the Appendix, we will need the above choice of Levelt form.

Remark 5.1. In this section we have given the analytic construction of the “proper” Levelt form, just starting from the analytic structure of fundamental solutions at a Fuchsian singularity. For the geometric viewpoint, see [1].

6 Strong Isomonodromy Deformations

We define a **central connection matrix** $C_0(\lambda)$ associated with $Y_0(z, \lambda)$ in (4.9) with $\nu = 0$, and with a fundamental solution $Y^{(0)}(z, \lambda)$ in Levelt form (2.1) at $z = 0$, by

$$Y_0(z, \lambda) = Y^{(0)}(z, \lambda)C_0(\lambda).$$

Notice that

$$Y_\nu(z, \lambda) = Y^{(0)}(z, \lambda) C_0 \mathbb{S}_0 \cdots \mathbb{S}_{\nu-1}.$$

Definition 6.1. *Let Assumption 1 hold and let system (1.1) be weakly isomonodromic in \mathbb{D} with holomorphic fundamental matrix solution $Y^{\text{hol}} = Y^{(0)}$ in Levelt form, so that Theorem 3.1 holds, Assumption 2 is satisfied and the essential monodromy data*

$$L^{(0)}, \quad D^{(0)} \quad \text{are constant.}$$

*If also Assumption 3 holds, system (1.1) is said to be **strongly isomonodromic** on \mathbb{D} when also the remaining essential monodromy data are constant, namely*

$$d\mathbb{S}_\nu = 0, \quad dL = 0, \quad D \text{ is constant}, \quad dC_0 = 0.$$

Remark 6.1. For a strongly isomonodromic system, the relations (4.8), if any, are satisfied by definition, so that the fundamental matrices $Y_\nu(z, \lambda)$ are holomorphic on $\mathcal{R} \times \mathbb{D}$.

Theorem 6.1.

Part I. *System (1.1) is strongly isomonodromic in \mathbb{D} if and only if the fundamental matrix solutions $Y^{(0)}(z, \lambda)$ and $Y_\nu(z, \lambda)$ satisfy for every $\nu \in \mathbb{Z}$ the integrable Pfaffian system (3.1) of the specific form*

$$dY = \omega(z, \lambda)Y, \quad \omega(z, \lambda) = \left(\Lambda + \frac{A}{z} \right) dz + \sum_{j=1}^s \left(zE_{p_j} + \tilde{\omega}_j(\lambda) \right) d\lambda_j, \quad (6.1)$$

where

$$\tilde{\omega}_j(\lambda) = \omega_j(\lambda) + \frac{\partial \mathcal{T}(\lambda)}{\partial \lambda_j} \cdot \mathcal{T}(\lambda)^{-1},$$

and $\omega_j(\lambda)$ has blocks

$$\omega_{[a,a]}^{(j)}(\lambda) = 0, \quad \omega_{[a,b]}^{(j)}(\lambda) = \frac{A_{[a,b]}(\lambda) (\delta_{aj} - \delta_{bj})}{\lambda_a - \lambda_b}, \quad a \neq b = 1, \dots, s, \quad (6.2)$$

while

$$\mathcal{T}(\lambda) = \mathcal{T}_1(\lambda) \oplus \dots \oplus \mathcal{T}_s(\lambda)$$

is a holomorphic matrix reducing to Jordan form the diagonal blocks of A as in (4.1). Strong isomonodromy deformations with constant \mathcal{T} are allowed. In this case,

$$\tilde{\omega}_j(\lambda) = \omega_j(\lambda).$$

Part II. If system (1.1) is strongly isomonodromic in \mathbb{D} , then

$$\frac{\partial A}{\partial \lambda_j} = [\tilde{\omega}_j(u), A], \quad j = 1, \dots, s. \quad (6.3)$$

In particular,

$$\frac{\partial A_{[1,1]}}{\partial \lambda} = \frac{\partial A_{[2,2]}}{\partial \lambda} \dots = \frac{\partial A_{[s,s]}}{\partial \lambda} = 0,$$

and so the block-diagonal part of A and the Jordan forms J_k in (4.1) are constant on \mathbb{D} .

For a strong isomonodromy deformation, PART II says that $A_{[1,1]} \oplus \dots \oplus A_{[s,s]}$ is constant, so it can be reduced to Jordan form by a constant block-diagonal matrix \mathcal{T}_0 . If $\mathcal{T} = \mathcal{T}_1 \oplus \dots \oplus \mathcal{T}_s$ is another matrix satisfying (4.1), but not constant, then it has the structure

$$\mathcal{T}(\lambda) = \mathcal{T}_0 \mathfrak{B}(\lambda), \quad \mathfrak{B}(\lambda) = \mathfrak{B}_1(\lambda) \oplus \dots \oplus \mathfrak{B}_s(\lambda), \quad \text{with} \quad [\mathfrak{B}_j(\lambda), J_j] = 0.$$

The isomonodromic fundamental matrix solutions $Y_\nu(z, \lambda)$ which satisfy the Pfaffian system (6.1) have structure (4.9) with constant $\mathcal{T} = \mathcal{T}_0$ if and only if system (6.1) is of the specific form with coefficients $\tilde{\omega}_j(\lambda) = \omega_j(\lambda)$.

In other words, if a differential system

$$\frac{dY}{dz} = \left(\Lambda_0 + \frac{A_0}{z} \right) Y$$

is given at $\lambda = \lambda_0$, where Λ_0 has repeated eigenvalues, then it can have different isomonodromy deformations (6.1), differing by the specific $\sum_j \tilde{\omega}_j(\lambda) d\lambda_j$, namely by the specific $\mathcal{T}(\lambda)$. For all these deformations, the diagonal blocks are constant and equal to those of A_0 , but the off-diagonal blocks of $A(\lambda)$ are different for different deformations, satisfying different systems (6.3), with different⁴ matrix coefficients $\tilde{\omega}_j$ depending on the choice of $\mathcal{T}(\lambda)$.

⁴This allows, in case $n = 3$ and $s = 2$, to have isomonodromy deformations with constant A , see Section 7. For $n \geq 4$, it is not possible to reach a constant A by a gauge transformation, see Remark 1.2.

Two isomonodromy deformations with different \mathcal{T} are related by a gauge transformation. Suppose that Y satisfies (6.1) with matrices $A(\lambda)$ and $\tilde{\omega}_j(\lambda) = \omega_j(\lambda) + \partial_j \mathcal{T} \cdot \mathcal{T}^{-1}$, where ω_j is in (6.2). Consider the gauge transformation

$$Y = \mathcal{T}(\lambda) \check{\mathcal{T}}^{-1}(\lambda) \check{Y},$$

where $\check{\mathcal{T}}$ is another matrix Jordanizing the block-diagonal part of $A(\lambda)$. Then, \check{Y} satisfies a system (6.1) of the form

$$d\check{Y} = \left[\left(\Lambda + \frac{\check{A}}{z} \right) dz + \sum_{j=1}^s \left(zE_{p_j} + \check{\omega}_j \right) d\lambda_j + d\check{\mathcal{T}} \cdot \check{\mathcal{T}}^{-1} \right] \check{Y}.$$

where

$$\check{A}(\lambda) := \check{\mathcal{T}}(\mathcal{T}^{-1}A(\lambda)\mathcal{T})\check{\mathcal{T}}^{-1}, \quad \check{\omega}_j(\lambda) = \check{\mathcal{T}}(\mathcal{T}^{-1}\omega_j(\lambda)\mathcal{T})\check{\mathcal{T}}^{-1}. \quad (6.4)$$

Notice that $\check{\omega}_j$ is the same as in definition (6.2) with A replaced by \check{A} (the block-diagonal parts of A and \check{A} are the same).

Remark 6.2. Suppose that we have a deformation with $\tilde{\omega}_j = \omega_j$ for all j . Then the dependence of Y_ν on λ is

$$Y_\nu(z, \lambda) = \mathcal{T} \hat{Y}_\nu(z; \lambda_2 - \lambda_1, \dots, \lambda_s - \lambda_1) z^D z^L e^{\Lambda z}.$$

Indeed, formula (6.2) implies that

$$\sum_{j=1}^s \omega_j(\lambda) = 0, \quad \implies \quad \sum_{j=1}^s \frac{\partial A}{\partial \lambda_j} = 0, \quad \implies \quad A = A(\lambda_2 - \lambda_1, \dots, \lambda_s - \lambda_1).$$

Moreover, let us write (4.9) as

$$Y_\nu(z, \lambda) = H_\nu(z, \lambda) e^{\Lambda z} \quad \text{with} \quad H_\nu := \mathcal{T} \hat{Y}_\nu z^D z^L.$$

Then $\frac{\partial H_\nu e^{\Lambda z}}{\partial \lambda_j} = \frac{\partial H_\nu}{\partial \lambda_j} e^{\Lambda z} + H_\nu \cdot z E_{p_j} e^{\Lambda z}$. Since Y_ν satisfies (6.1) with $\tilde{\omega}_j = \omega_j$, we also have $\frac{\partial H_\nu e^{\Lambda z}}{\partial \lambda_j} = (z E_{p_j} + \omega_j(\lambda)) H_\nu e^{\Lambda z}$. Thus,

$$\frac{\partial H_\nu}{\partial \lambda_j} = z[E_{p_j}, H_\nu] + \omega_j(\lambda) H_\nu, \quad \implies \quad \sum_{j=1}^s \frac{\partial H_\nu}{\partial \lambda_j} = 0,$$

so that $H_\nu = H_\nu(z; \lambda_2 - \lambda_1, \dots, \lambda_s - \lambda_1)$. Notice also that $z^D z^L$ commutes with E_{p_j} and that in the strong isomonodromic case $d\mathcal{T} = dD = dL = 0$, so that we also obtain

$$\frac{\partial \hat{Y}_\nu}{\partial \lambda_j} = z[E_{p_j}, \hat{Y}_\nu] + \omega_j(\lambda) \hat{Y}_\nu, \quad \implies \quad \sum_{j=1}^s \frac{\partial \hat{Y}_\nu}{\partial \lambda_j} = 0,$$

This concludes.

Proof of Theorem 6.1. Let system (1.1) be strongly isomonodromic in \mathbb{D} (Definition 6.1), so that $Y^{\text{hol}} = Y^{(0)}$ in Levelt form is holomorphic, and by Assumption 3 all the Y_ν are holomorphic. Thus, we can take differentials. We define

$$\omega(z, \lambda) := dY^{(0)} \cdot (Y^{(0)})^{-1} \underset{dC^{(0)}=0}{=} dY_0 \cdot (Y_0)^{-1} \underset{\text{all } d\mathbb{S}_\nu=0}{=} dY_\nu \cdot (Y_\nu)^{-1}, \quad \forall \nu \in \mathbb{Z}.$$

This is single valued for the counter-clockwise loop $z \mapsto ze^{2\pi i}$, because the monodromy $e^{2\pi i L^{(0)}}$ of $Y^{(0)}$ and the monodromy $e^{2\pi i L} (\mathbb{S}_\nu \mathbb{S}_{\nu+1})^{-1}$ of Y_ν are constant. Its singularities may only be located at $z = 0, \infty$. We find the structure of ω at $z = 0$ and $z = \infty$ respectively.

Structure at $z = 0$. Let us decompose the differential as $d = d_z + d_\lambda$, the former being the component on dz , the latter on $d\lambda_1, \dots, d\lambda_s$. Firstly, we compute

$$\begin{aligned} d_z Y^{(0)} \cdot (Y^{(0)})^{-1} &= G^{(0)} d_z \hat{Y}^{(0)} \cdot (\hat{Y}^{(0)})^{-1} + G^{(0)} \hat{Y}^{(0)} \frac{D^{(0)} + z^{D^{(0)}} L z^{-D^{(0)}}}{z} (G^{(0)} \hat{Y}^{(0)})^{-1} dz \\ &= \left(\frac{A}{z} + \text{reg}_1(z, \lambda) \right) dz, \quad \text{reg}_1(z, \lambda) = O(1), \quad z \rightarrow 0, \end{aligned}$$

where we have used (3.9). Here, $\text{reg}_1(z, \lambda)$ is holomorphic for $z \in \mathbb{C}$ and $\lambda \in \mathbb{D}$. Then, we compute

$$\begin{aligned} d_\lambda Y^{(0)} \cdot (Y^{(0)})^{-1} &\underset{dD^{(0)}=dL^{(0)}=0}{=} d_\lambda G^{(0)} \cdot (G^{(0)})^{-1} + G^{(0)} d_\lambda \hat{Y}^{(0)} \cdot (\hat{Y}^{(0)})^{-1} (G^{(0)})^{-1} \\ &= d_\lambda G^{(0)} \cdot (G^{(0)})^{-1} + \text{reg}(z, \lambda), \quad \text{reg}(z, \lambda) = O(z) \rightarrow 0, \quad z \rightarrow 0, \end{aligned}$$

where $\text{reg}(z, \lambda)$ is holomorphic for $z \in \mathbb{C}$ and $\lambda \in \mathbb{D}$.

Structure at $z = \infty$. Firstly, we compute

$$\begin{aligned} d_z Y_\nu \cdot Y_\nu^{-1} &= \mathcal{T} d_z \hat{Y}_\nu \cdot \hat{Y}_\nu^{-1} \mathcal{T}^{-1} + \\ &+ \left(\mathcal{T} \hat{Y}_\nu \frac{D + z^D L z^{-D}}{z} \hat{Y}_\nu^{-1} \mathcal{T}^{-1} + \mathcal{T} \hat{Y}_\nu z^D z^L \Lambda z^{-L} z^{-D} \hat{Y}_\nu^{-1} \mathcal{T}^{-1} \right) dz. \end{aligned}$$

Due to the block structure of D and L and diagonality of Λ , we have $z^D z^L \Lambda z^{-L} z^{-D} = \Lambda$, while by (4.3), (4.6), (4.7) we have $(D + z^D L z^{-D})/z = J/z + O(z^{-2})$. Hence,

$$d_z Y_\nu \cdot Y_\nu^{-1} = \left(\Lambda + \widetilde{\text{reg}}(z^{-1}, \lambda) \right) dz, \quad \widetilde{\text{reg}}(z^{-1}, \lambda) = O\left(\frac{1}{z}\right) \rightarrow 0, \quad z \rightarrow \infty,$$

being $\widetilde{\text{reg}}(1/z, \lambda)$ analytic for $z \in \overline{\mathbb{C}} \setminus \{0\}$ and $\lambda \in \mathbb{D}$. Then, we compute

$$d_\lambda Y_\nu \cdot Y_\nu^{-1} \underset{dD=dL=0}{=} d_\lambda \mathcal{T} \cdot \mathcal{T}^{-1} + \mathcal{T} d_\lambda \hat{Y}_\nu \cdot \hat{Y}_\nu^{-1} \mathcal{T}^{-1} + z \cdot \mathcal{T} \hat{Y}_\nu z^D z^L d\Lambda z^{-L} z^{-D} (\mathcal{T} \hat{Y}_\nu)^{-1}.$$

As before, from the diagonality of $d\Lambda$ we receive $z^D z^L d\Lambda z^{-L} z^{-D} = d\Lambda$, so that

$$d_\lambda Y_\nu \cdot Y_\nu^{-1} = d_\lambda \mathcal{T} \cdot \mathcal{T}^{-1} + z d\Lambda + \mathcal{T} [F_1, d\Lambda] \mathcal{T}^{-1} + \widehat{\text{reg}}(z^{-1}, \lambda),$$

where $\widehat{\text{reg}}(z^{-1}, \lambda)$ is a 1-form in $d\lambda_1, \dots, d\lambda_n$, analytic for $z \in \overline{\mathbb{C}} \setminus \{0\}$ and $\lambda \in \mathbb{D}$, with behaviour

$$\widehat{\text{reg}}(z^{-1}, \lambda) = O\left(\frac{1}{z}\right) \rightarrow 0, \quad z \rightarrow \infty,$$

Notice that $d\Lambda = E_{p_1}d\lambda_1 + \dots + E_{p_s}d\lambda_s$. By Liouville theorem and the above behaviours at $z = 0, \infty$ we conclude that

$$\omega = \left(\Lambda + \frac{A}{z} \right) dz + \sum_{j=1}^s \left(zE_{p_j} + [\mathcal{T}F_1\mathcal{T}^{-1}, E_{p_j}] + \frac{\partial \mathcal{T}}{\partial \lambda_j} \mathcal{T}^{-1} \right) d\lambda_j. \quad (6.5)$$

It remains to show that $[\mathcal{T}F_1(\lambda)\mathcal{T}^{-1}, E_{p_j}]$ equals (6.2). The computations of section 4.1 of [13], Proposition 4,1, yield the off-diagonal blocks

$$F_{[i,j]}^{(1)} = \frac{\mathcal{A}_{[i,j]}}{\lambda_j - \lambda_i}, \quad 1 \leq i \neq j \leq s. \quad (6.6)$$

They suffice to evaluate $[\mathcal{T}F_1\mathcal{T}^{-1}, E_{p_j}]$, since the diagonal blocks do not contribute. From the definition of \mathcal{A} in (4.4) and (6.6) we receive

$$F_{[i,j]}^{(1)} = \frac{\mathcal{T}_i^{-1} A_{[i,j]} \mathcal{T}_j}{\lambda_j - \lambda_i} \implies (\mathcal{T}F_1\mathcal{T}^{-1})_{[i,j]} = \mathcal{T}_i F_{[i,j]}^{(1)} \mathcal{T}_j^{-1} = \frac{A_{[i,j]}}{\lambda_j - \lambda_i}.$$

Using the last formula, we conclude that

$$[\mathcal{T}F_1(\lambda)\mathcal{T}^{-1}, E_{p_k}]_{[a,a]} = 0, \quad [\mathcal{T}F_1(\lambda)\mathcal{T}^{-1}, E_{p_k}]_{[a,b]} = \frac{A_{[a,b]}(\lambda) (\delta_{ak} - \delta_{bk})}{\lambda_a - \lambda_b}, \quad a \neq b.$$

It remains to show that isomonodromy deformations with constant \mathcal{T} are possible. This will be proved after Lemma 6.1.

• Conversely, we assume that all the fundamental matrices $Y^{(0)}$ and Y_ν , $\nu \in \mathbb{Z}$, of system (1.1) also satisfy

$$dY = \left[\left(\Lambda + \frac{A}{z} \right) dz + \sum_{j=1}^s \left(zE_{p_j} + \tilde{\omega}_j(\lambda) \right) d\lambda_j \right] Y,$$

with holomorphic $\tilde{\omega}_j(\lambda)$ and $A(\lambda)$. In particular, this means that $dY^{(0)} \cdot (Y^{(0)})^{-1}$ and $dY_\nu \cdot Y_\nu^{-1}$ depend homomorphically on λ .

Since ω has Fuchsian singularity at $z = 0$, by Proposition 3.2 we know that indeed it has holomorphic solution $Y^{(0)}$ in Levelt form, and⁵

$$D^{(0)}, \quad L^{(0)} \quad \text{are constant.}$$

The fact that $dC_0 = d\mathbb{S}_\nu = 0$ is straightforward. Indeed, since all fundamental solutions satisfy $dY = \omega Y$, we have

$$dY_{\nu+1} \cdot Y_{\nu+1}^{-1} = dY_\nu \cdot Y_\nu^{-1} \iff d\mathbb{S}_\nu = 0.$$

⁵This can also be seen directly by taking

$$\begin{aligned} d_\lambda Y^{(0)} \cdot (Y^{(0)})^{-1} &= d_\lambda (G^{(0)} \hat{Y}^{(0)}) \cdot (G^{(0)} \hat{Y}^{(0)})^{-1} + \\ &+ (G^{(0)} Y^{(0)}) \left(d\Delta^{(0)} \ln z + z^{\Delta^{(0)}} \sum_{k=1}^{\bar{k}_0} \frac{d(N^{(0)})^k}{k!} (\ln z)^k z^{-N^{(0)}} z^{-\Delta^{(0)}} \right) (G^{(0)} Y^{(0)})^{-1}. \end{aligned}$$

Since ω does not contain terms in $\ln z$, it follows that $d\Delta^{(0)} = dN^{(0)} = 0$, so that $D^{(0)}$ and $L^{(0)}$ are constant. Here, $\Delta^{(0)}$ and $N^{(0)}$ are the analogues of Δ and N in (5.5).

$$dY_0 \cdot Y_0^{-1} = dY^{(0)} \cdot (Y^{(0)})^{-1} \iff dC_0 = 0.$$

Finally, we show that D and L are constant. Recall from Section 5 that we can write $Y_\nu = \mathcal{T}\widehat{Y}_\nu z^\Delta z^N$, with diagonal Δ and nilpotent N . By assumption

$$\begin{aligned} \sum_{j=1}^s (zE_{p_j} + \tilde{\omega}_j(\lambda)) d\lambda_j &= d_\lambda Y_\nu \cdot Y_\nu^{-1} = d\mathcal{T} \cdot \mathcal{T}^{-1} + \mathcal{T} d_\lambda \widehat{Y}_\nu \cdot \widehat{Y}_\nu^{-1} \mathcal{T}^{-1} + \\ &+ \mathcal{T} \widehat{Y}_\nu \left(\ln z d\Delta + z^\Delta \sum_{k=1}^{\bar{k}} \frac{dN^k}{k!} (\ln z)^k z^{-N} z^{-\Delta} \right) (\mathcal{T} \widehat{Y}_\nu)^{-1} + \\ &+ z \cdot \mathcal{T} \widehat{Y}_\nu d\Lambda (\mathcal{T} \widehat{Y}_\nu)^{-1}. \end{aligned}$$

Since logarithmic terms cannot occur, necessarily $d\Delta = dN = 0$, so that D and L are constant. From the dominant terms at $z = \infty$ in the above computation we receive

$$\tilde{\omega}_j(\lambda) = [\mathcal{T} F_1 \mathcal{T}^{-1}, E_{p_j}] + \frac{\partial \mathcal{T}}{\partial \lambda_j} \mathcal{T}^{-1}.$$

• PART II. Suppose the system is strongly isomonodromic. By PART I, the matrices $Y^{(0)}$ and $Y^{(\nu)}$ solve a Pfaffian system $dY = \omega Y$ where ω has structure

$$\omega = \left(\Lambda + \frac{A}{z} \right) dz + \sum_{j=1}^s \left(zE_{p_j} + \tilde{\omega}_j(\lambda) \right) d\lambda_j.$$

We write for short

$$\omega = \sum_{\alpha=0}^s \varphi_\alpha(x) dx^\alpha, \quad (x^0, x^1, \dots, x^s) := (z, \lambda_1, \dots, \lambda_s).$$

Thus, ω is integrable, i.e. $d\omega = \omega \wedge \omega$, which explicitly is

$$\frac{\partial \varphi_\beta}{\partial x^\alpha} + \varphi_\beta \varphi_\alpha = \frac{\partial \varphi_\alpha}{\partial x^\beta} + \varphi_\alpha \varphi_\beta, \quad \alpha \neq \beta = 0, 1, \dots, s, \quad (6.7)$$

For $\beta = 0$ and $\alpha = j \in \{1, \dots, s\}$, (6.7) is

$$\frac{\partial}{\partial \lambda_j} \left(\Lambda + \frac{A}{z} \right) + \left(\Lambda + \frac{A}{z} \right) (zE_{p_j} + \tilde{\omega}_j(\lambda)) = \frac{\partial}{\partial z} (zE_{p_j} + \tilde{\omega}_j(\lambda)) + (zE_{p_j} + \tilde{\omega}_j(\lambda)) \left(\Lambda + \frac{A}{z} \right).$$

Expanding, we see that the equality is true if and only if the coefficients of z^{-1} and z^0 are respectively equal, namely

$$\frac{\partial A}{\partial \lambda_j} = [\tilde{\omega}_j(\lambda), A], \quad (6.8)$$

$$[\Lambda, \tilde{\omega}_j(\lambda)] = [E_{p_j}, A]. \quad (6.9)$$

The equations (6.9) have general solution

$$[\tilde{\omega}_j(\lambda)]_{\text{block } a,a} \text{ arbitrary, } [\tilde{\omega}_j(\lambda)]_{\text{block } a,b} = \frac{A_{[a,b]}(\lambda) (\delta_{aj} - \delta_{bj})}{\lambda_a - \lambda_b}, \quad a \neq b = 1, \dots, s.$$

Thus,

$$\tilde{\omega}_j(\lambda) = \omega_j(\lambda) + \mathcal{D}_j(\lambda), \quad (6.10)$$

where $\omega_j(\lambda)$ is (6.2), while $\mathcal{D}_j(\lambda)$ is an arbitrary block-diagonal matrix, which in our case must be

$$\mathcal{D}_j = \frac{\partial \mathcal{T}}{\partial \lambda_j} \cdot \mathcal{T}^{-1}.$$

The integrability condition (6.7) for $\alpha = k$, $\beta = j$, with $j \neq k \in \{1, \dots, s\}$ is

$$\frac{\partial}{\partial \lambda_k} \left(z E_{p_j} + \tilde{\omega}_j(\lambda) \right) + \left(z E_{p_j} + \tilde{\omega}_j(\lambda) \right) \left(z E_{p_k} + \tilde{\omega}_k(\lambda) \right) = \text{the same with } j, k \text{ exchanged}$$

This is true if and only if

$$[E_{p_j}, \tilde{\omega}_k(\lambda)] = [E_{p_k}, \tilde{\omega}_j(\lambda)], \quad (6.11)$$

$$\frac{\partial \tilde{\omega}_j(\lambda)}{\partial \lambda_k} + \tilde{\omega}_j(\lambda) \tilde{\omega}_k(\lambda) = \frac{\partial \tilde{\omega}_k(\lambda)}{\partial \lambda_j} + \tilde{\omega}_k(\lambda) \tilde{\omega}_j(\lambda), \quad 1 \leq j \neq k \leq s. \quad (6.12)$$

The equalities (6.11) are automatically satisfied⁶ by (6.10) and (6.2), while (6.12) implies the Frobenius integrability of (6.8). Notice that the block-diagonal part of (6.12) is (see Lemma 6.2 for details)

$$\frac{\partial \mathcal{D}_j(\lambda)}{\partial \lambda_k} + \mathcal{D}_j(\lambda) \mathcal{D}_k(\lambda) = \frac{\partial \mathcal{D}_k(\lambda)}{\partial \lambda_j} + \mathcal{D}_k(\lambda) \mathcal{D}_j(\lambda), \quad j, k = 1, \dots, s,$$

and admits in particular the holomorphic solution $\mathcal{D}_j = \partial \mathcal{T} / \partial \lambda_j \cdot \mathcal{T}^{-1}$ for some holomorphic $\mathcal{T}(\lambda) = \mathcal{T}_1(\lambda) \oplus \dots \oplus \mathcal{T}_s(\lambda)$, in accordance with the required structure (6.5), which necessarily holds in case of strong isomonodromic deformations.

Let $A_{[k,k]}$ denote as usual a diagonal-block of A , and let $\mathcal{D}_{[k,k]}^{(j)}$ be a diagonal-block of \mathcal{D}_j , $k = 1, \dots, s$. The block diagonal part of (6.8) now reduces to

$$\frac{\partial A_{[k,k]}}{\partial \lambda_j} = [\mathcal{D}_{[k,k]}^{(j)}, A_{[k,k]}], \quad k = 1, \dots, s. \quad (6.13)$$

⁶First, notice that each \mathcal{D}_j commutes with Λ and each E_{p_k} , so that

$$[\Lambda, \tilde{\omega}_j(\lambda)] = [E_{p_j}, A], \quad [E_{p_j}, \tilde{\omega}_k(\lambda)] = [E_{p_k}, \tilde{\omega}_j(\lambda)], \quad 1 \leq j \neq k \leq s,$$

are equivalent to

$$[\Lambda, \omega_j(\lambda)] = [E_{p_j}, A], \quad [E_{p_j}, \omega_k(\lambda)] = [E_{p_k}, \omega_j(\lambda)], \quad 1 \leq j \neq k \leq s.$$

Then, we show that

$$[\Lambda, \omega_j(\lambda)] = [E_{p_j}, A], \implies [E_{p_j}, \omega_k(\lambda)] = [E_{p_k}, \omega_j(\lambda)], \quad 1 \leq j \neq k \leq s.$$

Indeed, the $[\Lambda, \omega_j(\lambda)] = [E_{p_j}, A]$ imply that the matrices ω_j are as in (6.2). Substituting (6.2) we obtain the blocks

$$\begin{aligned} & \left([E_{p_j}, \omega_k] - [E_{p_k}, \omega_j] \right)_{[a,b]} = \delta_{ja} \omega_{[a,b]}^{(k)} - \omega_{[a,b]}^{(k)} \delta_{jb} - \delta_{ka} \omega_{[a,b]}^{(j)} + \omega_{[a,b]}^{(j)} \delta_{kb} \\ & = \frac{A_{[a,b]}}{\lambda_a - \lambda_b} \left(\delta_{ja} (\delta_{ka} - \delta_{kb}) - \delta_{jb} (\delta_{ka} - \delta_{kb}) - \delta_{ka} (\delta_{ja} - \delta_{jb}) + \delta_{kb} (\delta_{ja} - \delta_{jb}) \right) = 0. \end{aligned}$$

If $\mathcal{D} = 0$, namely \mathcal{T} is constant, the above implies constancy of the block diagonal part of A . If \mathcal{T} is not constant, we need the following technical Lemma, proved in the Appendix.

Lemma 6.1. *Let $\mathcal{T} = \mathcal{T}_1 \oplus \cdots \oplus \mathcal{T}_s$ be a matrix yielding a Jordan form $J = J_1 \oplus \cdots \oplus J_s = \mathcal{T}^{-1} \left(A_{[1,1]} \oplus \cdots \oplus A_{[s,s]} \right) \mathcal{T}$, where each J_k is as in (4.2). If the deformation is strongly isomonodromic, then*

$$[\mathcal{T}^{-1} d\mathcal{T}, J] = 0.$$

In case of strong isomonodromic deformation, we have $\mathcal{D}_j = \partial\mathcal{T}/\partial\lambda_j \cdot \mathcal{T}^{-1}$. Using Lemma 6.1 and (6.13), we prove that $A_{[1,1]}, \dots, A_{[s,s]}$ are constant. Indeed

$$\begin{aligned} \frac{\partial A_{[k,k]}}{\partial\lambda_j} &= [\mathcal{D}_{[k,k]}^{(j)}, A_{[k,k]}] \\ &= \frac{\partial\mathcal{T}_k}{\partial\lambda_j} \mathcal{T}_k^{-1} A_{[k,k]} - A_{[k,k]} \frac{\partial\mathcal{T}_k}{\partial\lambda_j} \mathcal{T}_k^{-1} \\ &= \frac{\partial\mathcal{T}_k}{\partial\lambda_j} J_k \mathcal{T}_k^{-1} - \mathcal{T}_k J_k \left(\mathcal{T}_k^{-1} \frac{\partial\mathcal{T}_k}{\partial\lambda_j} \right) \mathcal{T}_k^{-1} \\ &\stackrel{\text{Lemma 6.1}}{=} \frac{\partial\mathcal{T}_k}{\partial\lambda_j} J_k \mathcal{T}_k^{-1} - \mathcal{T}_k \mathcal{T}_k^{-1} \frac{\partial\mathcal{T}_k}{\partial\lambda_j} J_k \mathcal{T}_k^{-1} = 0. \end{aligned}$$

This proves PART II.

By the constancy of the diagonal blocks of A for a strong isomonodromy deformation, it is possible to consider deformations with \mathcal{T} constant. Conversely, if \mathcal{T} is constant, so that all $\mathcal{D}_j = 0$, then (6.13) implies that all the $A_{[k,k]}$ are constant. \square

It is convenient to point out from the proof above that $[\Lambda, \tilde{\omega}_j] = [E_{p_j}, A]$ is equivalent to

$$\tilde{\omega}_j(\lambda) = \omega_j(\lambda) + \mathcal{D}_j(\lambda) \tag{6.14}$$

where $\omega_j(\lambda)$ is (6.2) and

$$\mathcal{D}_j = \mathcal{D}_{[1,1]}^{(j)} \oplus \cdots \oplus \mathcal{D}_{[s,s]}^{(j)} \quad \text{arbitrary block-diagonal matrix.} \tag{6.15}$$

It is also convenient to state the following

Lemma 6.2. *Assume that the matrices $\tilde{\omega}_j(\lambda)$ are defined by*

$$[\Lambda, \tilde{\omega}_j] = [E_{p_j}, A], \quad j = 1, \dots, s,$$

so that they have structure (6.14). Let $\partial_j := \frac{\partial}{\partial\lambda_j}$. The following facts hold.

a) The system

$$\partial_i \tilde{\omega}_j - \partial_j \tilde{\omega}_i = [\tilde{\omega}_i, \tilde{\omega}_j], \quad i, j = 1, \dots, s,$$

is equivalent to

$$\begin{cases} \partial_i \omega_j - \partial_j \omega_i = [\omega_i, \omega_j] + [\omega_i, \mathcal{D}_j] + [\mathcal{D}_j, \omega_i] & \text{block off-diagonal,} \\ \partial_i \mathcal{D}_j - \partial_j \mathcal{D}_i = [\mathcal{D}_i, \mathcal{D}_j] & \text{block diagonal.} \end{cases}$$

b) Assume moreover that

$$dA = \left[\sum_{j=1}^n \tilde{\omega}_j(\lambda) d\lambda_j, A \right], \quad (6.16)$$

then

$$\begin{cases} \partial_i \tilde{\omega}_j - \partial_j \tilde{\omega}_i - [\tilde{\omega}_i, \tilde{\omega}_j] = \partial_i \mathcal{D}_j - \partial_j \mathcal{D}_i - [\mathcal{D}_i, \mathcal{D}_j], \\ \partial_i \omega_j - \partial_j \omega_i - \left([\omega_i, \omega_j] + [\omega_i, \mathcal{D}_j] + [\mathcal{D}_j, \omega_i] \right) = 0; \end{cases}$$

and

$$dA^D = [\mathcal{D}, A^D],$$

where

$$A^D(\lambda) := A_{[1,1]}(\lambda) \oplus \cdots \oplus A_{[s,s]}(\lambda) \quad (6.17)$$

Notice that part b) above implies that if (6.16) holds, then the system $\partial_i \tilde{\omega}_j - \partial_j \tilde{\omega}_i = [\tilde{\omega}_i, \tilde{\omega}_j]$ is equivalent to $\partial_i \mathcal{D}_j - \partial_j \mathcal{D}_i = [\mathcal{D}_i, \mathcal{D}_j]$, $i, j = 1, \dots, s$.

Proof. a) Given any matrix $\tilde{\omega}_j$, we can write it as $\tilde{\omega}_j = \omega_j + \mathcal{D}_j$, a sum of a block off-diagonal term ω_j + block diagonal term \mathcal{D}_j . Then

$$\begin{aligned} & \partial_i \tilde{\omega}_j - \partial_j \tilde{\omega}_i - [\tilde{\omega}_i, \tilde{\omega}_j] = \\ & = \underbrace{\left\{ \underbrace{\partial_i \omega_j - \partial_j \omega_i}_{(*1*)} - \underbrace{\left([\omega_i, \omega_j] + [\omega_i, \mathcal{D}_j] + [\mathcal{D}_j, \omega_i] \right)}_{(*2*)} \right\}}_{(\bullet)} + \underbrace{\left\{ \partial_i \mathcal{D}_j - \partial_j \mathcal{D}_i - [\mathcal{D}_i, \mathcal{D}_j] \right\}}_{(*3*)} \end{aligned}$$

Clearly, (*1*) is block off-diagonal and (*3*) is block diagonal. It is easy to see that in (*2*), also $[\omega_i, \mathcal{D}_j] + [\mathcal{D}_j, \omega_i]$ is block off-diagonal. Indeed

$$[\omega_i, \mathcal{D}_j]_{[a,a]} = \omega_{[a,a]}^{(i)} \mathcal{D}_{[a,a]}^{(j)} - \mathcal{D}_{[a,a]}^{(j)} \omega_{[a,a]}^{(i)} \Big|_{\omega_{[a,a]}^{(i)}=0} = 0.$$

If the matrices $\tilde{\omega}_j$ are defined by $[\Lambda, \tilde{\omega}_j(\lambda)] = [E_{p_j}, A]$ for $j = 1, \dots, s$, namely if ω_j is as in (6.2), then the whole (\bullet) is block off-diagonal. Indeed,

$$\begin{aligned} [\omega_i, \omega_j]_{[a,a]} &= \sum_{b \neq a} (\omega_{[a,b]}^{(i)} \omega_{[b,a]}^{(j)} - \omega_{[a,b]}^{(j)} \omega_{[b,a]}^{(i)}) \\ &\stackrel{(6.2)}{=} \sum_{b \neq a} \frac{A_{[a,b]} A_{[b,a]}}{(\lambda_a - \lambda_b)(\lambda_b - \lambda_a)} ((\delta_{ia} - \delta_{ib})(\delta_{jb} - \delta_{ja}) - (\delta_{ja} - \delta_{jb})(\delta_{ib} - \delta_{ia})) = 0. \end{aligned}$$

This proves a).

b) We consider the blocks of (\bullet) and substitute $\tilde{\omega}_k = \omega_k + \mathcal{D}_k$ as in (6.14). Then, where derivatives of the blocks $\omega_{[a,b]}^{(k)}$ occur, we express the $\omega_{[a,b]}^{(k)}$ in terms of A using (6.2), and then we re-substitute $\hat{\partial}_k A_{[a,b]} = [\omega_k + \mathcal{D}_k, A]_{[a,b]}$. After a lengthy computation we receive

$$(\bullet) = 0.$$

This proves the first part of b). The blocks of (6.16) are

$$\frac{\partial A_{[a,b]}}{\partial \lambda_j} = \sum_{c=1}^s \left(\omega_{[a,c]}^{(j)} A_{[c,b]} - A_{[a,c]} \omega_{[c,b]}^{(j)} \right) + \mathcal{D}_a^{(j)} A_{[a,b]} - A_{[a,b]} \mathcal{D}_b^{(j)}.$$

For $a = b$, the structure (6.2) implies that

$$\sum_{c=1}^s \left(\omega_{[a,c]}^{(j)} A_{[c,a]} - A_{[a,c]} \omega_{[c,a]}^{(j)} \right) = 0, \quad \implies \quad \frac{\partial A_{[a,a]}}{\partial \lambda_j} = [\mathcal{D}_a^{(j)}, A_{[a,a]}].$$

□

We state the converse of Part II of Theorem 6.1.

Theorem 6.2. *Let A and $\tilde{\omega}_1, \dots, \tilde{\omega}_s$ satisfy the system*

$$[\Lambda, \tilde{\omega}_j] = [E_{p_j}, A], \quad j = 1, \dots, s, \quad (\text{so that } \tilde{\omega}_j = \omega_j + \mathcal{D}_j \text{ as in (6.14)}); \quad (6.18)$$

$$dA = \left[\sum_{j=1}^n \tilde{\omega}_j(\lambda) d\lambda_j, A \right]; \quad (6.19)$$

$$\partial_i \mathcal{D}_j - \partial_j \mathcal{D}_i = [\mathcal{D}_i, \mathcal{D}_j], \quad 1 \leq i \neq j \leq s; \quad (6.20)$$

Then, the following facts hold.

1. System (6.19) is Frobenius integrable.
2. The connection

$$\omega(z, \lambda) = \left(\Lambda + \frac{A}{z} \right) dz + \sum_{j=1}^s \left(z E_{p_j} + \tilde{\omega}_j(\lambda) \right) d\lambda_j. \quad (6.21)$$

with matrices (6.2), (6.14) and (6.15) is Frobenius integrable. The Pfaffian system $dY = \omega Y$ has a fundamental matrix solution $Y^{(0)}(z, \lambda)$ in Levelt form (2.1), with constant exponents as in (3.6).

3. Assumption 3 holds and $A^D(\lambda)$ defined in (6.17) admits constant Jordan form

$$\mathcal{T} A^D \mathcal{T}^{-1} = J \equiv J_1 \oplus \dots \oplus J_s,$$

where $\mathcal{T}(\lambda)$ is a holomorphic invertible matrix solution of the integrable Pfaffian system

$$d\mathcal{T} = \mathcal{D}(\lambda) \mathcal{T}, \quad (6.22)$$

where $\mathcal{D}(\lambda) := \sum_{j=1}^n \mathcal{D}_j(\lambda) d\lambda_j$. Such a $\mathcal{T}(\lambda)$ can be chosen block diagonal as J .

4. If there are no partial resonances (i.e. all the blocks $A_{[k,k]}$, $k = 1, \dots, s$, are non-resonant), then every $Y_\nu(z, \lambda)$ in (4.9) with \mathcal{T} as in point 3. satisfies

$$dY_\nu = \omega(z, \lambda) Y_\nu,$$

so that system (1.1) is strongly isomonodromic.

Remark 6.3. We can choose the solution $\mathcal{D} = 0$ of (6.20), and in this case \mathcal{T} at point 3. is constant.

Remark 6.4. In the proof of Theorem 6.2, we will see that for every $\nu \in \mathbb{Z}$ there exists a holomorphic matrix valued one-form $K_\nu(\lambda)$ such that each $Y_\nu(z, \lambda)$ in (4.9), with \mathcal{T} as at point 2., satisfies the system

$$dY_\nu = \omega(z, \lambda)Y_\nu + Y_\nu K_\nu(\lambda);$$

If all the blocks $A_{[k,k]}$, $k = 1, \dots, s$ are non-resonant, then we will see that $K_\nu = 0$.

Corollary 6.1. *If Λ has pairwise distinct eigenvalues, system (1.1) is strongly isomonodromic if and only if (6.18)-(6.19) are satisfied.*

Proof of Corollary 6.1. At point 3. the non-resonance condition always holds if Λ has pairwise distinct eigenvalues $\lambda_1, \dots, \lambda_n$, so that $A_{[k,k]} = A_{kk}$, $k = 1, \dots, n$. \square

Proof of Theorem 6.2.

1. A computation gives

$$\frac{\partial^2 A}{\partial \lambda_i \partial \lambda_j} - \frac{\partial^2 A}{\partial \lambda_j \partial \lambda_i} \stackrel{(6.19)}{=} [\partial_i \tilde{\omega}_j - \partial_j \tilde{\omega}_i - [\tilde{\omega}_i, \tilde{\omega}_j], A], \quad 1 \leq i \neq j \leq s.$$

From Lemma 6.2, if (6.20) holds, we receive $\partial_i \tilde{\omega}_j - \partial_j \tilde{\omega}_i - [\tilde{\omega}_i, \tilde{\omega}_j] = 0$, which is (6.12). Therefore $\partial_i \partial_j A - \partial_j \partial_i A = 0$, so that (6.19) is integrable.

2. Equations (6.11) are automatically satisfied by the structure of the $\tilde{\omega}_j$ as in (6.14). In 1. we have seen that (6.12) holds. Now, (6.11)-(6.12) and (6.18)-(6.19) are the Frobenius integrability conditions of (6.21). The last statement at point 2. is proved by Proposition 3.2.

3. By (6.20), system (6.22) is integrable and admits a holomorphic fundamental matrix solution, call it $\mathfrak{T}(\lambda)$. Any fundamental matrix solution is $T(\lambda) = \mathfrak{T}(\lambda)T_0$, for an invertible constant matrix T_0 . Take any such $T(\lambda)$. Using (6.22), we receive

$$d(T^{-1}A^D T) = T^{-1} \left(dA^D + [A^D, \mathcal{D}] \right) T.$$

By Lemma 6.2, $dA^D + [A^D, \mathcal{D}] = 0$. Consequently,

$$d(T^{-1}A^D T) = 0.$$

So, $T^{-1}A^D T$ is constant. Therefore, there is a choice of T_0 such that

$$\mathcal{T}^{-1}(\lambda)A^D(\lambda)\mathcal{T}(\lambda) = J \quad \text{Jordan constant.}$$

Since A^D is block diagonal, we can take a block diagonal fundamental matrix solution $\mathfrak{T}(\lambda) = \mathfrak{T}_1(\lambda) \oplus \dots \oplus \mathfrak{T}_s(\lambda)$. Therefore, also T_0 can be taken block diagonal, and so is $\mathcal{T}(\lambda)$.

4. Since Assumption 3 holds by point 3. and J is constant, there are fundamental matrices $Y_\nu(z, \lambda)$ as in (4.9) satisfying

$$\frac{dY_\nu}{dz} = \left(\Lambda + \frac{A(\lambda)}{z} \right) Y_\nu,$$

which are holomorphic in $\mathcal{R}(\overline{\mathbb{C}} \setminus \{0, \infty\}) \times \mathbb{D}$. Therefore,

$$\varphi_\nu(z, \lambda) := d_\lambda Y_\nu - \sum_{j=1}^s (zE_{p_j} + \tilde{\omega}_j) d\lambda_j Y_\nu,$$

is well defined, where d_λ is the differential w.r.t. $\lambda_1, \dots, \lambda_s$. Using $d_\lambda \partial_z = \partial_z d_\lambda$ and (6.19) we obtain

$$\frac{\partial \varphi_\nu}{\partial z} = \left(\Lambda + \frac{A(\lambda)}{z} \right) \varphi_\nu + \sum_{j=1}^s \left([A, E_{p_j}] + [\Lambda, \tilde{\omega}_j] \right) d\lambda_j Y_\nu.$$

By (6.18), the above reduces to

$$\frac{\partial \varphi_\nu}{\partial z} = \left(\Lambda + \frac{A(\lambda)}{z} \right) \varphi_\nu.$$

Therefore, there is on \mathbb{D} a holomorphic matrix 1-form $K_\nu(\lambda)$ (not necessarily invertible) such that

$$\varphi_\nu = Y_\nu K_\nu.$$

We rewrite (4.9) with the specific Levelt form as in Section 5:

$$Y_\nu(z, \lambda) = \mathcal{T}(\lambda) \hat{Y}_\nu(z, \lambda) z^\Delta z^{N(\lambda)} e^{\Lambda z}.$$

From point 2. (constancy of J) we know already that Δ is constant. Nevertheless, we will indicate $d\Delta$ in the following, also if it is zero. We compute the structure of

$$K_\nu = Y_\nu^{-1} d_\lambda Y_\nu - Y_\nu^{-1} \sum_{j=1}^s (zE_{p_j} + \tilde{\omega}_j) d\lambda_j Y_\nu.$$

We have

$$\begin{aligned} & Y_\nu^{-1} d_\lambda Y_\nu = \\ & = e^{-\Lambda z} \left(z^{-N} z^{-\Delta} (\mathcal{T} \hat{Y}_\nu)^{-1} d_\lambda (\mathcal{T} \hat{Y}_\nu) z^\Delta z^N + z^{-N} d\Delta \ln z z^N + z^{-N} \sum_{k=1}^{\bar{k}} \frac{dN^k}{k!} (\ln z)^k z^N + z d\Lambda \right) e^{\Lambda z}. \end{aligned}$$

Notice that $d\Lambda = \sum_j E_{p_j} d\lambda_j$, so that

$$\begin{aligned} e^{\Lambda z} K_\nu e^{-\Lambda z} & = z^{-N} z^{-\Delta} \left[(\mathcal{T} \hat{Y}_\nu)^{-1} \left(d_\lambda (\mathcal{T} \hat{Y}_\nu) - \sum_j (zE_{p_j} + \tilde{\omega}_j) d\lambda_j \mathcal{T} \hat{Y}_\nu \right) \right] z^\Delta z^N \\ & + z^{-N} \left(d\Delta \ln z + \sum_{k=1}^{\bar{k}} \frac{dN^k}{k!} (\ln z)^k \right) z^N + \sum_j z E_{p_j} d\lambda_j. \end{aligned}$$

Notice that $[\Delta, d\Delta] = 0$ (also in case $d\Delta \neq 0$, because Δ is diagonal), and recall that $[\Delta, \Lambda] = [\Delta, d\Lambda] = 0$ and $[N, \Lambda] = [N, d\Lambda] = 0$, so that

$$\begin{aligned} z^\Delta z^N e^{\Lambda z} K_\nu e^{-\Lambda z} z^{-N} z^{-\Delta} &= (\mathcal{T}\hat{Y}_\nu)^{-1} \left(d_\lambda(\mathcal{T}\hat{Y}_\nu) - \sum_j (zE_{p_j} + \tilde{\omega}_j) d\lambda_j \mathcal{T}\hat{Y}_\nu \right) + \\ &+ d\Delta \ln z + z^\Delta \left(\sum_{k=1}^{\bar{k}} \frac{dN^k}{k!} (\ln z)^k \right) z^{-\Delta} + \sum_j zE_{p_j} d\lambda_j. \end{aligned}$$

The asymptotic expansion (4.10) holds in a sector \mathcal{S}_ν of amplitude greater than π . In such a sector, we obtain

$$\begin{aligned} z^\Delta z^N e^{\Lambda z} K_\nu e^{-\Lambda z} z^{-N} z^{-\Delta} &= - \sum_j zE_{p_j} d\lambda_j + \mathcal{T}^{-1} d\mathcal{T} + \mathcal{T}^{-1} \left(\sum_j [\mathcal{T}F_1 \mathcal{T}^{-1}, E_{p_j}] d\lambda_j \right) \mathcal{T} + \\ &- \mathcal{T}^{-1} \left(\sum_j \omega_j d\lambda_j \right) \mathcal{T} - \mathcal{T}^{-1} \mathcal{D}\mathcal{T} + \sum_j zE_{p_j} d\lambda_j + \\ &+ d\Delta \ln z + z^\Delta \left(\sum_{k=1}^{\bar{k}} \frac{dN^k}{k!} (\ln z)^k \right) z^{-\Delta} + O\left(\frac{1}{z}\right). \end{aligned}$$

In the computations in the proof of Theorem 6.1 we have seen that $[\mathcal{T}F_1 \mathcal{T}^{-1}, E_{p_j}] = \omega_j$. By point 2., $d\mathcal{T} = \mathcal{D}(\lambda)\mathcal{T}$. Therefore

$$z^\Delta z^N e^{\Lambda z} K_\nu e^{-\Lambda z} z^{-N} z^{-\Delta} = d\Delta \ln z + z^\Delta \left(\sum_{k=1}^{\bar{k}} \frac{dN^k}{k!} (\ln z)^k \right) z^{-\Delta} + O\left(\frac{1}{z}\right).$$

The off-diagonal blocks of the r.h.s. are of order $O(1/z)$, because N and Δ are block-diagonal. The l.h.s. is

$$e^{(\lambda_a - \lambda_b)z} z^{\Delta_a} z^{N_{[a,a]}} K_{[a,b]}^{(\nu)} z^{-N_{[b,b]}} z^{-\Delta_b}, \quad a \neq b = 1, \dots, s.$$

Since $e^{(\lambda_a - \lambda_b)z}$ diverges exponentially in a subsector of \mathcal{S}_ν , while the r.h.s. does not, necessarily

$$K_{[a,b]}^{(\nu)} = 0, \quad a \neq b = 1, \dots, s.$$

The diagonal blocks are

$$z^{\Delta_a} z^{N_{[a,a]}} K_{[a,a]}^{(\nu)} z^{-N_{[a,a]}} z^{-\Delta_a} = d\Delta_a \ln z + z^{\Delta_a} \left(\sum_{k=1}^{\bar{k}} \frac{dN_{[a,a]}^k}{k!} (\ln z)^k \right) z^{-\Delta_a} + O\left(\frac{1}{z}\right).$$

Namely

$$\begin{aligned} z^{\Delta_a} K_{[a,a]}^{(\nu)} z^{-\Delta_a} + z^{\Delta_a} ([N_{[a,a]}, K_{[a,a]}^{(\nu)}] \ln z + \text{terms in } (\ln z)^r \text{ with } r \geq 2) z^{-\Delta_a} &= \\ = d\Delta_a \ln z + z^{\Delta_a} (dN_{[a,a]} \ln z + \text{terms in } (\ln z)^r, r \geq 2) z^{-\Delta_a} + O(1/z). \end{aligned}$$




Figure 1: The structure of the sub-blocks \mathcal{K}_q of $K_{[a,a]}^{(\nu)}$ corresponding to \mathcal{N}_q , \mathcal{D}_q of figure 3 and $\Sigma_q = \sigma_q I_q$. They are split into sub-blocks.

Therefore, it is necessary that

$$z^{\Delta_a} K_{[a,a]}^{(\nu)} z^{-\Delta_a} = O(1/z). \quad (6.23)$$

Recall that $\Delta = D + \Sigma$. Proceeding as in the Appendix, we see that $N_{[a,a]}$ has diagonal block structure, and correspondingly so has $K_{[a,a]}^{(\nu)}$. Let for simplicity $\mathcal{K} := K_{[a,a]}^{(\nu)}$ for a fixed a . Then $\mathcal{K} = \mathcal{K}_1 \oplus \dots \oplus \mathcal{K}_\ell$, as in the Appendix. Since the integer diagonal entries in \mathcal{D}_q , $q = 1, \dots, \ell$ are an increasing sequence as in figure 3, each diagonal sub-block \mathcal{K}_q has structure as in figure 1.

In case of no resonance in $A_{[a,a]}$, then $\mathcal{D}_q = d_q I_q$ in figure 3, where d_q is integer and I_q is an identity matrix of suitable dimension. Thus, the diagonal sub-blocks of (6.23) reduce to

$$\mathcal{K}_q = O(1/z), \quad q = 1, \dots, \ell.$$

This implies that $K_{[a,a]}^{(\nu)} = 0$. □

7 The 3-dimensional case

Let $n = 3$. The isomonodromy problem of the case with no coalescences, namely $\Lambda = \text{diag}((\lambda_1, \lambda_2, \lambda_3), \lambda_i \neq \lambda_j)$, is highly transcendental. For example, for a certain class of matrices $A(\lambda)$, the isomonodromy deformation equations (1.6) are equivalent to the sixth Painlevé equation [23, 17, 27, 5, 15]. Coalescences of pairs of eigenvalues correspond to the fixed singularities of the sixth Painlevé equation.

The opposite situation is the trivial case $\Lambda = \text{diag}(\lambda_1, \lambda_1, \lambda_1)$. All the fundamental matrix solutions of system (1.1) are

$$Y(z, \lambda_1) = e^{z\lambda_1} z^{A(\lambda_1)} C(\lambda_1), \quad \det C(\lambda_1) \neq 0.$$

The system is isomonodromic if and only if A is constant.

The only non-trivial case we need to consider along a stratum of the coalescence locus is, up to permutation,

$$\frac{dY}{dz} = \left(\begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_2 \end{pmatrix} + \frac{A}{z} \right) Y. \quad (7.1)$$

The gauge $Y = e^{\lambda_2 z} \tilde{Y}$ and the change of variable $\zeta = xz$, where $x := \lambda_1 - \lambda_2$, yield

$$\frac{d\tilde{Y}}{d\zeta} = \left(\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \frac{A}{\zeta} \right) \tilde{Y}. \quad (7.2)$$

Suppose that A is constant. Then the above admits fundamental matrix solutions with constant essential monodromy data. Thus, the fundamental matrix solutions of the starting system also have constant data, if x varies in a sufficiently small domain away from $x = 0$. Therefore, the starting system is strongly isomonodromic.

We can obtain this result also from the point of view of Theorem 6.1. Since the diagonal blocks $A_{[1,1]} \equiv A_{11}$ and $A_{[2,2]}$ are always constant in the isomonodromic case, we are allowed to consider an isomonodromic deformation with

$$\mathcal{T}(\lambda) = \mathcal{T}_0 \mathfrak{B}(\lambda), \quad \mathfrak{B}(\lambda) = \mathfrak{B}_1(\lambda) \oplus \mathfrak{B}_2(\lambda), \quad [\mathfrak{B}_j(\lambda), J_j] = 0, \quad j = 1, 2,$$

where \mathcal{T}_0 is constant. We can also choose $\mathfrak{B}(\lambda) \equiv \mathfrak{B}(\lambda_1 - \lambda_2)$. This implies that

$$\frac{\partial \mathcal{T}}{\partial \lambda_2} = -\frac{\partial \mathcal{T}}{\partial \lambda_1}.$$

Moreover

$$\omega_1 = \frac{1}{\lambda_1 - \lambda_2} \begin{pmatrix} 0 & A_{12} & A_{13} \\ A_{21} & 0 & 0 \\ A_{31} & 0 & 0 \end{pmatrix}, \quad \omega_2 = -\omega_1.$$

Since $\tilde{\omega}_1 + \tilde{\omega}_2 = 0$, the same arguments of Remark 6.2 imply that

$$A = A(x), \quad x = \lambda_1 - \lambda_2,$$

so that $\omega_j = \omega_j(x)$. Therefore, the gauge $Y = e^{\lambda_2 z} \tilde{Y}$ transforms the Pfaffian system (6.1) into

$$d\tilde{Y} = \left\{ \left[\begin{pmatrix} x & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \frac{A}{z} \right] dz + \left(zE_1 + \tilde{\omega}_1(x) \right) dx \right\} \tilde{Y}, \quad \tilde{\omega}_1(x) = \omega_1(x) + \frac{d\mathcal{T}(x)}{dx} \mathcal{T}(x)^{-1},$$

where $E_1 = \text{diag}(1, 0, 0)$. Its integrability condition (6.3) reduces to

$$\frac{dA}{dx} = \left[\omega_1(x) + \frac{d\mathcal{T}(x)}{dx} \mathcal{T}(x)^{-1}, A \right]. \quad (7.3)$$

Now, a suitable choice of $\mathcal{T}(x)$ can be made such that

$$\left[\omega_1(x) + \frac{d\mathcal{T}(x)}{dx} \mathcal{T}(x)^{-1}, A \right] = 0. \quad (7.4)$$

This choice of \mathcal{T} is obtained by setting $A = A_0$ constant, calculating by linear algebra a constant \mathcal{T}_0 which Jordanizes $A_{11}^{(0)} \oplus A_{[2,2]}^{(0)}$:

$$\mathcal{T}_0^{-1}(A_{11}^{(0)} \oplus A_{[2,2]}^{(0)})\mathcal{T}_0 = J_1 \oplus J_2,$$

and the general $\mathfrak{B}(x)$ such that $[\mathfrak{B}_j(x), J_j] = 0$. The so obtained $\mathcal{T}(x) = \mathcal{T}_0\mathfrak{B}(x)$ must be substituted into (7.4), which can be solved as a system of differential equations for the entries of $\mathfrak{B}(x)$.

Example. We consider for simplicity matrices of the form

$$A = \begin{pmatrix} 0 & A_{12} & A_{13} \\ A_{21} & 0 & A_{23} \\ A_{31} & A_{32} & 0 \end{pmatrix}.$$

In the isomonodromic case, A_{23} and A_{32} are constant, due to constancy of the diagonal blocks of A . To simplify computations, we suppose that $A_{23}A_{32} \neq 0$, so that the block $A_{[2,2]}$ is diagonalizable, with eigenvalues $\pm\sqrt{A_{32}A_{23}}$. Thus, the general form of $\mathcal{T}(x)$ is

$$\mathcal{T}(x) = \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{A_{23}}{\sqrt{A_{23}A_{32}}} & -\frac{A_{23}}{\sqrt{A_{23}A_{32}}} \\ 0 & 1 & 1 \end{pmatrix}}_{=: \mathcal{T}_0^{\text{particular}}} \cdot \underbrace{\begin{pmatrix} a(x) & 0 & 0 \\ 0 & b(x) & 0 \\ 0 & 0 & c(x) \end{pmatrix}}_{\mathfrak{B}(x)}, \quad (7.5)$$

where $\mathcal{T}_0^{\text{particular}}$ is a particular choice for a constant matrix diagonalizing the block-diagonal part of A . Equation (7.4) has solution

$$b(x) = b_0 x^\rho a(x), \quad c(x) = c_0 x^{-\rho} a(x), \quad b_0, c_0 \in \mathbb{C} \setminus \{0\}, \quad \rho := \sqrt{A_{32}A_{23}}. \quad (7.6)$$

With this choice of $\mathcal{T}(x)$, equation (7.3) becomes

$$\frac{dA}{dx} = 0 \quad \implies \quad A \text{ is constant.} \quad (7.7)$$

Notice that this introduces the integration constants $A_{12}, A_{13}, A_{21}, A_{31}$. If instead we choose a constant \mathcal{T} , namely

$$\mathcal{T} = \mathcal{T}_0 := \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{A_{23}}{\sqrt{A_{23}A_{32}}} & -\frac{A_{23}}{\sqrt{A_{23}A_{32}}} \\ 0 & 1 & 1 \end{pmatrix} \cdot \begin{pmatrix} a_0 & 0 & 0 \\ 0 & b_0 & 0 \\ 0 & 0 & c_0 \end{pmatrix} \quad \text{constant}, \quad (7.8)$$

then equation (7.3) has solution given by a non-constant $A(x)$. Indeed, now A must satisfy

$$\frac{dA(x)}{dx} = [\omega_1(x), A(x)] \equiv \begin{pmatrix} 0 & \frac{A_{32}}{x} A_{13}(x) & \frac{A_{23}}{x} A_{12}(x) \\ -\frac{A_{23}}{x} A_{31}(x) & 0 & 0 \\ -\frac{A_{32}}{x} A_{21}(x) & 0 & 0 \end{pmatrix}.$$

Since A_{23} and A_{32} are constant, the above is a linear system with Fuchsian singularity at $x = 0$, with general solution

$$\begin{aligned} A_{12}(x) &= c_1 x^\rho + c_2 x^{-\rho}, & A_{13}(x) &= \sqrt{\frac{A_{23}}{A_{32}}}(c_1 x^\rho - c_2 x^{-\rho}), \\ A_{21}(x) &= c_3 x^\rho + c_4 x^{-\rho}, & A_{31}(x) &= -\sqrt{\frac{A_{32}}{A_{23}}}(c_3 x^\rho - c_4 x^{-\rho}). \end{aligned}$$

Here, c_1, c_2, c_3, c_4 are the integration constants. It is a computation to check that the above $A(x)$ has structure

$$A(x) = \mathcal{T}_0(\mathcal{T}^{-1}(x)A_0\mathcal{T}(x))\mathcal{T}_0^{-1},$$

as predicted by the discussion leading to (6.4), where A_0 is a constant, i.e. a solution of (7.7), \mathcal{T}_0 is (7.8) and $\mathcal{T}(x)$ is (7.5) with the functions (7.6).

Remark 7.1. There is a precise correspondence, that we describe below, between constant monodromy data of (7.1) and of (7.2). We consider the case when A is *constant*. This can always be achieved by a gauge transformation, as explained before. In order to simplify computations, we further assume that A is diagonalizable and non-resonant, so that (2.1) is

$$Y^{(0)}(z, \lambda) = G^{(0)}(x) \left(I + \sum_{k=1}^{\infty} F_k^{(0)}(\lambda) z^k \right) z^{J^{(0)}}, \quad J^{(0)} = \text{diag}(\mu_1, \mu_2, \mu_3),$$

and assume that $A_{[2,2]}$ is non-resonant (i.e. $\pm\sqrt{A_{32}A_{23}}$ not a half integer), so that (4.5) is

$$Y_F(z, \lambda) = \mathcal{T}(x) \left(I + \sum_{j=1}^{\infty} F_j(\lambda) z^{-j} \right) z^{\text{diag}(0, \rho, -\rho)} e^{\Lambda z}.$$

Take $\mathcal{T}(x)$ in (7.5), where $\mathcal{T}_0^{\text{particular}} = \mathcal{T}_0$ has structure (7.8) and $\mathcal{B}(x) = \text{diag}(1, x^\rho, x^{-\rho})$. Then, $dG = \sum_{j=1}^2 (\omega_j + D_j) d\lambda_j G$ in Theorem 3.1, reduces to

$$\frac{dG}{dx} = \left(\omega_1(x) + \frac{d\mathcal{T}(x)}{dx} \mathcal{T}(x)^{-1} \right) G \equiv \frac{A}{x} G,$$

so that

$$G^{(0)}(x) = G_0^{(0)} x^{J^{(0)}}, \quad \text{with constant } G_0^{(0)} \text{ such that } (G_0^{(0)})^{-1} A G_0^{(0)} = J^{(0)}.$$

Then, it is easy to check that the essential monodromy data of system (7.1) with the solutions $Y^{(0)}(z, \lambda)$, $Y_F(z, u)$ above and an admissible direction $\arg z = \tau$ are the same data of the system (7.2) with admissible direction $\arg \zeta = \tau + \arg(\lambda_1 - \lambda_2)$, relative to the fundamental matrix solutions

$$\tilde{Y}^{(0)}(\zeta) = G_0^{(0)} \left(I + \sum_{k=1}^{\infty} \tilde{F}_k^{(0)} \zeta^k \right) \zeta^{J^{(0)}}, \quad \tilde{Y}_F(\zeta) = \mathcal{T}_0 \left(I + \sum_{j=1}^{\infty} \tilde{F}_j \zeta^{-j} \right) \zeta^{\text{diag}(0, \rho, -\rho)} e^{\zeta \cdot \text{diag}(1, 0, 0)},$$

where the matrices $\tilde{F}_k^{(0)}$, \tilde{F}_j are constant.

8 A few words on the 4-dimensional case

The 3-dimensional case is rigid, namely one can reduce to a constant A . The first non-trivial isomonodromic deformations occur with dimension 4, and are already highly transcendental. Apart from the trivial situation when Λ only has one eigenvalue and A is necessarily constant, the cases we need to consider are (up to permutation)

$$\text{Case (1): } \Lambda = \text{diag}(\underbrace{\lambda_1, \lambda_1}, \lambda_2, \lambda_3)$$

$$\text{Case (2): } \Lambda = \text{diag}(\underbrace{\lambda_1, \lambda_1, \lambda_1}, \lambda_2)$$

$$\text{Case (3): } \Lambda = \text{diag}(\underbrace{\lambda_1, \lambda_1}, \underbrace{\lambda_2, \lambda_2})$$

It can be proved analogously to the 3-dimensional case that cases (2) and (3) are rigid. Heuristically, we can see this by the gauge transformation $Y = e^{\lambda_1 z} \tilde{Y}$ and the change of variable $\zeta = (\lambda_2 - \lambda_1)z$, which respectively yield in the two cases

$$\frac{d\tilde{Y}}{d\zeta} = \left(\text{diag}(0, 0, 0, 1) + \frac{A}{\zeta} \right) \tilde{Y} \quad \text{and} \quad \frac{d\tilde{Y}}{d\zeta} = \left(\text{diag}(0, 0, 1, 1) + \frac{A}{\zeta} \right) \tilde{Y}.$$

Case (1) is non-trivial. Heuristically, we understand this by the gauge transformation $Y = e^{\lambda_1 z} \tilde{Y}$ and the change of variable $\zeta = (\lambda_3 - \lambda_1)z$, which yield

$$\frac{d\tilde{Y}}{d\zeta} = \left(\text{diag}(0, 0, x, 1) + \frac{A}{\zeta} \right) \tilde{Y}, \quad x := \frac{\lambda_2 - \lambda_1}{\lambda_3 - \lambda_1}. \quad (8.1)$$

The matrix $A(\lambda)$ must satisfy the isomonodromy deformation equations (1.6), which we show below to be highly transcendental.

To simplify the treatment, we consider a case when A is much simpler than the general situation. Up to a gauge transformation, we can assume that

$$\tilde{\omega}_j = \omega_j \quad \text{for all } j.$$

In general, for every n and s , we have seen in Section 6 that for an isomonodromic system the relations $[\Lambda, \omega_j] = [E_{p_j}, A]$, $j = 1, \dots, s$, hold. This, together with our assumption that $\omega_{[k,k]}^{(j)} = 0$, $\forall 1 \leq j, k \leq s$, yields

$$\sum_{j=1}^s \omega_j = 0, \quad \sum_{j=1}^s \lambda_j \omega_{[k,\ell]}^{(j)} = A_{[k,\ell]} \delta_{k\ell}, \quad \forall 1 \leq k, \ell \leq s.$$

Hence, the isomonodromy deformation equations (1.6) imply

$$\sum_{j=1}^s \frac{\partial A}{\partial \lambda_j} = 0, \quad \sum_{j=1}^s \lambda_j \frac{\partial A_{[k,\ell]}}{\partial \lambda_j} = A_{[k,\ell]} A_{[\ell,\ell]} - A_{[k,k]} A_{[k,\ell]}, \quad \forall 1 \leq k, \ell \leq s. \quad (8.2)$$

Successively, we assume that

$$A_{[j,j]} = 0 \quad \text{for all } j = 1, \dots, s,$$

so that (8.2) becomes

$$\sum_{j=1}^s \frac{\partial A}{\partial \lambda_j} = 0, \quad \sum_{j=1}^s \lambda_j \frac{\partial A_{[k,\ell]}}{\partial \lambda_j} = 0 \quad \forall 1 \leq k, \ell \leq s. \quad (8.3)$$

The above implies the functional dependence

$$A = A \left(\left\{ \frac{\lambda_j - \lambda_1}{\lambda_s - \lambda_1} \right\}_{j=2}^{s-1} \right).$$

In case (1), $s = 3$ and $n = 4$, the functional dependence reduces to

$$A = A(x), \quad x = \frac{\lambda_2 - \lambda_1}{\lambda_3 - \lambda_1}.$$

The deformation equations (1.6) reduce to

$$\frac{dA}{dx} = [\hat{\omega}_2, A], \quad (8.4)$$

where $\hat{\omega}_2 = \hat{\omega}_2(x)$ is

$$\hat{\omega}_2 = \begin{bmatrix} \mathbf{0}_2 & \hat{\omega}_{[1,2]}^{(2)} & \vec{0} \\ \hat{\omega}_{[2,1]}^{(2)} & 0 & \hat{\omega}_{[2,3]}^{(2)} \\ \vec{0}^T & \hat{\omega}_{[3,2]}^{(2)} & 0 \end{bmatrix},$$

and

$$\vec{0} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \quad \vec{0}^T = [0 \ 0], \quad \mathbf{0}_2 := \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix},$$

$$\hat{\omega}_{[1,2]}^{(2)} = \frac{A_{[1,2]}}{x}, \quad \hat{\omega}_{[2,1]}^{(2)} = \frac{A_{[2,1]}}{x}, \quad \hat{\omega}_{[2,3]}^{(2)} = \frac{A_{[2,3]}}{x-1}, \quad \hat{\omega}_{[3,2]}^{(2)} = \frac{A_{[3,2]}}{x-1},$$

More explicitly, equations (8.4) are

$$\begin{cases} \frac{dA_{[1,2]}}{dx} = \frac{A_{[1,3]}A_{[3,2]}}{1-x}, & \frac{dA_{[2,1]}}{dx} = \frac{A_{[3,1]}A_{[2,3]}}{x-1}, \\ \frac{dA_{[2,3]}}{dx} = \frac{A_{[2,1]}A_{[1,3]}}{x}, & \frac{dA_{[3,2]}}{dx} = \frac{A_{[3,1]}A_{[1,2]}}{-x}, \\ \frac{dA_{[1,3]}}{dx} = \frac{A_{[1,2]}A_{[2,3]}}{x(1-x)}, & \frac{dA_{[3,1]}}{dx} = \frac{A_{[2,1]}A_{[3,2]}}{x(x-1)} \end{cases} \quad (8.5)$$

where⁷

$$A_{[1,2]} = \begin{bmatrix} A_{13} \\ A_{23} \end{bmatrix}, \quad A_{[1,3]} = \begin{bmatrix} A_{14} \\ A_{24} \end{bmatrix},$$

$$A_{[2,1]} = [A_{31} \ A_{32}], \quad A_{[2,3]} = A_{34},$$

$$A_{[3,1]} = [A_{41} \ A_{42}], \quad A_{[3,3]} = A_{43},$$

⁷Matrices are partitioned into the blocks inherited from $\Lambda = \text{diag}(\lambda_1, \lambda_1, \lambda_2, \lambda_3) = \lambda_1 I_2 \oplus \lambda_2 I_1 \oplus \lambda_3 I_1$.

Suppose now that A is skew-symmetric, so that we write it as

$$A = \begin{pmatrix} 0 & 0 & \phi_1 & \phi_3 \\ 0 & 0 & \phi_2 & \phi_4 \\ 0 & 0 & 0 & \phi_5 \\ 0 & 0 & 0 & 0 \end{pmatrix} - \begin{pmatrix} 0 & 0 & \phi_1 & \phi_3 \\ 0 & 0 & \phi_2 & \phi_4 \\ 0 & 0 & 0 & \phi_5 \\ 0 & 0 & 0 & 0 \end{pmatrix}^T.$$

Thus, system (8.5) becomes

$$\begin{cases} \frac{d\phi_1}{dx} = \frac{\phi_3\phi_5}{1-x}, & \frac{d\phi_3}{dx} = \frac{\phi_1\phi_5}{x(1-x)}, \\ \frac{d\phi_2}{dx} = \frac{\phi_4\phi_5}{1-x}, & \frac{d\phi_4}{dx} = \frac{\phi_2\phi_5}{x(1-x)}, \\ \frac{d\phi_5}{dx} = -\frac{\phi_1\phi_3 + \phi_2\phi_4}{x} \end{cases}$$

If we further restrict to the simpler case $\phi_1 = \phi_2$ and $\phi_3 = \phi_4$, we receive the three equations

$$\begin{cases} \frac{d\phi_1}{dx} = \frac{\phi_3\phi_5}{1-x}, & \frac{d\phi_3}{dx} = \frac{\phi_1\phi_5}{x(1-x)}, \\ \frac{d\phi_5}{dx} = -\frac{2\phi_1\phi_3}{x}. \end{cases}$$

Setting $\Omega_1 := -i\phi_5$, $\Omega_2 := \sqrt{2}\phi_1$, $\Omega_3 := i\sqrt{2}\phi_3$, the above system becomes

$$\frac{d\Omega_1}{dx} = \frac{\Omega_2\Omega_3}{x}, \quad \frac{d\Omega_2}{dx} = \frac{\Omega_1\Omega_3}{1-x}, \quad \frac{d\Omega_3}{dx} = \frac{\Omega_1\Omega_2}{x(x-1)}. \quad (8.6)$$

This system appears in [17, 18], where it is proved to be equivalent to the sixth Painlevé equation⁸ with parameters $\alpha \in \mathbb{C}$, $\beta = \gamma = 0$, $\delta = 1/2$.

We conclude that already in the very simplified case with A skew-symmetric depending on only three independent entries ϕ_1, ϕ_3, ϕ_5 , the isomonodromy problem is as transcendental as the Painlevé equations.

9 Example: the Caustic of a Dubrovin-Frobenius manifold in a generic case

In this section, we give an important example where the theory of non generic isomonodromy deformations along a stratum of a coalescence locus applies. It concerns the *caustic* of a *semisimple Dubrovin-Frobenius manifolds*. The result of this section is to show that our Theorems 6.1 and 6.2 are realized at generic points of the caustic.

⁸In classical form

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{1}{2} \left[\frac{1}{y} + \frac{1}{y-1} + \frac{1}{y-x} \right] \left(\frac{dy}{dx} \right)^2 - \left[\frac{1}{x} + \frac{1}{x-1} + \frac{1}{y-x} \right] \frac{dy}{dx} \\ &+ \frac{y(y-1)(y-x)}{x^2(x-1)^2} \left[\alpha + \beta \frac{x}{y^2} + \gamma \frac{x-1}{(y-1)^2} + \delta \frac{x(x-1)}{(y-x)^2} \right] \end{aligned}$$

We cannot explain here the details of Dubrovin-Frobenius manifolds, introduced by B. Dubrovin to give a geometrical formulation of 2-D topological field theories. The reader is referred to [16, 17]. For the geometry of the caustic, we refer to [24, 31, 32] and in particular to [28], in whose setting we will work.

In a Dubrovin-Frobenius manifold M of dimension n , the tangent spaces $T_p M$ at each point $p \in M$ is a Frobenius algebra with multiplication \circ , analytically depending on p . If this algebra is semisimple⁹ on an open dense subset of M , the manifold is called semisimple. The subset $\mathcal{K} \subset M$ where \circ is not semisimple is called *caustic*. This is empty or a hypersurface [24]. A flat metric η is defined¹⁰ on M , compatible with the product:

$$\eta(\mathbf{u} \circ \mathbf{v}, \mathbf{w}) = \eta(\mathbf{u}, \mathbf{v} \circ \mathbf{w}), \quad (9.1)$$

for any vector fields $\mathbf{u}, \mathbf{v}, \mathbf{w}$.

At semisimple points there is a basis of idempotent vector fields π_1, \dots, π_n , such that $\pi_i \circ \pi_j = \delta_{ij} \pi$, which are orthogonal with respect to η . They commute, $[\pi_i, \pi_j] = 0$, so that locally there are coordinates $u = (u_1, \dots, u_n)$ such that each $\pi_j = \frac{\partial}{\partial u_j}$ (with abuse of notation). They are called *canonical coordinates*, being uniquely determined (up to permutation) as eigenvalues of a multiplication operator $E \circ$, where E is a preferred global *Euler* vector field of weight 1. This is a field satisfying $\text{Lie}_E(\mathbf{u} \circ \mathbf{v}) - \text{Lie}_E(\mathbf{u}) \circ \mathbf{v} - \mathbf{u} \circ \text{Lie}_E(\mathbf{v}) = \mathbf{u} \circ \mathbf{v}$ (in short notation, $\text{Lie}_E \circ = \circ$) and $\text{Lie}_E(\eta) = (2 - d)\eta$, where $d \in \mathbb{C}$.

A sufficient condition for a point to be semisimple is that $u_a \neq u_b$ for all $1 \leq a \neq b \leq n$.

A Frobenius manifold is essentially characterized by a “ z -deformed” connection $\tilde{\nabla}$ defined by Dubrovin [16] as follows

$$\tilde{\nabla}_{\frac{d}{dz}} \mathbf{v} := \frac{\partial \mathbf{v}}{\partial z} + E \circ \mathbf{v} - \frac{1}{z} \hat{\mu}(\mathbf{v}), \quad \tilde{\nabla}_{\mathbf{u}} \mathbf{v} := \nabla_{\mathbf{u}} \mathbf{v} + z \mathbf{u} \circ \mathbf{v}, \quad z \in \mathbb{C}^* := \mathbb{C} \setminus \{0\}, \quad (9.2)$$

on the vector bundle $\pi^* TM|_{\mathbb{C}^* \times M} \longrightarrow \mathbb{C}^* \times M$, where $\pi : \mathbb{C}^* \times M \longrightarrow M$ is the natural projection, and $\mathbf{u}, \mathbf{v} \in (\pi^* \mathcal{T}_M)(\mathbb{C}^* \times M)$. Here $\hat{\mu}(\mathbf{v}) := (1 - d/2)\mathbf{v} - \nabla_{\mathbf{v}} E$, and ∇ is the Levi-Civita connection of η .

Locally, at semisimple points where all the (u_1, \dots, u_n) are pairwise distinct, the flatness (the zero curvature condition) of the above connection is an integrable Pfaffian system¹¹, which Dubrovin writes as

$$d\mathcal{Y} = \left[\left(U + \frac{V(u)}{z} \right) dz + \sum_{j=1}^n \left(z E_j + V_j(u) \right) du_j \right] \mathcal{Y}, \quad (9.3)$$

Here

$$U = \text{diag}(u_1, \dots, u_n),$$

⁹A point $p \in M$ is a semisimple if there are no *nilpotent* vectors \mathbf{v} in $T_p M$, namely no vectors such that $\mathbf{v} \circ^m = 0$, for some integer m .

¹⁰We mean a symmetric non-degenerate bilinear form, non necessarily positive definite.

¹¹It is obtained considering the connection $\tilde{\nabla}$ acting on the cotangent bundle, by looking at flat coordinates \tilde{t} satisfying $\tilde{\nabla} \tilde{t} = 0$. System (9.3) is the representation of $\tilde{\nabla} \tilde{t} = 0$ on the basis of normalized idempotents.

is the matrix representing E° both on the basis π_1, \dots, π_n and on the normalized basis

$$f_j := \frac{\pi_j}{\eta_j}, \quad \eta_j := \sqrt{\eta(\pi_j, \pi_j)}, \quad j = 1, \dots, n,$$

(for a chosen sign of the square root), V represents on the normalized basis the operator $\hat{\mu}$ and is skew-symmetric

$$V^T = -V,$$

$E_j = \partial U / \partial u_j$ has all zero entries except for 1 in position (j, j) , and V_j has entries

$$V_{aa}^{(j)}(u) = 0, \quad V_{ab}^{(j)}(u) = \frac{V_{ab}(u) (\delta_{aj} - \delta_{bj})}{u_a - u_b}, \quad a \neq b = 1, \dots, n. \quad (9.4)$$

The above (9.3) is a particular case of (6.1) when all eigenvalues remain pairwise distinct, and the matrices V_j are the analogous of our ω_j . The integrability of (9.3) is

$$\frac{\partial V}{\partial u_j} = [V_j, V], \quad j = 1, \dots, n, \quad (9.5)$$

which is a particular case of the isomonodromy deformation equations (6.19). By Corollary 6.1, they are necessary and sufficient conditions for the strong isomonodromy of the differential system

$$\frac{d\mathcal{Y}}{dz} = \left(U + \frac{V(u)}{z} \right) \mathcal{Y}. \quad (9.6)$$

The local structure of the manifold can be explicitly constructed at semisimple points where all the (u_1, \dots, u_n) are pairwise distinct, in terms of a fundamental matrix solution of (9.6) in Levelt form at $z = 0$ (see [16, 17, 18]).

In [14], we have shown that the coalescence of a pair of canonical coordinates $u_a - u_b \rightarrow 0$, for some $1 \leq a \neq b \leq n$, corresponds to a semisimple point of M if and only if $\lim_{u_a - u_b \rightarrow 0} V_{ab}(u) = 0$ holomorphically.

In general, for a pair $a \neq b \in \{1, \dots, n\}$, the condition $V_{ab}(u) \rightarrow 0$ is not satisfied along solutions of (9.5) when the corresponding $u_a - u_b \rightarrow 0$. Consequently, the entries of $V(u)$ may have a branching at $u_a - u_b = 0$, and may diverge along any direction $u_a - u_b \rightarrow 0$. Such coalescences of canonical coordinates may not correspond to a point of the manifold. In case they do, then the point must belong to the caustic.

Due to the singularity of $V(u)$ at coalescence points $u_a = u_b$, the canonical coordinates are not the good ones to describe the manifold close to the caustic. *The purpose of this section is to show that in suitable local coordinates on M the flatness of the connection (9.2) is represented by a Pfaffian system which, restricted to the caustic, is a system of type (6.1), satisfying Theorem 6.1, where Λ has repeated eigenvalues and \mathcal{T} is non trivial. Moreover, the above mentioned local coordinates, restricted at the caustic, are deformation parameters for a strong isomonodromy deformation described by our Theorems 6.1 and 6.2. This purpose is realized in Proposition 9.1 and Remark 9.4 below.*

A geometric study of the caustic of a Frobenius manifold has been done in [28], under the assumptions that on an open dense subset of \mathcal{K} there are $n - 1$ idempotent vector fields,

and the metric η is non-degenerate on \mathcal{K} . These are the assumption we will also make here. Such cases are realized in singularity theory, or equivalently for manifolds given by the orbit space of Coxeter groups (see [28]).

The points of \mathcal{K} in the above dense subset of [28] correspond to the coalescence of two canonical coordinates. Let $p \in \mathcal{K}$ belong to this subset and, without loss in generality, suppose that in a sufficiently small neighbourhood $\mathcal{B} \subset M$ of p the caustic \mathcal{K} is reached when

$$u_1 - u_2 \rightarrow 0.$$

It has been shown in [28] that theorems 2.11 and 4.7 (classification of 2-dimensional F manifolds) of [24] allow to conclude that in a neighbourhood of p the germ of the manifold is (notations are borrowed from and explained in [24])

$$(M, p) = (I_2(m), p) \times \underbrace{(A_1, p) \times \cdots \times (A_1, p)}_{n-2 \text{ times}}. \quad (9.7)$$

These means the following. Local coordinates

$$(t_1, t_2, u_3, \dots, u_n) \quad (9.8)$$

are defined in a sufficiently small neighbourhood $\mathcal{B} \subset M$ of $p \in \mathcal{K}$, with local basis of vector fields

$$\underbrace{\frac{\partial}{\partial t_1}, \frac{\partial}{\partial t_2}}_{T_p I_2(m)}, \quad \underbrace{\pi_3 = \frac{\partial}{\partial u_3}, \dots, \pi_n = \frac{\partial}{\partial u_n}}_{\mathbb{C}^{n-2}},$$

satisfying the multiplication table

$$\left\{ \begin{array}{l} \frac{\partial}{\partial t_1} \circ \frac{\partial}{\partial t_1} = \frac{\partial}{\partial t_1}, \quad \frac{\partial}{\partial t_1} \circ \frac{\partial}{\partial t_2} = \frac{\partial}{\partial t_2}, \quad \frac{\partial}{\partial t_2} \circ \frac{\partial}{\partial t_2} = t_2^{m-2} \frac{\partial}{\partial t_1}, \\ \pi_i \circ \pi_j = \delta_{ij} \pi_i, \\ \frac{\partial}{\partial t_1} \circ \pi_j = \frac{\partial}{\partial t_2} \circ \pi_j = 0 \end{array} \right. \quad \begin{array}{l} m \geq 2, \text{ integer} \\ i, j \geq 3. \end{array} \quad (9.9)$$

The caustic $\mathcal{B} \cap \mathcal{K}$ around p corresponds to

$$t_2 = 0,$$

and $\partial/\partial t_2$ is nilpotent at $\mathcal{B} \cap \mathcal{K}$.

For points in $\mathcal{B} \setminus \mathcal{K}$, the idempotents π_1, \dots, π_n and pairwise distinct canonical coordinates (u_1, \dots, u_n) are well defined. The Euler vector field and the unit of \circ on $\mathcal{B} \setminus \mathcal{K}$ are

$$E = \sum_{j=1}^n u_j \frac{\partial}{\partial u_j}, \quad e = \pi_1 + \pi_2 + \pi_3 + \cdots + \pi_n.$$

When $\mathcal{K} \cap \mathcal{B}$ is reached, then $u_1 - u_2 \rightarrow 0$ and the good coordinates become (9.8), which include u_3, \dots, u_n . On the whole \mathcal{B} , we can write [28]

$$E = t_1 \frac{\partial}{\partial t_1} + \frac{2}{m} t_2 \frac{\partial}{\partial t_2} + \sum_{j \geq 3} u_j \frac{\partial}{\partial u_j}, \quad e = \frac{\partial}{\partial t_1} + \pi_3 + \cdots + \pi_n, \quad (9.10)$$

so that $\frac{\partial}{\partial t_1} = \pi_1 + \pi_2$ is well defined also at the caustic. Moreover, $\pi_1 + \pi_2, \pi_3, \dots, \pi_n$ are the $n-1$ idempotents defined on $\mathcal{K} \cap \mathcal{B}$. Notice that E_\circ is diagonalizable on $\mathcal{K} \cap \mathcal{B}$, with a repeated eigenvalue $t_1 \equiv u_1 = u_2$ and pairwise distinct eigenvalues u_3, \dots, u_n .

So far we have reviewed the local description of the manifold at a generic point of the caustic, according to [28]. We now show that the flatness of the Dubrovin deformed connection in the coordinates (9.8) is realized at the caustic by a strongly isomonodromic theory obeying our Theorems 6.1 and 6.2.

Using the multiplication table (9.9), E_\circ is represented on the basis

$$\mathbf{w}_1 := \frac{\partial}{\partial t_1}, \quad \mathbf{w}_2 := \frac{\partial}{\partial t_2}, \quad \mathbf{w}_j := f_j = \frac{\pi_j}{\sqrt{\eta_j}} \quad j \geq 3,$$

by a matrix

$$\mathcal{U} = \begin{pmatrix} \hat{\mathcal{U}} & 0 \\ 0 & U_{n-2} \end{pmatrix}, \quad (9.11)$$

where

$$\hat{\mathcal{U}} = \begin{pmatrix} t_1 & \frac{2}{m} t_2^{m-1} \\ \frac{2}{m} t_2 & t_1 \end{pmatrix}, \quad U_{n-2} = \text{diag}(u_3, \dots, u_n).$$

In $\mathcal{B} \setminus \mathcal{K}$, let Ψ be a matrix such that

$$\Psi \mathcal{U} \Psi^{-1} = U = \text{diag}(u_1, u_2, \dots, u_n),$$

with normalization

$$\Psi^T \Psi = \left(\eta(\mathbf{w}_\alpha, \mathbf{w}_\beta) \right)_{\alpha, \beta=1}^n. \quad (9.12)$$

The eigenvalues u_1, u_2 of $\hat{\mathcal{U}}$ are immediately computed:

$$\begin{cases} u_1 = t_1 + \frac{2}{m} t_2^{m/2} \\ u_2 = t_1 - \frac{2}{m} t_2^{m/2} \end{cases} \implies \begin{cases} t_1 = \frac{u_1 + u_2}{2}, \\ t_2 = \left(\frac{m}{4} (u_1 - u_2) \right)^{2/m}. \end{cases} \quad (9.13)$$

Since system (9.6) represents the z component of the deformed connection (9.2) on the basis of normalized idempotents f_1, \dots, f_n in $\mathcal{B} \setminus \mathcal{K}$, and \mathcal{U} represents E_\circ on the basis $\mathbf{w}_1, \dots, \mathbf{w}_n$, then the gauge transformation

$$\mathcal{Y} = \Psi Y,$$

gives a differential system representing the z -component of (9.2) on the basis $\mathbf{w}_1, \dots, \mathbf{w}_n$:

$$\frac{dY}{dz} = \left(\mathcal{U} + \frac{\mathcal{V}}{z} \right) Y, \quad \mathcal{V} := \Psi^{-1} V \Psi \quad (9.14)$$

In other words, Ψ gives the change of basis

$$\left[\frac{\partial}{\partial t_1}, \frac{\partial}{\partial t_2}, f_3, \dots, f_n \right] = [f_1, f_2, f_3, \dots, f_n] \Psi. \quad (9.15)$$

System (9.14) is expressed in the good coordinates $(t_1, t_2, u_3, \dots, u_n)$. The manifold structure is analytic at \mathcal{K} in these coordinates. Therefore, the coefficients of (9.14) analytically extend from $\mathcal{B} \setminus \mathcal{K}$ to the whole \mathcal{B} , and in particular $\mathcal{V}(t_1, t_2, u_3, \dots, u_n)$ must be well defined and holomorphic also at the caustic $t_2 = 0$.

The matrix Ψ diagonalizing \mathcal{U} is partitioned into blocks as \mathcal{U} :

$$\Psi = \begin{pmatrix} \hat{\Psi} & 0 \\ 0 & I_{n-2} \end{pmatrix}, \quad \left\{ \begin{array}{l} \hat{\Psi} = \begin{pmatrix} \frac{a}{\sqrt{2}} t_2^{(2-m)/4} & \frac{a}{\sqrt{2}} t_2^{(m-2)/4} \\ \frac{ib}{\sqrt{2}} t_2^{(2-m)/4} & -\frac{ib}{\sqrt{2}} t_2^{(m-2)/4} \end{pmatrix} \\ I_{n-2} = n - 2 \text{ dimensional identity matrix} \\ a \text{ and } b \text{ functions of } (t_1, t_2, u_3, \dots, u_n) \end{array} \right. \quad (9.16)$$

The condition (9.12) implies

$$a^2 = \tilde{\eta}_{12} + t_2^{\frac{m-2}{2}} \tilde{\eta}_{11}, \quad b^2 = \tilde{\eta}_{12} - t_2^{\frac{m-2}{2}} \tilde{\eta}_{11}. \quad (9.17)$$

where

$$\tilde{\eta}_{11} := \eta \left(\frac{\partial}{\partial t_1}, \frac{\partial}{\partial t_1} \right), \quad \tilde{\eta}_{12} := \eta \left(\frac{\partial}{\partial t_1}, \frac{\partial}{\partial t_2} \right), \quad \tilde{\eta}_{22} := \eta \left(\frac{\partial}{\partial t_2}, \frac{\partial}{\partial t_2} \right).$$

Notice that $\tilde{\eta}_{11}$, $\tilde{\eta}_{22}$ are well defined at $t_2 = 0$, because so is the metric η and the manifold structure is analytic in the coordinates $(t_1, t_2, u_3, \dots, u_n)$.

By (9.16), \mathcal{U} is not holomorphically diagonalizable at $t_2 = 0$ for $m \geq 3$, in accordance with the fact that the change of coordinates $(u_1, u_2; u_3, \dots, u_n) \mapsto (t_1, t_2; u_3, \dots, u_n)$ is singular at \mathcal{K} as in (9.13). For $m = 2$, \mathcal{U} is holomorphically diagonalizable, which is an example of the semisimple coalescence studied in [14].

Remark 9.1. By (9.1), (9.9) and the unit e in (9.10), we receive

$$\tilde{\eta}_{11} = \eta_1 + \eta_2, \quad \tilde{\eta}_{22} = t_2^{m-2} \tilde{\eta}_{11}. \quad (9.18)$$

Let $m \geq 3$. The above (9.18) implies that $\tilde{\eta}_{22}|_{t_2=0} = 0$, and thus

$$\tilde{\eta}_{12}|_{t_2=0} \neq 0,$$

otherwise the metric would be degenerate at $t_2 = 0$, which cannot be by assumption. Now let $m = 2$. In case $\tilde{\eta}_{12}|_{t_2=0} = 0$, then $\tilde{\eta}_{11}|_{t_2=0} = \tilde{\eta}_{22}|_{t_2=0} \neq 0$, otherwise the metric would be degenerate at $t_2 = 0$.

Let $\mathcal{V}_{[1,1]}$ be the 2×2 upper left block of \mathcal{V} . A lengthy but elementary computation proves the implication

$$\left\{ \begin{array}{l} V + V^T = 0, \\ \mathcal{V} = \Psi^{-1} V \Psi \quad \text{and (9.17)} \end{array} \right. \implies \mathcal{V}_{[1,1]} = \frac{iV_{12}(u)}{\sqrt{\tilde{\eta}_{12}^2 - \tilde{\eta}_{11}^2 t_2^{m-2}}} \begin{pmatrix} \tilde{\eta}_{12} & \tilde{\eta}_{11} t_2^{m-2} \\ -\tilde{\eta}_{11} & -\tilde{\eta}_{12} \end{pmatrix}.$$

The above matrix has eigenvalues $\pm iV_{12}(u)$. As already explained, $\mathcal{V}_{[1,1]}$, $\tilde{\eta}_{12}$, $\tilde{\eta}_{11}$, $\tilde{\eta}_{22}$ are well defined at $t_2 = 0$, namely for $u_1 = u_2$, as analytic functions of $t_1 (\equiv u_1 = u_2)$, u_3, \dots, u_n . Hence, the limit for $u_1 - u_2 \rightarrow 0$ of $V_{12}(u)$ is well defined as an analytic function of t_1, u_3, \dots, u_n . Let

$$\mathring{V}_{12} := \lim_{u_1 - u_2 \rightarrow 0} V_{12}(u) \quad \text{in } \mathcal{B}. \quad (9.19)$$

In principle, it depends holomorphically on t_1, u_3, \dots, u_n , but we will show that \mathring{V}_{12} is *constant*. Keeping the above discussion into account, in $\mathcal{B} \cap \mathcal{K}$ we have

$$\mathcal{V}_{[1,1]}|_{t_2=0} = \begin{cases} \left. \frac{i\mathring{V}_{12}}{\sqrt{\tilde{\eta}_{12}^2}} \begin{pmatrix} \tilde{\eta}_{12} & 0 \\ -\tilde{\eta}_{11} & -\tilde{\eta}_{12} \end{pmatrix} \right|_{t_2=0}, & m \geq 3. \\ \left. \frac{i\mathring{V}_{12}}{\sqrt{\tilde{\eta}_{12}^2 - \tilde{\eta}_{11}^2}} \begin{pmatrix} \tilde{\eta}_{12} & \tilde{\eta}_{11} \\ -\tilde{\eta}_{11} & -\tilde{\eta}_{12} \end{pmatrix} \right|_{t_2=0}, & m = 2. \end{cases} \quad (9.20)$$

Using (9.13) and the chain rule $\frac{\partial \mathcal{Y}}{\partial u_k} = \frac{\partial \mathcal{Y}}{\partial t_1} \frac{\partial t_1}{\partial u_k} + \frac{\partial \mathcal{Y}}{\partial t_2} \frac{\partial t_2}{\partial u_k}$ for $k = 1, 2$, the du_j components ($j \in \{1, \dots, n\}$) of (9.3) become

$$\frac{\partial Y}{\partial t_1} = \left(z \underbrace{\Psi^{-1}(E_1 + E_2)\Psi}_{(E_1 + E_2)} + \Psi^{-1}(V_1 + V_2)\Psi - \Psi^{-1} \frac{\partial \Psi}{\partial t_1} \right) Y. \quad (9.21)$$

$$\frac{\partial Y}{\partial t_2} = \left(z t_2^{\frac{m-2}{2}} \Psi^{-1}(E_1 - E_2)\Psi + t_2^{\frac{m-2}{2}} \Psi^{-1}(V_1 - V_2)\Psi - \Psi^{-1} \frac{\partial \Psi}{\partial t_2} \right) Y, \quad (9.22)$$

$$\frac{\partial Y}{\partial u_j} = \left(z \underbrace{\Psi^{-1}E_j\Psi}_{E_j} + \Psi^{-1}V_j\Psi - \Psi^{-1} \frac{\partial \Psi}{\partial u_j} \right) Y, \quad j \geq 3. \quad (9.23)$$

with

$$z t_2^{\frac{m-2}{2}} \Psi^{-1}(E_1 - E_2)\Psi = z \begin{pmatrix} 0 & t_2^{m-2} \\ 1 & 0 \\ & & I_{n-2} \end{pmatrix}.$$

By the same chain rule, it is easily seen that from the deformation equations (9.5) we receive

$$\frac{\partial \mathcal{V}}{\partial t_1} = \left[\Psi^{-1}(V_1 + V_2)\Psi - \Psi^{-1} \frac{\partial \Psi}{\partial t_1}, \mathcal{V} \right], \quad (9.24)$$

$$\frac{\partial \mathcal{V}}{\partial t_2} = \left[t_2^{\frac{m-2}{2}} \Psi^{-1}(V_1 - V_2)\Psi - \Psi^{-1} \frac{\partial \Psi}{\partial t_2}, \mathcal{V} \right], \quad (9.25)$$

$$\frac{\partial \mathcal{V}}{\partial u_j} = \left[\Psi^{-1}V_j\Psi - \Psi^{-1} \frac{\partial \Psi}{\partial t_2}, \mathcal{V} \right], \quad j \geq 3. \quad (9.26)$$

The Pfaffian system given by (9.14) and (9.21)-(9.23) represents the flatness of the connection (9.2) in the good coordinates $(t_1, t_2, u_3, \dots, u_n)$ in \mathcal{B} . All the coefficients must be

holomorphic at $t_2 = 0$ in these variables. The flatness of the connection (9.2) restricted at $\mathcal{B} \cap \mathcal{K}$ is then represented by (9.14), (9.21) and (9.23) restricted at $t_2 = 0$, namely

$$\begin{cases} \frac{dY}{dz} = \left(\mathcal{U}|_{t_2=0} + \frac{\mathcal{V}|_{t_2=0}}{z} \right) Y, & \mathcal{U}|_{t_2=0} = \text{diag}(t_1, t_1; u_3, \dots, u_n), \\ \frac{\partial Y}{\partial t_1} = \left(z(E_1 + E_2) + \mathcal{V}_2|_{t_2=0} - \Psi^{-1} \frac{\partial \Psi}{\partial t_1} \Big|_{t_2=0} \right) Y, & \mathcal{V}_2 := \Psi^{-1}(V_1 + V_2)\Psi, \\ \frac{\partial Y}{\partial u_j} = \left(zE_j + \mathcal{V}_j|_{t_2=0} - \Psi^{-1} \frac{\partial \Psi}{\partial u_j} \Big|_{t_2=0} \right) Y, \quad j \geq 3, & \mathcal{V}_j := \Psi^{-1}V_j\Psi. \end{cases} \quad (9.27)$$

The deformation equation (9.24) and (9.26) become

$$\begin{cases} \frac{\partial \mathcal{V}}{\partial t_1} = \left[\mathcal{V}_2 - \Psi^{-1} \frac{\partial \Psi}{\partial t_1} \Big|_{t_2=0}, \mathcal{V} \right], \\ \frac{\partial \mathcal{V}}{\partial u_j} = \left[\mathcal{V}_j - \Psi^{-1} \frac{\partial \Psi}{\partial u_j} \Big|_{t_2=0}, \mathcal{V} \right], \quad j \geq 3. \end{cases} \quad (9.28)$$

Proposition 9.1. *The Pfaffian system (9.27), representing the flatness of the Dubrovin deformed connection (9.2) restricted at caustic $\mathcal{K} \cap \mathcal{B}$, is exactly a system of type (6.1), with*

$$s = n - 1, \quad p_1 = 2, \quad p_3 = \dots = p_{n-1} = 1, \quad (\lambda_1, \dots, \lambda_{n-1}) := (t_1, u_3, \dots, u_n).$$

and

$$\Lambda := \text{diag}(\lambda_1, \lambda_1, \underbrace{\lambda_2, \dots, \lambda_{n-1}}_{n-2 \text{ distinct}}) \equiv \mathcal{U}|_{t_2=0}, \quad A(\lambda) := \mathcal{V}|_{t_2=0} \quad \text{as in (9.20)}.$$

With the above identification, system (9.28) is exactly the isomonodromy deformation equations (6.3).

The block-diagonal matrix $\mathcal{T}(\lambda)$ which reduces to Jordan form the block diagonal part $A_{[1,1]} \oplus A_{33} \oplus \dots \oplus A_{nn}$ of $A(\lambda)$ is

$$\mathcal{T}(\lambda) = \underbrace{\mathcal{T}_1(\lambda)}_{2 \times 2} \oplus \text{diag}(h_2, \dots, h_{n-1}), \quad (9.29)$$

where $\mathcal{T}_1(\lambda)$ brings the upper left 2×2 block $A_{[1,1]}$ to Jordan form, while the other h_ℓ , $2 \leq \ell \leq n - 1$, are arbitrary scalar constants. In (9.27), one exactly has

$$-\Psi^{-1} \frac{\partial \Psi}{\partial t_1} \Big|_{t_2=0} = \frac{\partial \mathcal{T}}{\partial t_1} \mathcal{T}^{-1}, \quad -\Psi^{-1} \frac{\partial \Psi}{\partial u_j} \Big|_{t_2=0} = \frac{\partial \mathcal{T}}{\partial u_j} \mathcal{T}^{-1}, \quad j \geq 3, \quad (9.30)$$

and

$$\mathcal{V}_2|_{t_2} = \omega_1, \quad \mathcal{V}_j|_{t_2} = \omega_{j-1}, \quad j = 3, \dots, n.$$

The proof will be given after Remark 9.4. Notice that $A(\lambda) := \mathcal{V}|_{t_2=0}$ is holomorphic in $\mathcal{B} \cap \mathcal{K}$, and \mathcal{B} can be made sufficiently small so that Assumption 1 of the Introduction is satisfied.

Remark 9.2. According to the block structure $\Lambda = \lambda_1 I_2 \oplus (\lambda_2 I_1 \oplus \cdots \oplus \lambda_{n-1} I_1)$, we have

$$E_{p_1} = E_1 + E_2, \quad E_{p_j} = E_{j+1}, \quad 2 \leq j \leq n-1,$$

and the matrix A must be partitioned into the following blocks. $A_{[1,1]}$ is a 2×2 block. Let $j, k \in \{2, \dots, n-1\}$. Then, the $A_{[j,1]}$ are 1×2 blocks; the $A_{[1,k]}$ are 2×1 blocks; the $A_{[j,k]} = A_{j+1,k+1}$ are matrix entries (1 dimensional blocks!). The structure (9.29) is required by this block decomposition to fulfil Theorem 6.1.

Proposition 9.1 not only tells us that the isomonodromy deformation theory developed in this paper applies at the caustics, but also allow us to predict properties of the caustic itself, as in the following

Corollary 9.1. *The block $\mathcal{V}_{[1,1]}|_{t_2=0}$ is constant. In particular, $\mathring{V}_{12} = \lim_{u_1 \rightarrow u_2} V_{12}(u)$ defined in (9.19) is constant, and $\tilde{\eta}_{11}|_{t_2=0}$ and $\tilde{\eta}_{12}|_{t_2=0}$ are constant.*

Proof of Corollary 9.1. It follows from PART II of Theorem 6.1, which applies thanks to Proposition 9.1, and from (9.20). \square

Remark 9.3 (on the value of \mathring{V}_{12}). Remarkably, the value of the constant \mathring{V}_{12} can be written explicitly:

$$\mathring{V}_{12} = \frac{i(m-2)}{2m},$$

(recall that $m \geq 2$). This result was first observed in [28], and can be proved in three ways.

- 1) It is proved in [28] *by purely geometric arguments*, starting from the decomposition (9.7).
- 2) It is rather laborious to show analytically that the isomonodromy deformation equations (9.5) admit a solution $V(u)$ with the property that $\lim_{u_2 \rightarrow u_1} V_{12}(u)$ exists finite, if and only if this solution is such that in a neighbourhood of $u_1 - u_2 = 0$

$$V_{12}(u) = \frac{i\sigma}{2} + \sum_{(h,k) \neq (0,0)}^{\infty} b_{hk}(x_3, \dots, x_{n-1}) x_2^{h+(1-\sigma)k} + \sum_{(h,q) \neq (0,0)}^{\infty} a_{hq}(x_3, \dots, x_{n-1}) x_2^{h+(1+\sigma)k},$$

where

$$\sigma \in \mathbb{C}, \quad 0 \leq \Re \sigma < 1; \quad x_j = \frac{u_j - u_1}{u_n - u_1}, \quad j = 2, \dots, n-1,$$

and the functions $a_{hk}(x_3, \dots, x_{n-1}), b_{hk}(x_3, \dots, x_{n-1})$ are analytic around p . Hence,

$$\mathring{V}_{12} = \frac{i\sigma}{2} \quad \text{is constant.}$$

Then, using again equations (9.5) one can prove that $\mathring{V}_{12} = i(m-2)/2m$. This is more laborious than the geometric proof of [28].

- 3) Using the approach of this paper. We have seen that the coefficients of equations (9.22) and (9.25) must be holomorphic at $t_2 = 0$ in the variables $(t_1, t_2, u_3, \dots, u_n)$. We prove at the end of this Section that

$$t_2^{\frac{m-2}{2}} \Psi^{-1}(V_1 - V_2)\Psi - \Psi^{-1} \frac{\partial \Psi}{\partial t_2},$$

is holomorphic at $t_2 = 0$ (as it must be) if and only if $\mathring{V}_{12} = i(m-2)/2m$.

Remark 9.4 (On the strong isomonodromy at the caustic). From Remark 9.3 it follows that $0 \leq \Re(\mathring{V}_{12}) < 1/2$. Thus, $\mathcal{V}_{[1,1]}|_{t_2=0}$ is non resonant, so that by Theorem 6.2 system (9.28) is a *necessary and sufficient condition for strong isomonodromy*. Therefore, likewise the case of semisimple points, also at the generic points of the caustic here considered, the Dubrovin flat connection is realized by a strongly isomonodromic system.

Proof of Proposition 9.1. The $z(E_1 + E_2)$ and zE_j terms in (9.27) correctly display $E_{p_1} = E_1 + E_2$ and $E_{p_j} = E_{j+1}$, $2 \leq j \leq n-1$.

- First, we show that the matrices \mathcal{V}_j , $j = 2, \dots, n$, in (9.27) are

$$\mathcal{V}_2|_{t_2=0} = \omega_1(\lambda), \quad \mathcal{V}_j|_{t_2=0} = \omega_{j-1}(\lambda), \quad j = 3, \dots, n.$$

with the structure (6.2). We start by computing \mathcal{V}_2 in \mathcal{B} .

$$\begin{aligned} \mathcal{V}_2 &= \Psi^{-1}(V_1 + V_2)\Psi = \Psi^{-1} \begin{pmatrix} 0 & 0 & \dots & \frac{V_{1k}}{u_1 - u_k} & \dots \\ 0 & 0 & \dots & \frac{V_{2k}}{u_2 - u_k} & \dots \\ \vdots & \vdots & 0 & \dots & 0 & \dots & 0 \\ \frac{V_{k1}}{u_1 - u_k} & \frac{V_{k2}}{u_2 - u_k} & 0 & \dots & 0 & \dots & 0 \\ \vdots & \vdots & 0 & \dots & 0 & \dots & 0 \end{pmatrix} \Psi \\ &= \begin{pmatrix} 0 & 0 & \widehat{\Psi}^{-1} \begin{bmatrix} \dots & \frac{V_{1k}}{u_1 - u_k} & \dots \\ \dots & \frac{V_{2k}}{u_2 - u_k} & \dots \end{bmatrix} \\ 0 & 0 & \begin{bmatrix} \vdots & \vdots \\ \frac{V_{k1}}{u_1 - u_k} & \frac{V_{k2}}{u_2 - u_k} \\ \vdots & \vdots \end{bmatrix} \widehat{\Psi} & 0 & \dots & 0 & \dots & 0 \\ & & & 0 & \dots & 0 & \dots & 0 \\ & & & 0 & \dots & 0 & \dots & 0 \end{pmatrix} \end{aligned}$$

We want to take the limit for $t_2 \rightarrow 0$, namely for $u_1 - u_2 \rightarrow 0$. We use (from (9.13) and $V = \Psi\mathcal{V}\Psi^{-1}$) the relations

$$u_1 - u_2 = \frac{4}{m} t_2^{m/2}, \quad \begin{cases} V_{1k} = \frac{a}{\sqrt{2}} \left(t_2^{\frac{2-m}{4}} \mathcal{V}_{1k} + t_2^{\frac{m-2}{4}} \mathcal{V}_{2k} \right) \\ V_{2k} = \frac{ib}{\sqrt{2}} \left(t_2^{\frac{2-m}{4}} \mathcal{V}_{1k} - t_2^{\frac{m-2}{4}} \mathcal{V}_{2k} \right) \end{cases} \quad k = 3, \dots, n,$$

and receive

$$\widehat{\Psi}^{-1} \begin{bmatrix} \dots & \frac{V_{1k}}{u_1 - u_k} & \dots \\ \dots & \frac{V_{2k}}{u_2 - u_k} & \dots \end{bmatrix} = \begin{pmatrix} \dots & \frac{\mathcal{V}_{1k}}{2} \frac{u_1 + u_2 - 2u_k}{(u_1 - u_k)(u_2 - u_k)} - \frac{2 t_2^{m-1}}{m(u_1 - u_k)(u_2 - u_k)} \mathcal{V}_{2k} & \dots \\ \dots & \frac{\mathcal{V}_{2k}}{2} \frac{u_1 + u_2 - 2u_k}{(u_1 - u_k)(u_2 - u_k)} - \frac{2 t_2}{m(u_1 - u_k)(u_2 - u_k)} \mathcal{V}_{1k} & \dots \end{pmatrix}$$

$$\xrightarrow{t_2 \rightarrow 0, u_1 - u_2 \rightarrow 0} \begin{pmatrix} \dots & \frac{\mathcal{V}_{1k}|_{t_2=0}}{u_1 - u_k} & \dots \\ \dots & \frac{\mathcal{V}_{2k}|_{t_2=0}}{u_1 - u_k} & \dots \end{pmatrix} \equiv \begin{pmatrix} \dots & \frac{A_{1k}}{\lambda_1 - \lambda_{k-1}} & \dots \\ \dots & \frac{A_{2k}}{\lambda_1 - \lambda_{k-1}} & \dots \end{pmatrix}, \quad \forall m \geq 2.$$

In the last step we have used $t_1 = u_1 = u_2$ at the caustic, the definition $\lambda_1 := t_1$, $\lambda_{k-1} := u_k$, $k = 3, \dots, n$, and $A(\lambda) := \mathcal{V}|_{t_2=0}$. Successively, we use

$$\begin{cases} V_{k1} = \frac{1}{\sqrt{2} a} \left(t_2^{\frac{2-m}{4}} \mathcal{V}_{k2} + t_2^{\frac{m-2}{4}} \mathcal{V}_{k1} \right) \\ V_{k2} = \frac{i}{\sqrt{2} b} \left(t_2^{\frac{2-m}{4}} \mathcal{V}_{k2} - t_2^{\frac{m-2}{4}} \mathcal{V}_{k1} \right) \end{cases} \quad k = 3, \dots, n,$$

and find

$$\begin{bmatrix} \vdots & \vdots \\ \frac{V_{k1}}{u_1 - u_k} & \frac{V_{k2}}{u_2 - u_k} \\ \vdots & \vdots \end{bmatrix} \widehat{\Psi} =$$

$$= \begin{pmatrix} \vdots & \vdots \\ \frac{\mathcal{V}_{k1}}{2} \frac{u_1 + u_2 - 2u_k}{(u_1 - u_k)(u_2 - u_k)} - \frac{2 t_2 \mathcal{V}_{k2}}{m(u_1 - u_k)(u_2 - u_k)} & \frac{\mathcal{V}_{k2}}{2} \frac{u_1 + u_2 - 2u_k}{(u_1 - u_k)(u_2 - u_k)} - \frac{2 t_2^{m-1} \mathcal{V}_{k1}}{m(u_1 - u_k)(u_2 - u_k)} \\ \vdots & \vdots \end{pmatrix}$$

$$\xrightarrow{t_2 \rightarrow 0, u_1 - u_2 \rightarrow 0} \begin{pmatrix} \vdots & \vdots \\ \frac{\mathcal{V}_{k1}|_{t_2=0}}{u_1 - u_k} & \frac{\mathcal{V}_{k2}|_{t_2=0}}{u_1 - u_k} \\ \vdots & \vdots \end{pmatrix} \equiv \begin{pmatrix} \vdots & \vdots \\ \frac{A_{k1}}{\lambda_1 - \lambda_{k-1}} & \frac{A_{k2}}{\lambda_1 - \lambda_{k-1}} \\ \vdots & \vdots \end{pmatrix}, \quad \forall m \geq 2.$$

Notice that the above computations confirm that \mathcal{V}_2 is holomorphic in a neighbourhood of

$t_2 = 0$. In conclusion, we have found that

$$\mathcal{V}_2|_{t_2=0} = \begin{pmatrix} 0 & 0 & \cdots & \frac{A_{1k}}{\lambda_1 - \lambda_{k-1}} & \cdots \\ 0 & 0 & \cdots & \frac{A_{2k}}{\lambda_1 - \lambda_{k-1}} & \cdots \\ \vdots & \vdots & 0 & \cdots & 0 & \cdots & 0 \\ \frac{A_{k1}}{\lambda_1 - \lambda_{k-1}} & \frac{A_{k2}}{\lambda_1 - \lambda_{k-1}} & 0 & \cdots & 0 & \cdots & 0 \\ \vdots & \vdots & 0 & \cdots & 0 & \cdots & 0 \end{pmatrix}.$$

The above is the correct form $\omega_1(\lambda)$, corresponding to the left upper block $\text{diag}(t_1, t_1) = t_1 I_2 \equiv \lambda_1 I_2$ of $\Lambda = \mathcal{U}|_{t_2=0}$. Analogous computations confirm that the \mathcal{V}_j are holomorphic in a neighbourhood of $t_2 = 0$ and

$$\mathcal{V}_j|_{t_2=0} = \left(\frac{\mathcal{V}_{rs}|_{t_2=0}(\delta_{rj} - \delta_{sj})}{u_r - u_s} \right)_{r,s=1}^n = \left(\frac{A_{rs}(\lambda) (\delta_{r-1,j-1} - \delta_{s-1,j-1})}{\lambda_{r-1} - \lambda_{s-1}} \right)_{r,s=1}^n, \quad j \geq 3.$$

which is the correct form of $\omega_{j-1}(\lambda)$ corresponding to the other 1-dimensional diagonal blocks of $\Lambda = \mathcal{U}|_{t_2=0}$.

• Next, we check if in (9.27) the correct term $\frac{\partial \mathcal{T}(\lambda)}{\partial \lambda_j} \cdot \mathcal{T}(\lambda)^{-1}$ appears, namely we prove the equalities (9.30), with $\mathcal{T}(\lambda) \equiv \mathcal{T}(t_1, u_3, \dots, u_n)$ having structure (9.29). Observe that in \mathcal{B}

$$\Psi^{-1} \frac{\partial \Psi}{\partial t_1} = \begin{pmatrix} \widehat{\Psi}^{-1} \frac{\partial \widehat{\Psi}}{\partial t_1} & 0 \cdots 0 \\ \vdots & \vdots \\ 0 & 0 & \mathbf{0} \\ \vdots & \vdots \end{pmatrix}, \quad \Psi^{-1} \frac{\partial \Psi}{\partial u_j} = \begin{pmatrix} \widehat{\Psi}^{-1} \frac{\partial \widehat{\Psi}}{\partial u_j} & 0 \cdots 0 \\ \vdots & \vdots \\ 0 & 0 & \mathbf{0} \\ \vdots & \vdots \end{pmatrix}, \quad j \geq 3, \quad (9.31)$$

The h_ℓ are a priori arbitrary functions, because they reduce the 1-dimensional blocks $A_{[\ell,\ell]} \equiv \mathcal{V}_{\ell+1,\ell+1}|_{t_2=0}$ to “diagonal form”! Corresponding to the $\mathbf{0}$ block in (9.31), equations (9.30) are satisfied if the h_ℓ are non-zero arbitrary constants. \mathcal{T}_1 must reduce the block $A_{[1,1]} = \mathcal{V}_{[1,1]}|_{t_2=0}$ in (9.20) to Jordan (diagonal, in our case) form. Thus, we have to solve for \mathcal{T}_1 the system

$$\begin{cases} \frac{\partial \mathcal{T}_1}{\partial t_1} \mathcal{T}_1^{-1} = -\widehat{\Psi}^{-1} \frac{\partial \widehat{\Psi}}{\partial t_1} |_{t_2=0} \\ \frac{\partial \mathcal{T}_1}{\partial u_j} \mathcal{T}_1^{-1} = -\widehat{\Psi}^{-1} \frac{\partial \widehat{\Psi}}{\partial u_j} |_{t_2=0}, \quad j \geq 3, \\ \mathcal{T}_1^{-1} \mathcal{V}_{[1,1]}|_{t_2=0} \mathcal{T}_1 = \begin{pmatrix} i\dot{V}_{12} & 0 \\ 0 & -i\dot{V}_{12} \end{pmatrix} \text{ “constraint”}. \end{cases} \quad (9.32)$$

The "known terms" $\widehat{\Psi}^{-1} \frac{\partial \widehat{\Psi}}{\partial t_1} \Big|_{t_2=0}$, $\widehat{\Psi}^{-1} \frac{\partial \widehat{\Psi}}{\partial u_j} \Big|_{t_2=0}$ in (9.32) all have the same structure. Indeed, let ξ be one of the variables t_1, u_3, \dots, u_n . Then, the corresponding "known term" is the value at $t_2 = 0$ of

$$\widehat{\Psi}^{-1} \frac{\partial \widehat{\Psi}}{\partial \xi} \stackrel{(9.16)}{=} \frac{1}{2(\tilde{\eta}_{12}^2 - t_2^{m-2} \tilde{\eta}_{11}^2)} \begin{pmatrix} \tilde{\eta}_{12} \frac{\partial \tilde{\eta}_{12}}{\partial \xi} - \tilde{\eta}_{11} \frac{\partial \tilde{\eta}_{11}}{\partial \xi} t_2^{m-2} & t_2^{m-2} \left(\tilde{\eta}_{12} \frac{\partial \tilde{\eta}_{11}}{\partial \xi} - \tilde{\eta}_{11} \frac{\partial \tilde{\eta}_{12}}{\partial \xi} \right) \\ \tilde{\eta}_{12} \frac{\partial \tilde{\eta}_{11}}{\partial \xi} - \tilde{\eta}_{11} \frac{\partial \tilde{\eta}_{12}}{\partial \xi} & \tilde{\eta}_{12} \frac{\partial \tilde{\eta}_{12}}{\partial \xi} - \tilde{\eta}_{11} \frac{\partial \tilde{\eta}_{11}}{\partial \xi} t_2^{m-2} \end{pmatrix}. \quad (9.33)$$

This is analytic in variables $(t_1, t_2, u_3, \dots, u_n)$ in a neighbourhood of $t_2 = 0$.

Case $m \geq 3$. In this case $\tilde{\eta}_{12} \neq 0$ (see Remark 9.1), so that

$$\widehat{\Psi}^{-1} \frac{\partial \widehat{\Psi}}{\partial \xi} \Big|_{t_2=0} = \frac{1}{2} \begin{pmatrix} \frac{1}{\tilde{\eta}_{12}} \frac{\partial \tilde{\eta}_{12}}{\partial \xi} & 0 \\ \left(\frac{1}{\tilde{\eta}_{12}} \frac{\partial \tilde{\eta}_{11}}{\partial \xi} - \frac{\tilde{\eta}_{11}}{\tilde{\eta}_{12}^2} \frac{\partial \tilde{\eta}_{12}}{\partial \xi} \right) & \frac{1}{\tilde{\eta}_{12}} \frac{\partial \tilde{\eta}_{12}}{\partial \xi} \end{pmatrix} \Big|_{t_2=0}.$$

The general solution of (9.32) is then

$$\mathcal{T}_1 = \begin{pmatrix} \frac{c_1}{\sqrt{\tilde{\eta}_{12}}} & 0 \\ -\frac{c_1}{2} \frac{\tilde{\eta}_{11}}{\tilde{\eta}_{12}^{3/2}} & \frac{c_2}{\sqrt{\tilde{\eta}_{12}}} \end{pmatrix} \Big|_{t_2=0}, \quad c_1, c_2 \in \mathbb{C} \setminus \{0\}. \quad (9.34)$$

Case $m = 2$. By Remark 9.3, $\mathcal{V}_{[1,1]} \Big|_{t_2=0} = 0$, so that we can take any solution of $\frac{\partial \mathcal{T}_1}{\partial \xi} \mathcal{T}_1^{-1} = -\widehat{\Psi}^{-1} \frac{\partial \widehat{\Psi}}{\partial \xi} \Big|_{t_2=0}$. The general solution is, for an arbitrary constant invertible matrix C ,

$$\begin{aligned} \mathcal{T}_1(t_1, u_3, \dots, u_n) &= \widehat{\Psi}^{-1}(t_1, 0, u_3, \dots, u_n) C \\ &\stackrel{(9.16), (9.17)}{=} \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -i \\ \frac{1}{\sqrt{\tilde{\eta}_{12} + \tilde{\eta}_{11}}} & \frac{-i}{\sqrt{\tilde{\eta}_{12} - \tilde{\eta}_{11}}} \\ 1 & i \\ \frac{1}{\sqrt{\tilde{\eta}_{12} + \tilde{\eta}_{11}}} & \frac{i}{\sqrt{\tilde{\eta}_{12} - \tilde{\eta}_{11}}} \end{pmatrix} \Big|_{t_2=0} \cdot C. \end{aligned}$$

Notice that if hypothetically $\mathring{V}_{12} \neq 0$, then the choice of \mathcal{T}_1 which diagonalizes $\mathcal{V}_{[1,1]} \Big|_{t_2=0}$ in (9.20) for $m = 2$ is

$$C = \begin{pmatrix} \alpha & \pm i\beta \\ \pm i\alpha & \beta \end{pmatrix} \implies \mathcal{T}_1 \cdot \mathcal{V}_{[1,1]}^{-1} \Big|_{t_2=0} \cdot \mathcal{T}_1 = \text{diag}(\pm i\mathring{V}_{12}, \mp i\mathring{V}_{12}).$$

□

Proof of point 3) of Remark 9.3. The coefficients of equations (9.22) and (9.25) are holomorphic at $t_2 = 0$ in variables $(t_1, t_2, u_3, \dots, u_n)$ if and only if so is the term

$$t_2^{\frac{m-2}{2}} \Psi^{-1} (V_1 - V_2) \Psi - \Psi^{-1} \frac{\partial \Psi}{\partial t_2}.$$

We have

$$\Psi^{-1}(V_1 - V_2)\Psi = \Psi^{-1} \begin{pmatrix} 0 & \frac{-2V_{12}}{u_2 - u_1} & \cdots & \frac{V_{1k}}{u_1 - u_k} & \cdots \\ \frac{2V_{21}}{u_1 - u_2} & 0 & \cdots & \frac{-V_{2k}}{u_2 - u_k} & \cdots \\ \vdots & \vdots & 0 & \cdots & 0 & \cdots & 0 \\ \frac{V_{k1}}{u_1 - u_k} & \frac{-V_{k2}}{u_2 - u_k} & 0 & \cdots & 0 & \cdots & 0 \\ \vdots & \vdots & 0 & \cdots & 0 & \cdots & 0 \end{pmatrix} \Psi$$

The analogous computations of the proof of Proposition 9.1 lead to

$$t_2^{\frac{m-2}{2}} \Psi^{-1}(V_1 - V_2)\Psi \underset{t_2 \rightarrow 0}{\sim} \begin{pmatrix} \mathcal{O}(1/t_2) & \mathcal{O}(t_2^{m-3}) & \cdots & \frac{A_{1k}}{\lambda_1 - \lambda_{k-1}} & \cdots \\ \mathcal{O}(1/t_2) & \mathcal{O}(1/t_2) & \cdots & \frac{-A_{2k}}{\lambda_1 - \lambda_{k-1}} & \cdots \\ \vdots & \vdots & 0 & \cdots & 0 & \cdots & 0 \\ \frac{A_{k1}}{\lambda_1 - \lambda_{k-1}} & \frac{-A_{k2}}{\lambda_1 - \lambda_{k-1}} & 0 & \cdots & 0 & \cdots & 0 \\ \vdots & \vdots & 0 & \cdots & 0 & \cdots & 0 \end{pmatrix}.$$

On the other hand, using (9.16) and (9.17) one finds

$$\Psi^{-1} \frac{\partial \Psi}{\partial t_2} = \begin{pmatrix} \widehat{\Psi}^{-1} \frac{\partial \widehat{\Psi}}{\partial t_2} & 0 \cdots 0 \\ \vdots & \vdots \\ 0 & 0 & \mathbf{0} \\ \vdots & \vdots \end{pmatrix} \underset{t_2 \rightarrow 0}{\sim} \begin{pmatrix} \mathcal{O}(1/t_2) & \mathcal{O}(t_2^{m-3}) & \cdots & 0 & \cdots \\ \mathcal{O}(1/t_2) & \mathcal{O}(1/t_2) & \cdots & 0 & \cdots \\ \vdots & \vdots & 0 & \cdots & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 & \cdots & 0 \\ \vdots & \vdots & 0 & \cdots & 0 & \cdots & 0 \end{pmatrix}.$$

All The \mathcal{O} terms are explicitly computed in terms of $\tilde{\eta}_{11}, \tilde{\eta}_{12}, V_{12}$. From this, one sees that in

$$t_2^{\frac{m-2}{2}} \Psi^{-1}(V_1 - V_2)\Psi - \Psi^{-1} \frac{\partial \Psi}{\partial t_2}$$

the divergences holomorphically cancel if and only if $\mathring{V}_{12} = i(m-2)/2m$, and in this case

$$t_2^{\frac{m-2}{2}} \Psi^{-1}(V_1 - V_2)\Psi - \Psi^{-1} \frac{\partial \Psi}{\partial t_2} \Big|_{t_2=0} = \begin{pmatrix} 0 & 0 & \cdots & \frac{A_{1k}}{\lambda_1 - \lambda_{k-1}} & \cdots \\ 0 & 0 & \cdots & \frac{-A_{2k}}{\lambda_1 - \lambda_{k-1}} & \cdots \\ \vdots & \vdots & 0 & \cdots & 0 & \cdots & 0 \\ \frac{A_{k1}}{\lambda_1 - \lambda_{k-1}} & \frac{-A_{k2}}{\lambda_1 - \lambda_{k-1}} & 0 & \cdots & 0 & \cdots & 0 \\ \vdots & \vdots & 0 & \cdots & 0 & \cdots & 0 \end{pmatrix}.$$

□

In conclusion, Proposition 9.1 shows that the isomonodromy deformation theory developed in this paper, summarized in Theorems 6.1 and 6.2, applies to the caustic of a Dubrovin-Frobenius manifold of the class studied in [28]. The representation of the Dubrovin flat connection on the basis $\partial/\partial t_1, \partial/\partial t_2, \pi_3, \dots, \pi_n$, and restricted to the caustic, is exactly a Pfaffian system of type (6.1), and the z -component is strongly isomonodromic (Remark 9.4). Moreover, the isomonodromy deformation theory developed in this paper allows us to predict some properties at the caustic, as Corollary 9.1 and Remark 9.3.

10 Appendix. Proof of Lemma 6.1

We consider the Jordan form $J = J_1 \oplus \cdots \oplus J_s$ of $A_{[1,1]} \oplus \cdots \oplus A_{[s,s]}$ and the corresponding $L = L_1 \oplus \cdots \oplus L_s$. We prove that if the deformation is strongly isomonodromic, then

$$\left[\mathcal{T}_k^{-1} \frac{\partial \mathcal{T}_k}{\partial \lambda_j}, L_k \right] = \left[\mathcal{T}_k^{-1} \frac{\partial \mathcal{T}_k}{\partial \lambda_j}, J_k \right] = 0, \quad \forall k = 1, \dots, s.$$

For a strong isomonodromy deformation, J and R , and equivalently D and L , are constant. The factor $z^D z^L$ in each

$$Y_\nu(z, \lambda) = \mathcal{T}(\lambda) \hat{Y}_\nu(z, \lambda) z^D z^L e^{z\Lambda}$$

corresponds to a fundamental solution in Levelt form for the system (4.1) of [13], which has a Fuchsian singularity in $z = \infty$. Up to a permutation, we can always assume that the matrices $Y_\nu(z, \lambda)$ are taken so that $z^D z^L$ satisfies the properties of Section 5 (to which we refer for notations).¹² Therefore, following Section 5, we can write

$$Y_\nu(z, \lambda) = \mathcal{T}(\lambda) \hat{Y}_\nu(z, \lambda) z^{D+\Sigma} z^N e^{z\Lambda}.$$

Step 1. First, we show that

$$[\mathcal{T}(\lambda)^{-1} d\mathcal{T}(\lambda), D + \Sigma] = 0, \tag{10.1}$$

¹²One can take a permutation matrix P , which does not change Λ because it permutes indexes inside the same block, which yields $\mathcal{T}P(P^{-1}\hat{Y}_\nu P)z^{P^{-1}DP}z^{P^{-1}LP}e^{z\Lambda}$ with the desired properties of Section 5.

and

$$[\mathcal{T}(\lambda)^{-1}d\mathcal{T}(\lambda), N] = 0. \quad (10.2)$$

In order to prove (10.1)-(10.2), recall (4.9)-(4.10) and let

$$\begin{aligned} \mathcal{Y}_\nu(z, \lambda) &:= Y_\nu(z, \lambda)\mathcal{T}(\lambda)^{-1}, & \mathfrak{D}(\lambda) &:= \mathcal{T}(\lambda)(D + \Sigma)\mathcal{T}(\lambda)^{-1} \\ \widehat{\mathcal{Y}}_\nu(z, \lambda) &:= \mathcal{T}(\lambda)\widehat{Y}_\nu(z, \lambda)\mathcal{T}(\lambda)^{-1}, & \mathfrak{N}(\lambda) &:= \mathcal{T}(\lambda)N\mathcal{T}(\lambda)^{-1} \end{aligned}$$

Then, we have

$$\begin{aligned} Y_\nu(z, \lambda) &= \widehat{\mathcal{Y}}_\nu(z, \lambda) z^{\mathfrak{D}(\lambda)} z^{\mathfrak{N}(\lambda)} e^{z\Lambda} \mathcal{T}(\lambda) = \mathcal{Y}_\nu(z, \lambda) \mathcal{T}(\lambda), \\ \widehat{\mathcal{Y}}_\nu(z, \lambda) &\sim I + \sum_{j=1}^{\infty} \mathcal{T}(\lambda) F_j(\lambda) \mathcal{T}(\lambda)^{-1} z^{-j}, \quad z \rightarrow \infty \text{ in } \mathcal{S}_\nu. \end{aligned}$$

Let d_λ be the differential w.r.t. $\lambda_1, \dots, \lambda_s$. From (6.5),

$$\sum_{j=1}^s \left(zE_{p_j} + \omega_j(\lambda) \right) d\lambda_j + d\mathcal{T} \cdot \mathcal{T}^{-1} = d_\lambda Y_\nu(z, \lambda) \cdot Y_\nu(z, \lambda)^{-1}$$

The right hand-side is

$$\begin{aligned} & d_\lambda \mathcal{Y}_\nu \cdot \mathcal{Y}_\nu^{-1} + \mathcal{Y}_\nu d\mathcal{T} \cdot \mathcal{T}^{-1} \mathcal{Y}_\nu^{-1} = \\ &= d_\lambda \widehat{\mathcal{Y}}_\nu \cdot \widehat{\mathcal{Y}}_\nu^{-1} + \widehat{\mathcal{Y}}_\nu \sum_{m=1}^{\infty} \frac{d(\mathfrak{D}^m)}{m!} (\ln z)^m z^{-\mathfrak{D}} \widehat{\mathcal{Y}}_\nu^{-1} + \widehat{\mathcal{Y}}_\nu z^{\mathfrak{D}} \sum_{k=1}^{\bar{k}} \frac{d(\mathfrak{N}^k)}{k!} (\ln z)^k z^{-\mathfrak{N}} z^{-\mathfrak{D}} \widehat{\mathcal{Y}}_\nu^{-1} + \\ & \quad + z \widehat{\mathcal{Y}}_\nu z^{\mathfrak{D}} z^{\mathfrak{N}} d\Lambda z^{-\mathfrak{N}} z^{-\mathfrak{D}} \widehat{\mathcal{Y}}_\nu^{-1} + \widehat{\mathcal{Y}}_\nu z^{\mathfrak{D}} z^{\mathfrak{N}} e^{\Lambda z} d\mathcal{T} \cdot \mathcal{T}^{-1} e^{-\Lambda z} z^{-\mathfrak{N}} z^{-\mathfrak{D}} \widehat{\mathcal{Y}}_\nu^{-1} \quad (10.3) \\ &= \underbrace{d_\lambda \widehat{\mathcal{Y}}_\nu \cdot \widehat{\mathcal{Y}}_\nu^{-1}}_{O(1/z)} + \widehat{\mathcal{Y}}_\nu \sum_{m=1}^{\infty} \frac{d(\mathfrak{D}^m)}{m!} (\ln z)^m z^{-\mathfrak{D}} \widehat{\mathcal{Y}}_\nu^{-1} + \widehat{\mathcal{Y}}_\nu z^{\mathfrak{D}} \sum_{k=1}^{\bar{k}} \frac{d(\mathfrak{N}^k)}{k!} (\ln z)^k z^{-\mathfrak{N}} z^{-\mathfrak{D}} \widehat{\mathcal{Y}}_\nu^{-1} + \\ & \quad + z \widehat{\mathcal{Y}}_\nu d\Lambda \widehat{\mathcal{Y}}_\nu^{-1} + \widehat{\mathcal{Y}}_\nu z^{\mathfrak{D}} z^{\mathfrak{N}} d\mathcal{T} \cdot \mathcal{T}^{-1} z^{-\mathfrak{N}} z^{-\mathfrak{D}} \widehat{\mathcal{Y}}_\nu^{-1}. \end{aligned}$$

In the last step, we have used the fact that $e^{\Lambda z}$ commutes with $d\mathcal{T} \cdot \mathcal{T}^{-1}$, \mathfrak{D} and \mathfrak{N} , due to the block structure. The absence of logarithmic singularities in $\sum_{j=1}^s (zE_{p_j} + \omega_j(\lambda)) d\lambda_j + d\mathcal{T} \cdot \mathcal{T}^{-1}$ requires that

$$d\mathfrak{D} = d\left(\mathcal{T}(\lambda)(D + \Sigma)\mathcal{T}(\lambda)^{-1}\right) = 0, \quad d\mathfrak{N} = d\left(\mathcal{T}(\lambda)N\mathcal{T}(\lambda)^{-1}\right) = 0 \quad (10.4)$$

Since $d(D + \Sigma) = dN = 0$ for the strong isomonodromy deformation, the above conditions are satisfied if and only if (10.1) and (10.2) respectively hold. In this way, also the last term in (10.3), namely

$$\widehat{\mathcal{Y}}_\nu z^{\mathfrak{D}} z^{\mathfrak{N}} d\mathcal{T} \cdot \mathcal{T}^{-1} z^{-\mathfrak{N}} z^{-\mathfrak{D}} \widehat{\mathcal{Y}}_\nu^{-1} \equiv \widehat{\mathcal{Y}}_\nu z^{D+\Sigma} z^N \mathcal{T}^{-1} d\mathcal{T} z^{-N} z^{-D-\Sigma} \widehat{\mathcal{Y}}_\nu^{-1},$$




Figure 2: Structure of the blocks L_k of L . Here, Σ_k is diagonal and N_k is nilpotent

does not contain logarithmic singularities when (10.1)-(10.2) hold, because it reduces to

$$\hat{Y}_\nu d\mathcal{T} \cdot \mathcal{T}^{-1} \hat{Y}_\nu^{-1} = d\mathcal{T} \cdot \mathcal{T}^{-1} + O(1/z).$$

Notice that the above behaviour is in agreement with $\sum_{j=1}^s (zE_{p_j} + \omega_j(\lambda))d\lambda_j + d\mathcal{T} \cdot \mathcal{T}^{-1}$.

Notice that if J is diagonal, $\mathcal{T}(\lambda)(D + \Sigma)\mathcal{T}(\lambda)^{-1} = \mathcal{T}(\lambda) J \mathcal{T}(\lambda)^{-1}$ in the first equation in (10.4)

Step 2. The relations (10.1)-(10.2) can be written for the individual blocks inherited from $L = L_1 \oplus \cdots \oplus L_s$ and $\mathcal{T} = \mathcal{T}_1 \oplus \cdots \oplus \mathcal{T}_s$. It suffices to consider a single block with label k .

As already explained, up to a $Y_\nu \mapsto Y_\nu P$ given by a suitable permutation matrix P , we assume that for each block L_k , $k = 1, \dots, s$, the Levelt structure explained in Section 5 applies. See figure 2.

We consider the problem at the level of a single block with label k . In order to avoid a proliferation of indices, from now on $\mathcal{J}, \mathcal{L}, \mathcal{R}, \mathcal{S}, \mathcal{D}, \Sigma$ and \mathcal{N} will respectively stand for $J_k, L_k, R_k, S_k, D_k, \Sigma_k$ and N_k . We will take the label k only for \mathcal{T}_k , in order not to confuse it with the full \mathcal{T} . To them, the structures of Section 5 apply. Now, we have $\mathcal{L} = \mathcal{L}_1 \oplus \cdots \oplus \mathcal{L}_\ell$ for some ℓ , and

$$\Sigma = \sigma_1 I_1 \oplus \cdots \oplus \sigma_\ell I_\ell,$$

with eigenvalues σ_q (with real part in $[0, 1)$). Hence, since \mathcal{D} is diagonal, (10.1) for the block k of $\mathcal{T}^{-1}d\mathcal{T}$ gives

$$[\mathcal{T}_k^{-1}d\mathcal{T}_k, \mathcal{D} + \Sigma] = 0 \implies \mathcal{T}_k = \mathcal{T}_1^{(k)} \oplus \cdots \oplus \mathcal{T}_\ell^{(k)}.$$

Now, $\mathcal{D} = \mathcal{D}_1 \oplus \cdots \oplus \mathcal{D}_\ell$, and $\mathcal{N} = \mathcal{N}_1 \oplus \cdots \oplus \mathcal{N}_\ell$. Notice that

$$[(\mathcal{T}_q^{(k)})^{-1}d\mathcal{T}_q^{(k)}, \mathcal{D}_q + \Sigma_q] = 0. \quad (10.5)$$

Each \mathcal{N}_q ($q = 1, \dots, \ell$) is upper triangular, it has zeros on the diagonal, and its diagonal blocks are elementary Jordan sub-blocks with 1's on the second upper diagonal, as in figure 3. Accordingly,




Figure 3: The structure of the sub-blocks \mathcal{N}_q and \mathcal{D}_q of \mathcal{N} and \mathcal{D} (that is, of a certain N_k and D_k), corresponding to $\Sigma_q = \sigma_q I_q$.

\mathcal{D}_q , which is diagonal with a non decreasing sequence of integer eigenvalues, has sub-blocks with the same eigenvalue corresponding to a Jordan sub-block in \mathcal{N}_q , as in figure 3.

The above facts and (10.5) imply that $(\mathcal{T}_q^{(k)})^{-1} d\mathcal{T}_q^{(k)}$ is divided into sub-blocks as $\Sigma_q + \mathcal{D}_q$, where the only non-zero sub-blocks are the block-diagonal part, as in Figure 4 .

It follows that $[\mathcal{T}_k(\lambda)^{-1} d\mathcal{T}_k(\lambda), \mathcal{N}] = 0$ (from (10.2)) reduces to

$$[(\mathcal{T}_q^{(k)})^{-1} d\mathcal{T}_q^{(k)}, \mathcal{N}_q] = 0, \quad (10.6)$$

Let $\mathcal{M}_q := (\mathcal{T}_q^{(k)})^{-1} d\mathcal{T}_q^{(k)}$, and let $\mathcal{M}_{[a,b]}^{(q)} = \mathcal{M}_{[a,b]}^{(q)} \delta_{ab}$ be a sub-block. From (10.6) we receive

$$\mathcal{M}_{[a,a]}^{(q)} \mathcal{N}_{[a,b]}^{(q)} - \mathcal{N}_{[a,b]}^{(q)} \mathcal{M}_{[b,b]}^{(q)} = 0.$$

In particular

$$\mathcal{M}_{[a,a]}^{(q)} \mathcal{N}_{[a,a]}^{(q)} - \mathcal{N}_{[a,a]}^{(q)} \mathcal{M}_{[a,a]}^{(q)} = 0,$$

which means that \mathcal{M}_q commutes with the block-diagonal matrix $\mathcal{N}_{[1,1]}^{(q)} \oplus \mathcal{N}_{[2,2]}^{(q)} \oplus \dots \oplus \mathcal{N}_{[\ell,\ell]}^{(q)}$ (= Jordan matrix obtained from the diagonal sub-blocks of \mathcal{N}_q in the left part of figure 3).

Now, observe that $\mathcal{D}_q + \Sigma_q + (\mathcal{N}_{[1,1]}^{(q)} \oplus \mathcal{N}_{[2,2]}^{(q)} \oplus \dots) = \mathcal{J}_q$. Since \mathcal{M}_q also commutes with $\mathcal{D}_q + \Sigma_q$, we conclude that it commutes with \mathcal{J}_q , namely

$$[(\mathcal{T}_q^{(k)})^{-1} d\mathcal{T}_q^{(k)}, \mathcal{J}_q] = 0.$$

Therefore,

$$[(\mathcal{T}_k)^{-1} d\mathcal{T}_k, \mathcal{J}] = 0.$$

Coming back to the original notations, the above is $[(\mathcal{T}_k)^{-1} d\mathcal{T}_k, J_k] = 0$. This is what we wanted to prove. \square




Figure 4: The sub-blocks-structure of the block $(\mathcal{T}_q^{(k)})^{-1} d\mathcal{T}_q^{(k)}$.

References

- [1] Anosov D. V., Bolibruch A. A., The Riemann-Hilbert Problem, A Publication of the Steklov Institute of Mathematics, Vieweg (1994).
- [2] Arnol'd, V. I.: On matrices depending on parameters, Russian Math. Surveys 26 , no. 2, (1971), 29-43.
- [3] Bertola, M., Mo, M. Y.: Isomonodromic deformation of resonant rational connections. IMRP Int. Math. Res. Pap. 2005, no. 11, 565-635.
- [4] Boalch, P.: G-bundles, isomonodromy, and quantum Weyl groups. Int. Math. Res. Not. 2002, no. 22, 1129-1166.
- [5] Boalch, P.: From Klein to Painlevé via Fourier, Laplace and Jimbo. Proc. London Math. Soc. (3) 90 (2005), 167-208.
- [6] Boalch, P.: Simply-laced isomonodromy systems. Publ. Math. Inst. Hautes Études Sci. 116 (2012), 1-68.
- [7] Boalch, P.: Geometry and braiding of Stokes data; fission and wild character varieties. Ann. of Math. (2) 179 (2014), 301-365.
66-69.
- [8] Bolibruch A. A., The fundamental matrix of a Pfaffian system of Fuchs type. Izv. Akad. Nauk SSSR Ser. Mat. 41 (1977), no. 5, 1084-1109, 1200.
- [9] Bolibruch A. A., On Isomonodromic Deformations of Fuchsian Systems, Journ. of Dynamical and Control Systems, 3, (1997), 589-604.
- [10] Bolibruch A. A., On Isomonodromic Confluence of Fuchsian Singularities, Proc, Stek. Inst. Math. 221, (1998), 117-132.
- [11] T. Bridgeland, V. Toledano Laredo: *Stokes factors and Multilogarithms*. J. reine und angew. Math. **682** (2013), 89-128.

- [12] Cotti G., Guzzetti D., Results on the Extension of Isomonodromy Deformations to the case of a Resonant Irregular Singularity, *Random Matrices Theory Appl.*, 7 (2018) 1840003, 27 pp.
- [13] Cotti G., Dubrovin B., Guzzetti D., Isomonodromy Deformations at an Irregular Singularity with Coalescing Eigenvalues, *Duke Math. J.*, (2019), arXiv:1706.04808 (2017).
- [14] Cotti G., Dubrovin B., Guzzetti D., Local Moduli of Semisimple Frobenius Coalescent Structures, *SIGMA* 16 (2020), 040, 105 pages. (2018).
- [15] Degano G., Guzzetti D., The sixth Painlevé equation as isomonodromy deformation of an irregular system: monodromy data, coalescing eigenvalues, locally holomorphic transcendents and Frobenius manifolds, arXiv:2108.07003 (2021).
- [16] Dubrovin B., Geometry of 2D topological field theories, *Lecture Notes in Math*, 1620, (1996), 120-348.
- [17] Dubrovin B., Painlevé transcendents in two-dimensional topological field theory, in “The Painlevé Property, One Century later” edited by R.Conte, Springer (1999).
- [18] Guzzetti D., Inverse problem and monodromy data for three-dimensional Frobenius manifolds. *Math. Phys. Anal. Geom.* 4 (2001), no. 3, 245-291.
- [19] Guzzetti D.: Notes on Non-Generic Isomonodromy Deformations, *SIGMA* 14 (2018), 087, 34 pages
- [20] Guzzetti D.: Deformations with a resonant irregular singularity, in Formal and analytic solutions of diff. equations, Springer Proc. Math. Stat., 256, (2018), 239-250.
- [21] Guzzetti D.: Isomonodromy deformations with coalescing eigenvalues and applications, in Complex Differential and Difference Equations, De Gruyter Proceedings in Mathematics (2020), 313-325.
- [22] Guzzetti, D.: Isomonodromic Laplace Transform with Coalescing Eigenvalues and Confluence of Fuchsian Singularities, *Letters in Mathematical Physics* (2021) 111:80, <https://doi.org/10.1007/s11005-021-01423-z>, arXiv:2101.03397.
- [23] Harnad, J.: Dual isomonodromic deformations and moment maps to loop algebras. *Comm. Math. Phys.* 166 (1994), 337-365.
- [24] Hertling, C.: *Frobenius Manifolds and Moduli Spaces for Singularities*, Cambridge University Press (2002)
- [25] Jimbo, M.; Miwa, T.; Mori, Y.; Sato, M.:Density matrix of an impenetrable Bose gas and the fifth Painlevé transcendent. *Phys. D* 1 (1980), no. 1, 80-158.
- [26] Jimbo M., Miwa T., Ueno K., Monodromy Preserving Deformations of Linear Ordinary Differential Equations with Rational Coefficients (I), *Physica, D2*, (1981), 306.
- [27] Mazzocco M.: Painlevé sixth equation as isomonodromic deformations equation of an irregular system, *CRM Proc. Lecture Notes*, **32**, Amer. Math. Soc. 219-238, (2002).
- [28] Reyes F.: Isomonodromic Deformations along the Caustic of a Dubrovin-Frobenius Manifold, arXiv:2209.01062, (2022).
- [29] Sabbah C., Integrable deformations and degenerations of some irregular singularities, *Publ. RIMS Kyoto Univ.* **57**, n. 3-4, (2021). arXiv:1804.00426 (2017)
- [30] Sibuya Y., Simplification of a System of Linear Ordinary Differential Equations about a Singular Point, *Funkcial. Ekvac*, 4, (1962), 29-56.
- [31] Strachan, I. A. B.: Frobenius submanifolds. *J. Geom. Phys.* 38, no. 3-4, (2001), 285-307.

- [32] Strachan, I. A. B.: Frobenius manifolds: natural submanifolds and induced bi-Hamiltonian structures. *Differential Geom. Appl.* 20, no. 1, (2004), 67-99.
- [33] Wasow W., *Asymptotic Expansions for Ordinary Differential Equations*, Dover (1965).
- [34] Yoshida M., Takano K., On a linear system of Pfaffian equations with regular singular points, *Funkcial. Ekvac.*, 19 (1976), no. 2, 175-189.